

Active detergent solubilized H⁺,K⁺-ATPase is a monomer*

Ingrid Dach^{1,2,3}, Claus Olesen^{1,3*}, Luca Signor^{4,5,6}, Poul Nissen^{1,2}, Marc le Maire^{7,8,9*}, Jesper V. Møller^{1,3*}, and Christine Ebel^{4,5,6*}

¹ From the Centre for Membrane Pumps in Cells and Diseases, Danish Research Foundation, Denmark,

² Department of Molecular Biology and Genetics, University of Aarhus, Denmark,

³ Department of Biomedicine, University of Aarhus, Denmark,

⁴ CEA, Institut de Biologie Structurale Jean-Pierre Ebel, Grenoble, France,

⁵ CNRS, Institut de Biologie Structurale, Grenoble, France,

⁶ Université Joseph Fourier Grenoble 1, Institut de Biologie Structurale, Grenoble, France,

⁷ CEA, iBiTec-S, Saclay, France,

⁸ CNRS, UMR 8221, France,

⁹ Université Paris-Sud, France.

*Running Title : *H⁺,K⁺-ATPase is a monomer*

*To whom correspondence should be addressed : Christine Ebel, Institut de Biologie Structurale Jean-Pierre Ebel, CEA, CNRS, and Université Joseph Fourier Grenoble 1, 41 rue Jules Horowitz, 38027 Grenoble Cedex 1, France, Tel.: (33) 4 38 78 95 70; Fax.: (33) 4 38 78 54 94; E-mail: christine.ebel@ibs.fr. Marc le Maire, DSV/iBiTec-S/SB2SM et UMR 8221 CEA, CNRS, Université Paris-Sud, PC 103, CEA de Saclay, F-91191 Gif sur Yvette Cedex, France ; Tel. : (33) 1 69 08 62 43 ; E-mail : marc.lemaire@cea.fr. Jesper V. Møller, Department of Biomedicine, University of Aarhus, Denmark Tel (45) 871 67754 ; E-mail : jvm@biophys.au.dk. Claus Olesen, Department of Biomedicine, University of Aarhus, Denmark Tel (45) 871 67753 ; E-mail : co@biophys.au.dk.

Keywords : P-ATPase, association state, purification, solubilization, detergent, activity

Background : The proton pump H⁺,K⁺-ATPase is responsible for the acidity of the gastric fluid. The minimum organization required for its activity is not established.

Results : We defined a solubilization protocol allowing us to obtain pig gastric H⁺, K⁺-ATPase as an active species.

Conclusion : Active detergent solubilized H⁺,K⁺-ATPase is unambiguously an α,β monomeric protomer.

Significance : The H⁺,K⁺-ATPase functioning is most probably similar to other P-ATPases

SUMMARY

The H⁺,K⁺-ATPase pumps protons or hydronium ions and is responsible for the acidification of the gastric fluid. It is made up of an α -catalytic- and a β -glycosylated subunit. The relation between cation translocation and the organization of the protein in the membrane are not well understood. We describe here how pure and functionally active pig gastric H⁺,K⁺-ATPase with an apparent

Stokes radius of 6.3 nm can be obtained after solubilization with the non-ionic detergent C₁₂E₈, followed by exchange of C₁₂E₈ with Tween 20 on a Superose 6 column. Mass spectroscopy indicates the β -subunit bears an excess mass of 9 Da attributable to glycosylation. From chemical analysis, there are 0.25 g phospholipids and around 0.024 g cholesterol bound per g protein. Analytical ultracentrifugation shows one main complex, sedimenting at $s_{20w} = 7.2 \pm 0.1$ S, together with minor amounts of irreversibly aggregated material. From these data a buoyant molecular mass is calculated, corresponding to an H⁺,K⁺-ATPase α,β -protomer of 147.3 kDa. Complementary sedimentation velocity with deuterated water gives a picture of an α,β -protomer with 0.9-1.4 g/g bound detergent and lipids and a reasonable frictional ratio of 1.5 corresponding to $R_s = 7.1$ nm. An α_2,β_2 dimer is rejected by the data. Light scattering coupled to gel filtration confirms the monomeric state of

solubilized H⁺,K⁺-ATPase. Thus, α,β H⁺,K⁺-ATPase is active at least in detergent, and may plausibly function as a monomer, as has been established for other P-type ATPases, Ca²⁺-ATPase and Na⁺,K⁺-ATPase.

The H⁺,K⁺-ATPase present in gastric mucosa is a P-type ATPase (1) capable of pumping protons or hydronium ions against very high concentration gradients ($\sim 10^6 : 1$) by an electroneutral K⁺ exchange mechanism (2-3). Like Na⁺,K⁺-ATPase, with which it shares high sequence similarity, H⁺,K⁺-ATPase is made up of an α -catalytic- and a β -glycosylated subunit. It is responsible for the acidification of the gastric fluid, with important pathophysiological consequences in cases of gastric hypersecretion. Extensive investigations on H⁺,K⁺-ATPase have led to the successful development of drugs of the omeprazole type, and of K⁺ competitive blockers like the Schering compound SCH28080, to inhibit H⁺,K⁺-ATPase *in vivo* (4-6). However, despite previous efforts, some fundamental aspects of the structure and function of H⁺,K⁺-ATPase are still not well understood, especially in relation to those of other P-type ATPases. This includes the relation between cation translocation and the organization of the protein in the membrane. Although analysis of ordered 2D crystals indicates overall structural features of H⁺,K⁺-ATPase similar to those of Na⁺,K⁺-ATPase and sarcoplasmic reticulum Ca²⁺-ATPase (7-9), other studies on H⁺,K⁺-ATPase have indicated that retention of enzyme activity in the detergent solubilized state depends on the enzyme being present in a α_2,β_2 dimer or higher oligomeric state of the α,β -protomer (10-18). The role of an obligate oligomeric organization of functional importance is underscored by functional evidence for half-of-the sites reactivity and other observations in the membraneous state (19, 18). Altogether these findings are in marked contrast to those emerging earlier on sarcoplasmic reticulum Ca²⁺-ATPase, in particular from studies on detergent solubilization (20-22), reconstitution in a surplus of phospholipids (23), and detailed studies of the kinetics and ligand stoichiometry (24, 21). Despite the probable presence of protein-protein interactions in the native membrane, which may affect enzyme activity (25-26), they all indicate monomeric Ca²⁺-ATPase as the fundamental Ca²⁺

transporting unit. For Na⁺,K⁺-ATPase, the evidence is more equivocal in that both monomeric and dimeric forms with retention of enzyme activity have been detected by analytical ultracentrifugation (27-30) and by HPLC (31-33). The requirement of an oligomeric structure as a *sine qua non* to preserve transport and enzyme activity as claimed for H⁺,K⁺-ATPase and reported in a recent review (34) is not indicated by either the detergent data or by the available 3D X-ray diffraction structures of Ca²⁺-ATPase (35-36), Na⁺,K⁺-ATPase (37-39), and H⁺-ATPase (40-41).

As a preliminary to attempts to crystallize H⁺,K⁺-ATPase for 3D structural analysis we have therefore felt the need to clarify the situation specifically for this transport protein. Could there be differences in the way by which H⁺,K⁺-ATPase is organized and works in the membrane, compared to that of other P-type ATPases? One possibility is that such differences could reflect relatively strong interactions between H⁺,K⁺-ATPase α,β -protomers that are retained after detergent solubilization. Furthermore, differences could be related to differences in transport mechanism due to the particular status of protons as the transported species. A dimeric state, for instance, might facilitate transfer through the membrane, either as the hydronium ion, H₃O⁺ by the Grothuis effect, or as H⁺ by a protein wire mechanism (42). In order to address these questions we have in the present communication attempted to apply our experience on the use of surfactants for solubilization of functionally active membrane proteins to H⁺,K⁺-ATPase in conjunction with structural studies by analytical ultracentrifugation and other advanced biophysical techniques (43). This has enabled us to provide a stringent analysis of the aggregational state of H⁺,K⁺-ATPase required to sustain activity.

EXPERIMENTAL PROCEDURES

H⁺,K⁺-ATPase preparation - Tubulovesicular preparations of gastric H⁺,K⁺-ATPase were prepared by differential centrifugation of gastric mucosa of actively secreting (pink) specimens of pig stomachs, with slight modifications of the method described by Skrabanja *et al.* (44). The pig stomachs were obtained from a local slaughterhouse (Danish Crown, Horsens) and kept cold after slaughtering and in all subsequent steps. After transfer to the laboratory, gastric mucosa

was scraped off from the stomach with a filet knife, minced twice with a garlic press, and homogenized at 1000 rpm with a home made motorized and Teflon piston based homogenizer in a buffer containing 0.25 M sucrose and 20 mM Tris-HCl (pH 7.4). The homogenate was centrifuged at 27,000 x g (15,000 rpm) for 20 min in a Sorvall SS-34 rotor and, after removal of the pellet, recentrifuged at 205,000 x g (46,000 rpm) in a Beckman Ti-45 angle rotor. The resulting pellet was resuspended with the buffer used for homogenization and layered on a two-step gradient consisting of 37 % sucrose (w/v) and 20 mM Tris-HCl (pH 7.4) at the bottom with 7 % Ficoll at the top (GE Healthcare, Denmark), 0.25 M sucrose and 20 mM Tris-HCl (pH 7.4). After centrifugation for 60 min at 217,000 x g (46,000 rpm) in a Beckman Ti.-70 swing-out rotor, H⁺,K⁺-ATPase containing vesicles were collected from the top of the Ficoll layer. As the last step the vesicles were diluted with 20 mM Tris buffer (pH 7.4) and pelleted at 217,000 x g followed by resuspension in the homogenization buffer at a final concentration of approx. 5 - 7 mg protein/ml, and kept frozen in small aliquots at -20°C.

Chromatographic preparation of active, solubilized H⁺,K⁺-ATPase - An aliquot of the vesicular H⁺,K⁺-ATPase membranes was spun down by centrifugation at 50,000 rpm for 35 min, and the pellet (approx 0.7 mg protein), resuspended with 0.7 ml 40 mM KCl, 40 mM Mes buffer (pH 6.0), 3 mM Mg²⁺, 1 mM EDTA, and 0.25 M sucrose. The membranes were solubilized by addition of C₁₂E₈ at a 5:1 detergent/protein weight ratio. After removal of insoluble residues by centrifugation at 50,000 rpm for 35 min, the supernatant, which had a content of 1.1 mg phospholipid/mg protein, was applied to a Superose 6 10x300 mm column, pre-equilibrated with the same buffer, except for the presence of Tween 20 (2 mg/ml) instead of C₁₂E₈. The sample was eluted at the rate of 0.25 ml/min with continuous monitoring of light absorption at 280 nm and refractive index.

Biochemical analyses - Protein concentrations were determined by the method of Lowry (45) as modified by Hartree (46), and phospholipid by the method of Bartlett (47). ATPase activities of the vesicular membranes were measured at 23°C by enzymatic spectrophotometry with an ATP regenerating system (48) in a 30 mM imidazole

(pH 7.0) buffer, containing, 5 mM Mg²⁺, 0.1 mM EDTA and 1 mM ATP to which had been added 0.42 µg alameticin/ µg protein to make the vesicles leaky to the substrates. Detergent solubilized ATPase activity was assayed in the column chromatographic solvent including the detergent.

Mass spectroscopy (MS) experiments - Membrane solubilization and purification of H⁺,K⁺-ATPase was done as described above, but in the presence of n-dodecyl-β-D-maltopyranoside (DDM) at 1 mg/ml. Three 500 µl fractions at 0.24, 0.27 and 0.35 mg/ml, eluting in the first part of the H⁺,K⁺-ATPase peak after the size exclusion chromatography, were concentrated to 15-20 µl and the DDM concentration was reduced by two cycles of dilution and concentration by ultrafiltration (Centricon cut off 100 kDa Amicon) in 40 mM Tris, 40 mM KCl, 0.2 mg/ml DDM. Previously, Lenoir *et al.* (49) and Cardi *et al.* (50) indeed have indicated that low salt and low DDM are appropriate for MALDI experiments. MALDI TOF MS spectra were measured on a Autoflex mass spectrometer (Bruker Daltonics) operated in linear positive ion mode. External calibration of the instrument was done using protein standard mixture II from Bruker Daltonics (BSA: 66.5 kDa, protein A: 44.6 kDa and trypsinogen: 23.9 kDa) providing usual mass accuracy ≤ 1000 ppm in this mass range. The samples were mixed in a ratio 1:2 (vol:vol) with sinapinic acid matrix (Sigma; 10 mg/ml in acetonitrile/water-0.1% TFA (50:50)) and 2 µl was deposited on the target. LC-ESI TOF MS analysis was carried out on a 6210 mass spectrometer interfaced with a binary pump liquid chromatography system (Agilent Technologies). The instrument was calibrated in the 300-3000 *m/z* range using ESI tuning mix from Agilent Technologies allowing a mass accuracy of about 50 ppm. All solvents were HPLC grade. Solvent A was 0.03% TFA in water and solvent B was acetonitrile 95%, TFA 0.03 %. Samples were desalted and washed on-line on a reverse phase cartridge (C4, Michrom Bioresources) with solvent A (2 min) and with 50 % solvent B (10 min.), at a flow rate of 0.3 ml/min. The sample was then separated and analysed on a reverse phase HPLC column (Jupiter-C4; 5µ, 300 Å, 50x1.0 mm, Phenomenex) with a linear gradient from 5 to 95 % B in 15 min.

Analytical ultracentrifugation sedimentation velocity (SV)¹ experiments - Sedimentation velocity experiments were done on an analytical ultracentrifuge XLI (Beckman Coulter, Palo Alto, USA) with a rotor speed of 42,000 rpm at 7 °C, using a rotor Anti-50 and double-sector cells (in Ti, Alu or epon) of optical path length 12 mm with sapphire windows. Cells were typically filled with 420 µl of sample and reference solvent. In some of the experiments, we used artificial boundary cells allowing the equilibration at 3,000 rpm of the levels of the two compartments prior stopping the ultracentrifuge and re-homogenisation of the sample prior the SV experiments: initial volumes of the channels were 430/420 µl for reference solvent and sample. The reference solvent was without detergent. Acquisitions were made using absorbance at 280 nm and interference optics. Sedimentation velocity analyses were performed by employing the Sedfit program, version 12.1, (<http://www.analyticalultracentrifugation.com>) in terms of continuous distributions $c(s)$ of sedimentation coefficients (s) (51). The solvent density and viscosity were measured at 20°C with a DMA 5000 density meter and AMVn viscosity meter (Paar, Graz, Austria), and extrapolated at 7°C using tabulated data for water. For experiments in deuterated buffer, three fractions of 0.9 or 1.3 ml of H^+,K^+ -ATPase from size exclusion chromatography in Tween 20 were recovered, and solvent exchanged to deuterated buffer (92.6% D₂O) of the same composition using desalting columns of 5 ml (PD10, Amersham). The theoretical background is detailed in (52, 43, 53) and in the text. For the analysis of the sedimentation in deuterated solvent, H-D exchange of exchangeable hydrogens in the protein by deuterium leads to a reformulation of the buoyant molar mass:

$$M_b = \sum M (M_D/M - \rho \bar{v}), \quad (1)$$

the summation standing for the decomposition of the particle in terms of its components, with M and \bar{v} being defined for the fully hydrogenated components.

Size exclusion chromatography coupled to light scattering (SEC-LS) experiments - H^+,K^+ -ATPase (30 µl) was injected into a high pressure liquid chromatography system LC20 AD (Shimadzu) in a KW 804 column (8 x 300 mm)

preceded by a guard column (Shodex) and equilibrated at 4°C, with a flow rate of 0.5 ml/min, with 40 mM KCl, 40 mM MES pH6, 0.25 M sucrose, 3 mM MgCl₂, 1mM EDTA, 2 mg/ml Tween 20, the solvent being filtered at 0.1 µm. The elution profile was followed on-line by refractive index (Δn) measurements with an Optilab rEX (Wyatt Technology), absorbance at 280 nm (A_{280}) with a Shimadzu SPD-M20A, and light scattering intensity (I) with a miniDAWN TREOS detector (Wyatt Technology) using laser emitting at $\lambda = 658$ nm. Data were analyzed using the ASTRA software (Wyatt Technology Corp.). The basic equation for light scattering, restricted to diluted solutions (corresponding to negligible interparticle interactions between particles) of small particles ($R_s < 20$ nm) reduces to:

$$I = K c M (\partial n / \partial c)^2, \quad (2)$$

where I is the excess intensity of scattered light, K is an optical parameter $= 4\pi^2 n^2 / (\lambda^4 N_A)$ and n is the refractive index. For a solute with known $(\partial n / \partial c)$, the concentration c is estimated from the refractive index measurements, $\Delta n = k c (\partial n / \partial c)$ with k being an instrumental constant.

For a complex described as a two component system (with indexes labelled 1 and 2 for the components), the concentrations of the component within the complex -thus its composition- are estimated from the combination of the absorbance, $A_{1cm, complex}$, and refractive index, $\Delta n_{complex}$, using as inputs the extinction coefficients $E_{0.1\%, 1}$ and $(\partial n / \partial c)$ for the two components:

$$\Delta n_{complex} = c_1 k (\partial n / \partial c)_1 + c_2 k (\partial n / \partial c)_2 \quad (3)$$

$$A_{1cm, complex} = c_1 E_{0.1\%, 1} + c_2 E_{0.1\%, 2} \quad (4)$$

The derived refractive index increment and concentration of the complex is combined with scattered light intensity for the calculation of its molecular mass in Eq. 2, where c , M and $(\partial n / \partial c)$ stand for the complex: $c = c_1 + c_2$; $M = M_1 + M_2$; $(\partial n / \partial c) = (\partial n / \partial c_1) c_1 / (c_1 + c_2) + (\partial n / \partial c_2) c_2 / (c_1 + c_2)$.

Numerical values used for the analysis of SV and SEC-LS experiments - We extrapolate to 7°C, from measurements done at 20°C, solvent densities for 40 mM KCl, 40 mM MES pH 6, 0.25 M sucrose, 3 mM MgCl₂, 1 mM EDTA in H₂O

and D_2O , of 1.035 and 1.126 g/ml, respectively, and solvent viscosities of 1.83 and 2.24 cP, respectively. The numerical values for the components of the H^+,K^+ -ATPase complex are reported in Table 1. Corrected $s_{20,w}$ values were calculated considering $\bar{v} = 0.83$ ml/g for the H^+,K^+ -ATPase complex and 0.87 ml/g for the Tween 20 or (Tween 20 + lipid) micelles.

RESULTS

Surfactant solubilization of tubular vesicular membranes and preparation of enzymatically active H^+,K^+ -ATPase. We prepared, as described under *Methods*, tubulovesicular fractions with a high H^+,K^+ -ATPase content by differential centrifugation of mucosal scrapings from actively acid secreting mucosae of pig stomachs. The use of surfactants to extract and solubilize membrane proteins in their molecular entities from the membrane matrix is indispensable for studying in detail structural aspects such as the role of monomeric/oligomeric organization, protein-protein interactions, for the functional properties of the membrane protein. In this way indeed, it was established that sarcoplasmic reticulum Ca^{2+} -ATPase, solubilized as a monomer by $C_{12}E_8$, retains all the basic features required to function as a Ca^{2+} transporting unit, regardless of the tendency of this Ca^{2+} pump to undergo reversible and irreversible self-association (20). For Na^+,K^+ -ATPase, this also appears to be the case (28-30), although dimerization of the α,β -protomer is a more pronounced feature of this ATPase than it is for Ca^{2+} -ATPase (27, 31-33). Application of the same surfactant strategy to H^+,K^+ -ATPase is fraught with the problem that it is difficult to avoid either irreversible or reversible inhibition of enzyme activity in the detergent solubilized state. Nevertheless, it has been found that for H^+,K^+ -ATPase inactivation can be reduced by partial solubilization with mild non-ionic surfactant under conditions that included cryoprotection with sucrose or glycerol, and a low ($4^\circ C$) temperature (see e.g. (15)). A key point for our study was the need to provide efficient solubilization by a potentially abrasive surfactant, to ensure formation of the minimal size of the H^+,K^+ -ATPase complex, with retention of function. After a number of preliminary trials with various non-ionic detergents (see Fig. S1) we found that these requirements were best met by the combined use

of $C_{12}E_8$ to initially solubilize the H^+,K^+ -ATPase membranes followed by exchange with the mild detergent Tween 20 to preserve activity in a following chromatographic separation. As can be seen from Fig. 1A, chromatography on Superose 6 resulted in the elution of three main peaks. In the first peak eluting at the void volume (V_0) there is some highly aggregated H^+,K^+ -ATPase. There was also non-solubilized phospholipid eluted at this position because of their reduced solubility in Tween 20 when compared to $C_{12}E_8$ (54). This conclusion can be drawn from the evidently smaller concentration of phospholipid present in the void volume for the sample shown in Fig. 1B where phospholipid was kept solubilized by $C_{12}E_8$ in the column eluant. With Tween 20 present in the eluant (Fig. 1A) the peak following the void volume has a high content of enzymatically active and essentially pure H^+,K^+ -ATPase that elutes partially delipidated, with 0.25 mg of phospholipids per mg protein. The third peak is mainly constituted by phospholipids that elute as mixed micelles with $C_{12}E_8$ (and probably also with some Tween 20) together with protein components of lower molecular mass present in the tubulovesicular preparation (Fig. 1E). Cholesterol was mainly detected in the third peak. However, there was also cholesterol in the protein peak, amounting to around 0.024 mg/mg protein (data not shown). The H^+,K^+ -ATPase present in the second peak (Fig. 1A) exhibited ATPase activity at a level similar to that of the membraneous preparation (see Fig S1C). The activity only slowly deteriorated over a period of 2 days when the column collected fractions were stored in the cold ($4^\circ C$) in the presence of 0.3 M sucrose (Fig. 1C). In contrast to these findings, when $C_{12}E_8$ was used instead of Tween 20 as the column eluant after initial solubilization with $C_{12}E_8$ (Fig. 1B), there was a large increase of protein in the void volume, and the H^+,K^+ -ATPase in the second peak was inactive, delipidated, and with an elution profile clearly revealing size heterogeneity. From column calibration data with water soluble standard proteins (Fig. 1D) we could deduce an apparent value for the Stokes radius of 6.3 nm for the enzymatically active H^+,K^+ -ATPase in the second peak in Fig. 1A.

The present protocol is similar to those previously used for preparation of $C_{12}E_8$ solubilized Ca^{2+} -ATPase, followed by exchange

with a milder detergent like Tween 80 (22) or an Amphipol polymer (55). In all of these cases, $C_{12}E_8$ has been used to break up the native membrane structure by partially solubilizing the membrane lipid, followed by exchange with a milder and less abrasive detergent of the Tween series (22) or by Amphipol treatment (55) to keep the ATPase soluble in an active form.

Molecular mass determination of the two subunits by MS. By MS we have sought to evaluate the effect of post-translational modifications, in particular on the carbohydrate content of our H^+,K^+ -ATPase preparations. The theoretical protein molecular masses deduced from the aminoacid sequences are 114,287 Da for the α -subunit and 33,076 Da for the β -subunit. These values are listed in Table 1 together with numerical values of the other components present in the solubilized H^+,K^+ -ATPase complex. Analysis by MS of H^+,K^+ -ATPase in Tween 20 at 2 mg/ml did not allow identification of the H^+,K^+ -ATPase polypeptide peaks: no protein peaks were detected due to the presence of this detergent. However, as shown in Fig. 2, it was possible to obtain clearly resolved mass spectra, albeit with low signal over noise ratios, for H^+,K^+ -ATPase prepared by solubilization in DDM, and then chromatographed at a column concentration of DDM of 0.2 mg/ml before concentration on Centricon 100 (Amicon). Analysis performed on two different samples by MALDI TOF resulted in registration of molecular masses up to 42,488 Da attributable to the β -subunit, and 114,781 Da for the α -subunit. For the α -subunit the molecular mass exceeds the theoretical value by 494 Da; we cannot exclude that at least part of this slight difference can be related to instrument inaccuracy, arising because the calibration was done with protein standards of lower mass in the 20-70 kDa range. Thus, LC-ESI TOF analysis gave slightly lower values, viz. 41,662 Da for the β -subunit and 114,112 Da for the α -subunit. Later in the LC elution, other contributions at 38,349 and 39,255 Da were detected. The origin of the latter two minor contributions is uncertain. We presume that they either represent protein unrelated to the β -subunit or that they correspond to β -subunits with different glycosylation levels; or perhaps to derivatives formed by partial proteolysis of the β -subunit during the preparation.

For the α -subunit, the molecular mass obtained from LC-ESI TOF is lower by 175 Da when compared to the theoretical mass value, and in this case a lower molecular mass is in accordance with known post-translational modifications, resulting in the removal of the N-terminal methionine (56) and with oxidation of some of the cysteine residues. Thus, we can conclude that the molecular masses measured by MALDI TOF and LC-ESI TOF for the α -subunit closely agrees with that computed from the aminoacid sequence. On the other hand, the molecular masses of the β -subunit as measured by the two ionization techniques are substantially higher than that calculated from the deduced aminoacid sequence (by 9346 Da in MALDI TOF and by 8586 Da in LC-ESI TOF). Therefore this subunit, according to our MS data on pig stomach H^+,K^+ -ATPase, includes a contribution from the glycosylated side chains that, based on a calculated protein molecular mass of 33076, is about 9 kDa corresponding to 0.06 g/g protein. This value is, in fact, lower than the value of 20 kDa glycosylation that we have been able to calculate from the model put forward by Tyargarajan *et al.* for rabbit gastric H^+,K^+ -ATPase (57) (see the Discussion).

Sedimentation velocity experiments after size exclusion chromatography - To study the size of the H^+,K^+ -ATPase, solubilized by Tween 20 in an enzymatically active form (Fig. 1A) we used analytical ultracentrifugation, which is a versatile and powerful method with which to assess sample homogeneity, and to characterize the monomeric/oligomeric states of proteins and their association equilibria (58). Previous experience has indicated that SV is to be preferred to sedimentation equilibrium in the study of the often inherently unstable detergent solubilized membrane proteins (43). With SV it is possible to follow in detail the evolution of sedimentation profiles by repeating the scans with short time intervals. This obviates the long time periods required for concluding the sedimentation equilibrium experiments (43, 53). Thus secondary aggregation of the detergent solubilized membrane protein can be avoided, or at least checked, since the development of size heterogeneity is very easily revealed by repeated sedimentation velocity scans. SV can be used in conjunction with continuous size distribution analysis for accurate

determination of sedimentation coefficients, s , for the different species present in a complex sample (51). The s -values are related to the molecular mass, M , the Stokes radius, R_s , and the buoyancy term, $(1 - \rho \bar{v})$, of the sedimenting species by the Svedberg relation, which for the present purpose can be formulated in the following way:

$$s = M(1 - \rho \bar{v}) / (N_A 6 \pi \eta R_s) \quad (5)$$

In this equation, N_A is Avogadro's number; ρ and η are the solvent density and viscosity and \bar{v} is the partial specific volume of the sedimenting species, which in our case includes bound lipid and detergent by the glycosylated protein. Because M and \bar{v} cannot be deconvoluted from SV measured at only one solvent density, we instead used the buoyant molar mass defined in Eq. 1 as the experimentally relevant parameter (see the results section and Eqs 7-9). The frictional ratio ff_{\min} that relates R_s to the radius, R_{\min} , of the anhydrous volume is determined by its shape and hydration:

$$ff_{\min} = R_s / R_{\min}. \quad (6)$$

In the $c(s)$ analysis, reasonable hypotheses on the values of \bar{v} , ff_{\min} , ρ and η are defined (ff_{\min} is fitted) allowing a plausible relationship between the buoyant mass and size of the particles. Furthermore, as will be shown, it is possible by combining SV data obtained with different optics (absorbance and interference optics), or by recording the sedimentation in solvents with different densities, to determine all the molecular parameters accessible to analytical ultracentrifugation. We have therefore based our study on the use of SV rather than on sedimentation equilibrium measurements, although the latter theoretically is more directly related to the determination of molecular mass.

In our experiments, SV analysis was performed at 7°C on samples prepared as in Fig. 1A by solubilization of H⁺,K⁺-ATPase in C₁₂E₈, followed by size exclusion chromatography on a Superose 6 column in the presence of Tween 20 and sucrose in the cold room. Three fractions, eluting from the column at different positions of the enzymatically active protein peak as indicated by the arabic numerals 1, 2, and 3 on Fig. 1A were

examined. Fig. 3A shows the UV absorption sedimentation profiles of the central top fraction 2, while Fig. 3B shows interference data obtained in the same experiment. The sedimentation profiles of Fig. 3A indicate the presence of one major boundary of the protein, while Fig. 3B obtained by interference optics shows the presence of one additional and slower moving component, which is attributable to the sedimentation of the detergent micelles. Global analysis of the entire SV scans as outlined above was done by subjecting the data to computer simulation to provide a continuous $c(s)$ distribution of sedimentation coefficients (51). Fig. 3C compares the properties of the three studied fractions. When investigated at 280 nm, they all show one main complex sedimenting at 3.23 +/- 0.03 S (corresponding to $s_{20w} = 7.2 +/- 0.1$ S). Other ill-defined and larger species are present and sediment at 5 +/- 0.2 S ($s_{20w} = 11.1 +/- 0.4$ S) and further up to 7 S ($s_{20w} = 15.5$ S). We found that the presence of the larger species was most prominent in samples collected from the first eluted fractions of the protein peak. These larger species appear as irreversibly aggregated material, since the s -values and proportions of the different species were not modified when the samples were diluted twice (data not shown). Thus, there is no evidence for the existence of any reversible equilibrium between the differently sized species. Furthermore, we noted in introductory SV experiments, which we had performed at the more commonly used rotor temperature of 20°C, that the proportion of the main $s_{20w} = 7.2$ S species was drastically reduced (more than 50%), and replaced by a predominant third species sedimenting at $s_{20w} = 15$ S. A higher proportion of the larger species at $s_{20w} = 11.1$ and 15 S was also observed in SV experiments of samples that had been stored for 18 hours at 7°C after chromatography, or for samples of H⁺,K⁺-ATPase obtained after an additional size exclusion chromatography column (using a silica column KW804 as described in the section on light scattering). Accordingly, the larger species most likely represent a time and temperature dependent deterioration of the sample that is paralleled by the decline of enzyme activity as a function of time (Fig. 1). The ratios of the sedimentation coefficients of the larger species to that of the main species are 1.5 and 1.9 for species sedimenting at $s_{20w} = 11.1$ and 15 S, respectively. This is close to the theoretical values of 1.59 and

2.08 expected for sedimentation of dimers and trimers of the same density and shape as the main species. The increased trend for aggregated ATPase to elute early from the column is apparent from Fig. 3D. In conclusion, the larger species probably correspond to irreversibly aggregated oligomeric forms (dimers, trimers and larger multimers) of the main species related to aging, a process which occurs more rapidly at 20°C than at 7°C.

The profiles obtained by interference optics (Fig. 3B) were analysed as depicted by the broken lines in Fig. 3C. They showed in addition to the protein species detected at 280 nm significant contributions of a boundary, sedimenting at 0.55 +/- 0.1 S ($s_{20w} = 1.3 +/- 0.2$ S), attributable to Tween micelles (see the physical chemical characterisation of Tween 20 in supplemental data Fig. S2). The contributions seen at lower s -values ($s < 0.3$ S) in Fig. 3C may be artefactual and were not analysed further (they may be solvent components or result from improper deconvolution of systematic noises). In addition, there were ill-defined contributions sedimenting between 1 and 2.7 S (Fig. 3C) with a mean value of 1.7 +/- 0.1 S ($s_{20w} = 4.1 +/- 0.2$ S) which we consider represents the presence of the mixed lipid/detergent micelles that are observed to elute at the trailing edge of the enzymatically active peak of H⁺,K⁺-ATPase (see Fig. 1A). In accordance with this view, the concentration of these species was larger in the protein fractions eluted late in the size exclusion chromatography (Fig. 3C). Furthermore, the J/A_{280} ratio, i.e. the ratio of the fringe shift to absorbance of these components at 280 nm was very large (>50), indicative of a predominant detergent and lipid composition, not compatible with a pure protein-detergent complex.

Information on the composition of the ATPase complex could also be gained from the interference data by comparison of the values for the absorbance at 280 nm (A_{280}) and fringe shift (J) signals (see the Methods section). As deduced from the $c(s)$ analysis, the J/A_{280} ratio for the main H⁺,K⁺-ATPase complex was 5.4 +/- 0.6 and 6.3 +/- 2.2 for the dimer species. These values are much larger than the values of 2.68 for the deglycosylated - and 2.81 for the glycosylated protein, which can be calculated for the detergent and lipid-free H⁺,K⁺-ATPase based on the values

for refractive index increment ($\partial n/\partial c$) and $E_{0.1\%}$ extinction coefficient at 280 nm given in Table 1. The excess of fringe shift observed for the solubilized complex is obviously related to the binding of appreciable amounts of detergent and lipid, which on this basis can be estimated to be present in a total amount of about 1.6 gram per gram of protein, considering 0.274 g/g bound lipid.

Hydrodynamic characterization based on combining the s and R_s - values - By the preceding analysis we have demonstrated that the main component of a Tween 20 solubilized and enzymatically active H⁺,K⁺-ATPase is present as a slowly moving species, sedimenting at $s_{20,w} = 7.2$ S. This is a low value when compared with previously reported values for sedimentation of related P-type ATPases, e.g. $s_{20,w} = 10.6$ S for the DDM solubilized monomer of α,β Na⁺,K⁺-ATPase (30) and $s_{20,w} = 7.5$ S for the sarcoplasmic reticulum Ca²⁺-ATPase, which has no β -subunit (43). We checked the consistency of our H⁺,K⁺-ATPase data by estimating the buoyant molecular mass of the enzymatically active peak by combining the measured sedimentation coefficient obtained in the sucrose containing solvent (3.23 S) with the Stokes radius, $R_s = 6.3$ nm, as determined from calibrated size exclusion chromatography. This can be done by use of the following equation (36):

$$M_b = 6 \pi N_a \eta R_s s \quad (7)$$

which with a solvent viscosity, $\eta = 1.83$ cP results in an estimate of the buoyant molar mass (M_b) of the main species of 42.3 kDa. M_b is related to the molecular mass (M) and partial specific volume (\bar{v}) of the sedimenting species by:

$$M_b = M (1 - \rho \bar{v}), \quad (8)$$

where ρ is the solvent density, which is 1.035 g/ml in our experimental conditions. For the H⁺,K⁺-ATPase complex, M_b in the above expression is related to the molecular mass of the protein component, M_p , and the sum of the buoyancies of the protein (P)-, glycosyl (G)-, lipid (L)-, and detergent (D) bound components by the following relation

$$M_b = M_p [(1 - \rho \bar{v}_p) + (1 - \rho \bar{v}_G) \delta_G +$$

$$(1 - \rho \bar{v}_L) \delta_L + (1 - \rho \bar{v}_D) \delta_D] \quad (9)$$

where δ_G is the mass of the glycosyl, δ_L of the lipid, and δ_D of the detergent associated components of the H^+,K^+ -ATPase protein, expressed on a gram/gram basis. From these equations and using the numerical values given in Table 1, we calculate that, for a H^+,K^+ -ATPase α,β -protomer, the protein content (with a molar mass of 147.3 kDa) contributes with 35.3 kDa and glycosylation (9 kDa) with 3.5 kDa to M_b . Bound phospholipid and cholesterol (with a global molar mass of 40.3 kDa) makes essentially no contribution (-0.1 kDa), because $1/\bar{v}_L$ is close to ρ . The sum of these contributions in terms of buoyant mass, 38.7 kDa, for a H^+,K^+ -ATPase α,β -protomer thus approaches that estimated according to Eq. 7 experimentally for the buoyancy of the solubilized H^+,K^+ -ATPase (42.3 kDa), except for a small residual (3.6 kDa) that may be attributed to bound Tween 20. The data are thus consistent with the elution of the Tween 20 solubilized H^+,K^+ -ATPase as an unaggregated α,β -protomer with a moderate amount of bound detergent (of the order 0.2 – 0.3 g detergent per g protein). This is a lower value for detergent binding than suggested from the J/A_{280} analysis given above. However, it should be noted that the calculation of detergent binding from the buoyancy terms in Eq. 9 is subject to uncertainty, being based on the difference between two large numbers (42.3 kDa – 38.7 kDa). In particular the calculation is critically dependent on the correctness of the value for the column based determination of Stokes radius, where it is known that the R_S values of membrane proteins, as compared to those of the watersoluble standard proteins used for column calibrations, can be influenced by the type of membrane protein and by the detergent in the column eluent (see (43)).

Sedimentation velocity experiments in deuterated solvent - In order to proceed with the hydrodynamic analysis and arrive at a firm conclusion as to the aggregational state and other hydrodynamic parameters of the solubilized H^+,K^+ -ATPase, we have made supplementary determinations of the sedimentation coefficients after exchange of the solvent in our samples with deuterated water. Previously, introducing density variation by exchange with heavy water has been established in sedimentation equilibrium

experiments as a useful method to estimate the contribution that the partial specific volumes makes to M_b , the buoyant molar mass of the solubilized complex (59). In appropriate cases this method can be used, by detergent density matching, to blank out the effect of detergent from the buoyancy term in sedimentation equilibrium experiments of membrane proteins (60). Recently, D_2O exchange has also been used to extend the scope for $c(s)$ sedimentation velocity analysis, as a means to unravel the hydrodynamic properties of samples that are not homogeneous or stable (61-62), and to deal with the complexity of solubilized membrane proteins (63).

In our experiments H^+,K^+ -ATPase exchange with deuterated solvent was done by subjecting different fractions of the enzymatically active H^+,K^+ -ATPase to solvent exchange on a PD10 column. The $c(s)$ analysis shows that for three D_2O/H_2O exchanged fractions, there is, as in hydrogenated solvent, one main peak and some aggregates, mainly dimers of the main complex, that as expected are less prominent in the later eluted fractions (shown in Fig. S3 in the supplemental data). The s -values were logically reduced when compared to those obtained in hydrogenated solvent, due to the increase in density of the deuterated solvent (rising from 1.035 to 1.124 g/ml) and of the viscosity (rising from 1.83 cP to 2.24 cP). For the main peak, we determined $s = 1.6 \pm 0.1$ S, while the sedimentation rate for the larger particles were in the range 2.3 - 5 S. The analysis by interference could not be performed here in a satisfactory way, because the modelling was made difficult by the presence of Tween 20, which floats at -0.005 S under these conditions. Analysis of the sedimentation of the protein was done according to:

$$s_H \eta_H / s_D \eta_D = (1 - \bar{v} \rho_H) / (M_D/M_H - \bar{v} \rho_D) \quad (10)$$

where subscripts H and D refer to the relevant values in hydrogenated and deuterated solvents for the sedimentation coefficients, viscosities, densities, and molecular masses. Note that the denominator on the right hand side incorporates a correction for the increase in molecular mass of the protein complex as a result of the D/H exchange. With these data and considering the

increase in molecular mass, M_D/M_H , caused by D/H exchange in the H^+,K^+ -ATPase complex, to be 1.014 (intermediate between that of the protein and the Tween 20 / lipid component, see Table 1), we calculate from Eq. 10 a value for the partial specific volume of the complex of 0.818 +/- 0.015 ml/g (errors being evaluated from the imprecision on s_H and s_D). For a α,β -protomer of H^+,K^+ -ATPase binding the amounts of carbohydrate, phospholipid and cholesterol shown in Table 1, we calculate for this ensemble, a molecular mass of 196.7 kDa and a partial specific volume of 0.779 ml/g. By the use of Eq. 9 this can be used to derive an estimate of the amount of bound Tween 20 of 1.0 +/-0.7 g/g. The minimum radius (R_{min}) that can be calculated from this particle composition is 4.8 +/- 0.4 nm, which also by the use of Eq. 5 leads to a revised value for the Stokes radius (R_S) of 7.9 +/- 1.3 nm ($fff_{min}=1.6 +/- 0.2$).

From the above analysis in D_2O/H_2O , we have thus obtained a set of solutions for all of the hydrodynamic parameters of the detergent solubilized H^+,K^+ -ATPase. This includes an estimate of the amount of bound Tween 20, arising as a side gain from the determination of the buoyancy of the whole complex. In order to explore further the range of values consistent with our data we have followed a procedure described by Nury *et al.* (63). According to this the detergent solubilized complex is considered to be comprised of two pseudo-components: (i) the glycosylated protein and (ii) the detergent + lipid component. In this approach the lipid and detergent are thus lumped together and the analysis considers all possible solutions for δ_{D+L} , for different hypotheses of association states of the protein (monomer, dimer, etc), and frictional ratios. The frictional ratio ($fff_{min} = R_S/R_{min}$) is a shape factor that includes a contribution from hydration, while R_{min} is defined as the minimum radius of the anhydrous volume (defined by the product $M\bar{v}$). For proteins the magnitude of fff_{min} is restricted to a rather limited range. Values of 1.15-1.25 define globular and compact proteins (64). Coil-like disordered proteins have fff_{min} in the range 1.5 to 3 depending on their masses (64). Typical values for glycolysated proteins are 1.5-1.8. From the Svedberg equation (Eq. 1), and incorporating a correction for the slight increase of mass of the components due to H-D exchange in deuterated solvents, the following equation is derived:

$$s = M_{PG} [((M_D/M_H)_{PG} - \rho \bar{v}_{PG}) + \delta_{D+L}(M_D/M_H)_{D+L} - \rho \bar{v}_{D+L}] / [6\pi N_a \eta fff_{min} ((3/4\pi N_a)(M_{PG} \bar{v}_{PG} + \delta_{D+L} \bar{v}_{D+L}))^{1/3}] \quad (11)$$

The equation should be satisfied in different solvents (with defined values of s , $(M_D/M_H)s$, ρ and η). If the oligomeric state (M_{PG}) and shape factor (fff_{min}) are appropriately chosen, a unique pair of (δ_{D+L} , \bar{v}_{D+L}) values must account for sedimentation in H_2O and D_2O buffers. We used the commercial computer algebra system *Maple* to draw the implicit function $\delta_{D+L} = f(\bar{v}_{D+L})$, based on different assumptions as to the magnitude of fff_{min} and aggregational state. We considered at the outset that H^+, K^+ -ATPase is a monomer or a dimer and assume different values of the frictional ratio. Fig. 4 shows the results for four hypotheses: H^+, K^+ -ATPase as a monomer with $fff_{min}=1.3$ (Panel A), or as a monomer with $fff_{min}=1.5$ (Panel B) or $fff_{min}=1.8$ (Panel C); and as a dimer with $fff_{min}=1.5$ (Panel D). On each panel, four curves $\delta_{D+L} = f(\bar{v}_{D+L})$ are superimposed, representing the limiting values of the experimental sedimentation coefficients ($s \pm \Delta s$) in the hydrogenated and deuterated solvents, thus a total of four s -values. The areas delimited by the four curves (coloured in grey in Fig. 4A-C) define all possible mathematical solutions for $\delta_{D+L} = f(\bar{v}_{D+L})$ that are compatible with the experimental data. Among them, we exclude some irrelevant ones, since \bar{v}_{D+L} has to be within the range defined by pure Tween 20 (0.87 ml/g) and lipid (0.98 ml/g). Real solutions for our ATPase will be further restricted to \bar{v}_{D+L} within an even narrower range, since we know that both components will be present in the actual complex. We find that with these constraints, the s_H/s_D data are definitively not compatible with a globular compact monomer ($fff_{min}=1.25$, not shown). Fig. 4A therefore presents the analysis for the minimal $fff_{min}=1.3$ value compatible with our data, which includes \bar{v}_{D+L} values that are just above 0.96 ml/g -i.e. mainly lipids -, with δ_{D+L} being in the range 0.55-0.85 g/g, and $R_S=5.7$ nm. The largest (mathematically found) anisotropy of $fff_{min}=1.8$ (Fig. 4C) would correspond to 1.7 +/- 0.3 g/g of bound detergent + lipids (with $\bar{v}_{D+L} = 0.878 +/- 0.005$ ml/g -i.e.

this, the present data offer evidence for the view that the detergent solubilized H^+,K^+ -ATPase α,β -protomer is the fundamental unit for H^+,K^+ -ATPase function. In addition to the strong homology to Na^+,K^+ -ATPase and Ca^{2+} -ATPase, combined with many functional data, this is a suggestion which is supported by recent progress on the structure of H^+,K^+ -ATPase, obtained on 2D crystals of this protein with a resolution of 6.5 Å (8-9, 65). These structural data suggest that the H^+,K^+ -ATPase α,β -protomer with a few structural adaptations operates on the same general structural principles as Ca^{2+} -ATPase (35-36) and Na^+,K^+ -ATPase (37-39). The differences include an intraprotomeric protein-protein interaction of the β -unit with the phosphorylation domain of the α -subunit, for H^+,K^+ -ATPase. It would account for the "ratchet" mechanism, proposed as a special mechanism that prevents reversal of the proton transport mechanism (9). Such a mechanism, by preventing the E2P conformation from becoming reversibly dephosphorylated by backward reaction with ADP via E1P (65), is considered necessary to explain the very high proton membrane gradients ($\sim 10^6 : 1$) achievable by ATP phosphorylation with H^+,K^+ -ATPase (66), E1P and E2P being the two main conformations of phosphorylated P-ATPases, see *e.g.* (36). Unique features of H^+,K^+ -ATPase also comprise a modified E1P conformation, approaching that of E2P, and lack of evidence for wide opening of the intramembranous helices towards the extracytosolic side in E2P ground state in the BeF stabilized crystals (65), in contrast to the luminal opening observed for the same kind of crystals in Ca^{2+} -ATPase (67). These differences can probably be attributed to the particular status of the proton which leaves this ion with more possibilities for a facile traverse through the H^+,K^+ -ATPase membrane by hydronium ion and/or proton wiring mechanisms (37).

However, it should be noted that since the new data do not tell us much about the actual state of self-association of H^+,K^+ -ATPase in the native membrane, they do not exclude either the possibility that quaternary organization could be of regulatory importance for H^+ transport through membranes as has been inferred from kinetic experiments suggesting half-of-the-sites reactivity (18-19). The main argument for half-of-the-sites reactivity of a membraneous H^+,K^+ -ATPase

revolves around the issue of the maximal amount that can be obtained by phosphorylation with ATP in K^+ free and hence dephosphorylation-blocked medium. The value obtained is around 2.5 nmol EP/mg protein (see *e.g.* (18)). Does this value represent 50 % of that which would be possible for a monoprotomeric and fully active protein (*i.e.* approx 5 nmol/mg for H^+,K^+ -ATPase)? An affirmative answer to this question presupposes the availability of fully purified and active preparations of the enzyme. However, it is common experience within the P-type ATPase field that even the best "pure" preparations of *e.g.* Ca^{2+} -ATPase and Na^+,K^+ -ATPase contain considerable amounts of inactive ATPase (see *e.g.* (36, 68-69)). For these two enzymes the level of inactive ATPase is close to 50% in the native membranes. Reliable stoichiometries are also made difficult by other methodological uncertainties such as the determination of absolutely correct protein concentrations and upper values for ligand binding with low affinity. For instance the error in protein determination using BSA as a standard for P-ATPase or other membrane proteins is of the order of 20% (see *e.g.* (70-71)). The incompletely active proteins and methodological difficulties are the two factors that may explain the extremely large variability of results for stoichiometries obtained by different authors for describing the properties of ATPases, *e.g.* Na^+,K^+ -ATPase (72).

So far the use of detergents to study the properties of the gastric acid pump has been hampered by the fact that H^+,K^+ -ATPase, in comparison to other P-type ATPase cation pumps, is exquisitely sensitive to deterioration by exposure to detergents. This prevented us to employ detergents ordinarily used for this purpose to prepare solubilized H^+,K^+ -ATPase in a structurally defined and functional state. We here introduced Tween 20 as a new detergent for keeping the ATPase in a solubilized and functionally active state. This was also the case for DDM (14) which could have been the only viable alternative for performing these experiments, but the enzyme was prepared with much less ATPase activity after HPLC (See Fig. S1C). Tween 20 is a sorbitol based polyoxyethylene glycol detergent that is chemically heterogeneous (73). It is being used extensively as a membrane transferring and surface blocking agent in *e.g.* Western blot and

ELISA experiments, but so far is rather inefficient a solubilizer of biological membranes (54). This is probably due to the presence of a long and branched hydrophilic chain leading to a high HLB (Hydrophilic-Lipophilic Balance) number. Despite this, we had noticed in previous experiments that Ca^{2+} -ATPase could be slowly solubilized as monomers from sarcoplasmic reticulum membranes by prolonged (overnight) treatment with Tween 20 (unpublished data). However, proper solubilization of gastric H^+,K^+ -ATPase membrane vesicles could not be done by treatment with Tween 20 alone (Fig. S1A). In order to achieve solubilization we used an initial treatment with $C_{12}E_8$ to break up the membrane structure and partially delipidate the H^+,K^+ -ATPase and combined this step with a quick exchange of $C_{12}E_8$ by Tween 20 to avoid inactivation of the enzyme. The fractionated separation on a Superose column indeed allowed us to maintain the H^+,K^+ -ATPase in a solubilized and enzymatically active state (Fig. 1). This two-stage strategy is similar to that previously employed to keep Ca^{2+} -ATPase in a solubilized and enzymatically active state with Tween 80 (22) and Amphipol (74) after pretreatment with $C_{12}E_8$.

With this preparation we determined by chemical analysis that lipids with our procedure were retained at a level of 0.25 g/g for phospholipids, and 0.024 g/g for cholesterol. Analysis of the phospholipid content of the tubulovesicular membranes indicated a usual overall fatty acid distribution, but with a high content of phosphatidylethanolamine (53%) (Table S1). We determined also a rather low cholesterol content (7.5 % (w/w) of total phospholipids). We consider that all lipid eluting together with H^+,K^+ -ATPase by gel filtration is actually bound by the protein. To retain some lipid on the ATPase is important for the preservation of protein activity after solubilization, although too much will be expected to favour oligomerisation (75). In addition, it is known that a suitable amount of lipid will be required to facilitate membrane proteins crystallisation -see *e.g.* (76-77).

As a second important point we determined the glycosylation of the pig H^+,K^+ -ATPase β -subunit to contribute with 9 kDa to the mass of the H^+,K^+ -ATPase β -subunit as determined by MS of the complete protein (Fig. 2). This value is significantly lower than the glycosylation level

that we could estimate from the model for the β -subunit by Tyagarajan et al. (57) based on rabbit gastric H^+,K^+ -ATPase. From this model we calculate a glycosylation mass of 20 kDa considering the masses of the main glycopeptides analyzed by MS analysis. Glycosylation heterogeneity was observed in the rabbit on all the N-glycosylation sites (an additional O-glycosylation site is also mentioned but not characterized). The probable presence of microheterogeneity in the glycosylation pattern for pig gastric H^+,K^+ -ATPase studied here is suggested from the smear observed on SDS PAGE, while an inordinately high apparent molecular mass, resulting in an electrophoretic migration close to that of the α -subunit, could suggest that the β -subunit is only slowly released from the α,β complex during the SDS-PAGE procedure (Fig. 1E). The large difference in the glycosylation level estimated between pig and rabbit gastric H^+,K^+ -ATPase could also arise for a number of other reasons. First it is known that protein glycosylation is a highly regulated process that strongly depends on species, tissue localisation, and cell metabolism (see *e.g.* (78-79), and references therein). In addition, we estimate a reduced number of potential glycosylation sites in the pig: 6 sites, including a marginal one, from the NetNGlyc 1.0 Server program (www.cbs.dtu.dk/services/NetNGlyc) to be compared to seven N-glycosylation sites in H^+,K^+ -ATPase from rabbit. Furthermore, we cannot exclude either that partial deglycosylation has occurred during purification or that there has been incomplete glycosylation during the transit from the endoplasmic reticulum to the plasma membrane. It is interesting to note that while glycosylation of H^+,K^+ -ATPase and Na^+,K^+ -ATPase is important for targeting of these proteins in polarized cells, their exact role in folding, stabilisation and transport function is still a debated issue (see (80) and references therein).

The determination of the association state of membrane proteins by hydrodynamic methods is not a simple procedure, because solubilized membrane proteins usually are multicomponent complexes (43, 53). This may explain part of the controversy concerning different proteins (see *e.g.* (81, 63)), including H^+,K^+ -ATPase, whose association state was inferred from a variety of techniques (reviewed in (16)), for example native

gel electrophoresis (14-15), gel filtration, see *e.g.* (16), or cross-linking (13). To these difficulties is added sample instability in the detergent solubilized state. It may explain the structural heterogeneity noticed for H⁺,K⁺-ATPase by chromatography, or in single-molecule detection using total internal reflection fluorescence microscopy (16-17) which has given rise to extreme difficulty in assigning which association state is the functional one. Our choice of sedimentation velocity allowed us to conduct the experiments within a time frame sufficiently short not to significantly perturb aggregation and other molecular properties as well as enzyme activity. A main species was identified, together with minor amounts of aggregates that clearly were attributable to sample aging. Our solubilized H⁺,K⁺-ATPase in addition to glycosylation and lipid contains bound Tween 20 for which we had no specific analytical procedure. Tween 20 bound to H⁺,K⁺-ATPase was first inferred from refractive index and UV absorption measurements, with suitable deductions for the changes of these parameters caused by the lipid and protein content (Table 1). This resulted in an estimate of 1.6 g/g the precision of which however is strongly dependent on the exactness of the extinction coefficients (Table 1). Combining the values of the sedimentation coefficient and of the Stokes radius (estimated with undefined uncertainty to be 6.3 nm by the elution from calibrated Superose columns) did not allow to define the detergent content, but strongly suggested the sedimenting H⁺,K⁺-ATPase species to be in a monomeric α,β state. Furthermore, we were able to estimate the partial specific volume of the solubilized complex from the decrease in $s_{20,w}$ following H₂O/D₂O solvent exchange. From the determined value of 0.813 ml/g and integrating this with our knowledge of

the carbohydrate and lipid content, we can estimate the binding of Tween 20 to be 1.0 +/- 0.7 g/g. In view of its sedimentation, a particle with this composition would correspond to a molecular dimension slightly larger than estimated from gel filtration. Analysis of the sedimentation in H₂O and D₂O, in the absence of any hypothesis on the lipid content allowed us to confirm the description the solubilized H⁺,K⁺-ATPase as a compact α,β glycosylated monomer, with a frictional ratio of 1.5 +/- 0.1 and 0.9-1.4 g/g bound detergent plus lipids. Finally, independent measurements, employing gel filtration coupled to light scattering, absorbance and refractive index detections supported the view of an α,β monomer solvated with 1.1 +/-0.2 g/g bound detergent plus lipids, based on the extinction coefficients calculated from sequence.

In conclusion, by our combined approach a consistent picture is emerging from these measurements. In fact, we consider the present work as a showpiece how biophysical methods can be developed and adapted to answer specific questions of both structural and functional importance for membrane proteins. By these means we have defined a solubilization protocol allowing to obtain solubilized pig gastric H⁺, K⁺-ATPase as an active species over days, which was unambiguously present as a α,β monomeric protomer. We would like to point out that, by using similar experimental approaches, dimers were observed in solution, for membrane proteins described to function as dimers in the membrane, for example, cytochrome C oxydase (81) or ABC transporters (82-83). We conclude that the general functioning of H⁺,K⁺-ATPase is most probably similar to that of the Ca²⁺-ATPase and Na⁺, K⁺-ATPase.

References :

1. Pedersen, P. L., and Carafoli, E. (1987) Ion motive ATPases. 1. Ubiquity, properties, and significance to cell-function. *Trends Biochem.Sci.* **12**, 146-150.
2. Sachs, G., Munson, K., Balaji, V. N., Aures-Fischer, D., Hersey, S. J., and Hall, K. (1989) Functional domains of the gastric HK ATPase. *J. Bioenerg. Biomembr.* **21**, 573-588.
3. Läuger, P. (1991) *Electrogenic ion pumps*, Sinauer Associates, Sunderland, Mass.
4. Sachs, G., Shin, J. M., Vagin, O., Lambrecht, N., Yakubov, I., and Munson, K. (2007) The gastric H,K ATPase as a drug target - Past, present, and future. *J. Clin. Gastroenterol.* **41**, S226-S242.
5. Bamford, M. (2009) H⁺/K⁺ ATPase inhibitors in the treatment of acid-related disorders. *Prog. Med. Chem.* **47**, 75-162.

6. Mullin, J. M., Gabello, M., Murray, L. J., Farrell, C. P., Bellows, J., Wolov, K. R., Kearney, K. R., Rudolph, D., and Thornton, J. J. (2009) Proton pump inhibitors: actions and reactions. *Drug Discov. Today* **14**, 647-660.
7. Xian, Y., and Hebert, H. (1997) Three-dimensional structure of the porcine gastric H,K-ATPase from negatively stained crystals. *J. Struct. Biol.* **118**, 169-177.
8. Nishizawa, T., Abe, K., Tani, K., and Fujiyoshi, Y. (2008) Structural analysis of 2D crystals of gastric H⁺,K⁺-ATPase in different states of the transport cycle. *J. Struct. Biol.* **162**, 219-228.
9. Abe, K., Tani, K., Nishizawa, T., and Fujiyoshi, Y. (2009) Inter-subunit interaction of gastric H⁺,K⁺-ATPase prevents reverse reaction of the transport cycle. *Embo J.* **28**, 1637-1643.
10. Soumarmon, A., Grelac, F., and Lewin, M. J. (1983) Solubilization of active (H⁺ + K⁺)-ATPase from gastric membrane. *Biochim. Biophys. Acta* **732**, 579-585.
11. Soumarmon, A., Robert, J. C., and Lewin, M. J. (1986) Depolymerization of solubilized gastric (H⁺ + K⁺)-ATPase by n-octylglucoside or cholate. *Biochim. Biophys. Acta* **860**, 109-117.
12. Rabon, E. C., Gunther, R. D., Bassilian, S., and Kempner, E. S. (1988) Radiation inactivation analysis of oligomeric structure of the H,K-ATPase. *J. Biol. Chem.* **263**, 16189-16194.
13. Rabon, E. C., Bassilian, S., and Jakobsen, L. J. (1990) Glutaraldehyde cross-linking analysis of the C₁₂E₈ solubilized H,K-ATPase. *Biochim. Biophys. Acta* **1039**, 277-289.
14. Shin, J. M., and Sachs, G. (1996) Dimerization of the gastric H⁺,K⁺-ATPase. *J. Biol. Chem.* **271**, 1904-1908.
15. Lacapère, J. J., Robert, J. C., and Thomas-Soumarmon, A. (2000) Efficient solubilization and purification of the gastric H⁺,K⁺-ATPase for functional and structural studies. *Biochem. J.* **345**, 239-245.
16. Abe, K., Kaya, S., Hayashi, Y., Imagawa, T., Kikumoto, M., Oiwa, K., Katoh, T., Yazawa, M., and Taniguchi, K. (2003) Correlation between the activities and the oligomeric forms of pig gastric H/K-ATPase. *Biochemistry* **42**, 15132-15138.
17. Abe, K., Kaya, S., Taniguchi, K., Hayashi, Y., Imagawa, T., Kikumoto, M., Oiwa, K., and Sakaguchi, K. (2005) Evidence for a relationship between activity and the tetraprotomeric assembly of solubilized pig gastric H/K-ATPase. *J. Biochem.* **138**, 293-301.
18. Shin, J. M., Grundler, G., Senn-Bilfinger, J., Simon, W. A., and Sachs, G. (2005) Functional consequences of the oligomeric form of the membrane-bound gastric H,K-ATPase. *Biochemistry* **44**, 16321-16332.
19. Abe, K., Kaya, S., Imagawa, T., and Taniguchi, K. (2002) Gastric H/K-ATPase liberates two moles of P(i) from one mole of phosphoenzyme formed from a high-affinity ATP binding site and one mole of enzyme-bound ATP at the low-affinity site during cross-talk between catalytic subunits. *Biochemistry* **41**, 2438-2445.
20. le Maire, M., and Møller, J. V. (1986) Protein-protein and protein-lipid interactions of the sarcoplasmic reticulum Ca²⁺-ATPase, in *Sarcoplasmic Reticulum in Muscle Physiology* Vol. 1, pp 101-126, C.R.C. Press, Boca Raton, FLA, USA.
21. Lund, S., and Møller, J. V. (1988) Biphasic kinetics of sarcoplasmic-reticulum Ca⁺⁺-ATPase and the detergent-solubilized monomer. *J. Biol. Chem.* **263**, 1654-1664.
22. le Maire, M., Lind, K. E., Jorgensen, K. E., Roigaard, H., and Møller, J. V. (1978) Enzymatically active Ca²⁺ ATPase from sarcoplasmic-reticulum membranes, solubilized by non-ionic detergents - Role of lipid for aggregation of protein. *J. Biol. Chem.* **253**, 7051-7060.
23. Heegaard, C. W., le Maire, M., Gulik-Krzywicki, T., and Møller, J. V. (1990) Monomeric state and Ca²⁺ transport by sarcoplasmic-reticulum Ca²⁺-ATPase, reconstituted with an excess of phospholipid. *J. Biol. Chem.* **265**, 12020-12028.
24. Andersen, J. P. (1989) Monomer-oligomer equilibrium of sarcoplasmic-reticulum Ca-ATPase and the role of subunit interaction in the Ca²⁺ pump mechanism. *Biochim. Biophys. Acta* **988**, 47-72.
25. Mahaney, J. E., Albers, R. W., Waggoner, J. R., Kutchai, H. C., and Froehlich, J. P. (2005) Intermolecular conformational coupling and free energy exchange enhance the catalytic

- efficiency of cardiac muscle SERCA2a following the relief of phospholamban inhibition. *Biochemistry* **44**, 7713-7724.
26. Mahaney, J. E., Thomas, D. D., and Froehlich, J. P. (2004) The time-dependent distribution of phosphorylated intermediates in native sarcoplasmic reticulum Ca^{2+} -ATPase from skeletal muscle is not compatible with a linear kinetic model. *Biochemistry* **43**, 4400-4416.
 27. Esmann, M., Christiansen, C., Karlsson, K. A., Hansson, G. C., and Skou, J. C. (1980) Hydrodynamic properties of solubilized $(Na^+ + K^+)$ -ATPase from rectal glands of *Squalus-acanthias*. *Biochim. Biophys. Acta* **603**, 1-12.
 28. Brotherus, J. R., Jacobsen, L., and Jorgensen, P. L. (1983) Soluble and enzymatically stable $(Na^+ + K^+)$ -ATPase from mammalian kidney consisting predominantly of protomer alpha- beta- units - Preparation, assay and reconstitution of active Na^+,K^+ transport. *Biochim. Biophys. Acta* **731**, 290-303.
 29. Vilsen, B., Andersen, J. P., Petersen, J., and Jorgensen, P. L. (1987) Occlusion of $Na-22^+$ and $Rb-86^+$ in membrane-bound and soluble protomeric alpha- beta- units of Na,K -ATPase. *J. Biol. Chem.* **262**, 10511-10517.
 30. Cohen, E., Goldshleger, R., Shainskaya, A., Tal, D. M., Ebel, C., le Maire, M., and Karlsh, S. J. D. (2005) Purification of Na^+,K^+ -ATPase expressed in *Pichia pastoris* reveals an essential role of phospholipid-protein interactions. *J. Biol. Chem.* **280**, 16610-16618.
 31. Hayashi, Y., Mimura, K., Matsui, H., and Takagi, T. (1989) Minimum enzyme unit for Na^+/K^+ -ATPase is the alpha-beta-protomer - Determination by low-angle laser-light scattering photometry coupled with high-performance gel chromatography for substantially simultaneous measurement of ATPase activity and molecular-weight. *Biochim. Biophys. Acta* **983**, 217-229.
 32. Kobayashi, T., Tahara, Y., Takenaka, H., Mimura, K., and Hayashi, Y. (2007) Na^+ - and K^+ - dependent oligomeric interconversion among alpha beta-protomers, diprotomers and higher oligomers in solubilized Na^+/K^+ -ATPase. *J. Biochem.* **142**, 157-173.
 33. Mimura, K., Tahara, Y., Shinji, N., Tokuda, E., Takenaka, H., and Hayashi, Y. (2008) Isolation of stable (alpha beta)(4)-tetraprotomer from Na^+/K^+ -ATPase solubilized in the presence of short-chain fatty acids. *Biochemistry* **47**, 6039-6051.
 34. Shin, J. M., Munson, K., Vagin, O., and Sachs, G. (2009) The gastric HK-ATPase: structure, function, and inhibition. *Pflugers Arch.* **457**, 609-622.
 35. Toyoshima, C. (2009) How Ca^{2+} -ATPase pumps ions across the sarcoplasmic reticulum membrane. *Biochim. Biophys. Acta-Mol. Cell Res.* **1793**, 941-946.
 36. Møller, J. V., Olesen, C., Winther, A. M. L., and Nissen, P. (2010) The sarcoplasmic Ca^{2+} -ATPase: design of a perfect chemi-osmotic pump. *Q. Rev. Biophys.* **43**, 501-566.
 37. Morth, J. P., Pedersen, B. P., Toustrup-Jensen, M. S., Sorensen, T. L. M., Petersen, J., Andersen, J. P., Vilsen, B., and Nissen, P. (2007) Crystal structure of the sodium-potassium pump. *Nature* **450**, 1043-U1046.
 38. Shinoda, T., Ogawa, H., Cornelius, F., and Toyoshima, C. (2009) Crystal structure of the sodium-potassium pump at 2.4 angstrom resolution. *Nature* **459**, 446-U167.
 39. Morth, J. P., Pedersen, B. P., Buch-Pedersen, M. J., Andersen, J. P., Vilsen, B., Palmgren, M. G., and Nissen, P. (2011) A structural overview of the plasma membrane Na^+, K^+ -ATPase and H^+ -ATPase ion pumps. *Nat. Rev. Mol. Cell Biol.* **12**, 60-70.
 40. Pedersen, B. P., Buch-Pedersen, M. J., Morth, J. P., Palmgren, M. G., and Nissen, P. (2007) Crystal structure of the plasma membrane proton pump. *Nature* **450**, 1111-1114.
 41. Bublitz, M., Morth, J. P., and Nissen, P. (2011) P-type ATPases at a glance. *J. Cell Sci.* **124**, 2515-2519.
 42. Morii, M., Yamauchi, M., Ichikawa, T., Fujii, T., Takahashi, Y., Asano, S., Takeguchi, N., and Sakai, H. (2008) Involvement of the H_3O^+ -Lys-164-Gln-161-Glu-345 charge transfer pathway in proton transport of gastric H^+, K^+ -ATPase. *J. Biol. Chem.* **283**, 16876-16884.
 43. le Maire, M., Arnou, B., Olesen, C., Georgin, D., Ebel, C., and Møller, J. V. (2008) Gel chromatography and analytical ultracentrifugation to determine the extent of detergent binding

- and aggregation, and Stokes radius of membrane proteins using sarcoplasmic reticulum Ca²⁺-ATPase as an example. *Nature protocols* **3**, 1782-1795.
44. Skrabanja, A. T. P., Vanderhijden, H., and Depont, J. (1987) Transport ratios of reconstituted (H⁺+K⁺)-ATPase. *Biochim. Biophys. Acta* **903**, 434-440.
 45. Lowry, O. H., Rosebrough, N. J., Farr, A. L., and Randall, R. J. (1951) Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**, 265-275.
 46. Hartree, E. F. (1972) Determination of protein: a modification of the Lowry method that gives a linear photometric response. *Anal. Biochem.* **48**, 422-427.
 47. Bartlett, G. R. (1959) Phosphorous assay in column chromatography. *J. Biol. Chem.* **234**, 466-468.
 48. Møller, J. V., Lind, K. E., and Andersen, J. P. (1980) Enzyme-kinetics and substrate stabilization of detergent-solubilized and membraneous (Ca⁺⁺ Mg⁺⁺)-activated ATPase from sarcoplasmic-reticulum - Effect of protein-protein interactions. *J. Biol. Chem.* **255**, 1912-1920.
 49. Lenoir, G., Picard, M., Gauron, C., Montigny, C., Le Marechal, P., Falson, P., le Maire, M., Møller, J. V., and Champeil, P. (2004) Functional properties of sarcoplasmic reticulum Ca²⁺-ATPase after proteolytic cleavage at Leu(119)-Lys(120), close to the A-domain. *J. Biol. Chem.* **279**, 9156-9166.
 50. Cardi, D., Pozza, A., Arnou, B., Marchal, E., Clausen, J. D., Andersen, J. P., Krishna, S., Møller, J. V., le Maire, M., and Jaxel, C. (2010) Purified E255L mutant SERCA1a and purified PfATP6 are sensitive to SERCA-type inhibitors but insensitive to artemisinins. *J. Biol. Chem.* **285**, 26406-26416.
 51. Schuck, P. (2000) Size-distribution analysis of macromolecules by sedimentation velocity ultracentrifugation and Lamm equation modeling. *Biophys. J.* **78**, 1606-1619.
 52. Salvay, A. G., Santamaria, M., le Maire, M., and Ebel, C. (2007) Analytical Ultracentrifugation Sedimentation Velocity for the Characterization of Detergent-Solubilized Membrane Proteins Ca⁺⁺-ATPase and ExbB. *J. Biol. Physics* **33** 399-419.
 53. Ebel, C. (2011) Sedimentation velocity to characterize surfactants and solubilized membrane proteins. *Methods* **54**, 56-66.
 54. de Foresta, B., le Maire, M., Orłowski, S., Champeil, P., Lund, S., Møller, J. V., Michelangeli, F., and Lee, A. G. (1989) Membrane solubilization by detergent: use of brominated phospholipids to evaluate the detergent-induced changes in Ca²⁺-ATPase/lipid interaction. *Biochemistry* **28**, 2558-2567.
 55. Champeil, P., Menguy, T., Tribet, C., Popot, J. L., and le Maire, M. (2000) Interaction of amphipols with sarcoplasmic reticulum Ca²⁺-ATPase. *J. Biol. Chem.* **275**, 18623-18637.
 56. Møller, J. V., Juul, B., and le Maire, M. (1996) Structural organization, ion transport, and energy transduction of P-type ATPases. *Biochim. Biophys. Acta-Rev. Biomembr.* **1286**, 1-51.
 57. Tyagarajan, K., Lipniunas, P. H., Townsend, R. R., and Forte, J. G. (1997) The N-linked oligosaccharides of the beta-subunit of rabbit gastric H,K-ATPase: Site localization and identification of novel structures. *Biochemistry* **36**, 10200-10212.
 58. Lebowitz, J., Lewis, M. S., and Schuck, P. (2002) Modern analytical ultracentrifugation in protein science: A tutorial review. *Protein Sci.* **11**, 2067-2079.
 59. Edelstein, S. J., and Schachman, H. K. (1967) The simultaneous determination of partial specific volumes and molecular weights with microgram quantities. *J. Biol. Chem.* **242**, 306-311.
 60. Reynolds, J. A., and Tanford, C. (1976) Determination of molecular weight of the protein moiety in protein-detergent complexes without direct knowledge of detergent binding. *Proc. Natl. Acad. Sci. U. S. A.* **73**, 4467-4470.
 61. Gohon, Y., Pavlov, G., Timmins, P., Tribet, C., Popot, J.-L., and Ebel, C. (2004) Partial specific volume and solvent interactions of amphipol A8-35. *Anal. Biochem.* **334**, 318-334.
 62. Brown, P. H., Balbo, A., Zhao, H., Ebel, C., and Schuck, P. (2011) Density contrast sedimentation velocity for the determination of protein partial-specific volumes. *PLoS One* **6**, e26221.

63. Nury, H., Manon, F., Arnou, B., le Maire, M., Pebay-Peyroula, E., and Ebel, C. (2008) Mitochondrial bovine ADP/ATP carrier in detergent is predominantly monomeric but also forms multimeric species. *Biochemistry* **47**, 12319-12331.
64. Manon, F., and Ebel, C. (2010) Analytical ultracentrifugation, a useful tool to probe intrinsically disordered proteins, in *Instrumental Analysis of Intrinsically Disordered Proteins: Assessing Structure and Conformation*, pp 433-449, John Wiley and Sons, Hoboken.
65. Abe, K., Tani, K., and Fujiyoshi, Y. (2010) Structural and functional characterization of H^+,K^+ -ATPase with bound fluorinated phosphate analogs. *Journal of Structural Biology* **170**, 60-68.
66. Durr, K. L., Abe, K., Tavraz, N. N., and Friedrich, T. (2009) E2P State Stabilization by the N-terminal Tail of the H,K -ATPase beta-Subunit Is Critical for Efficient Proton Pumping under in Vivo Conditions. *J. Biol. Chem.* **284**, 20147-20154.
67. Olesen, C., Picard, M., Winther, A. M., Gyrop, C., Morth, J. P., Oxvig, C., Moller, J. V., and Nissen, P. (2007) The structural basis of calcium transport by the calcium pump. *Nature* **450**, 1036-1042.
68. Fedosova, N. U., Cornelius, F., and Klodos, I. (1998) E(2)P phosphoforms of Na,K -ATPase. I. Comparison of phosphointermediates formed from ATP and P(i) by their reactivity toward hydroxylamine and vanadate. *Biochemistry* **37**, 13634-13642.
69. Fedosova, N. U., Champeil, P., and Esmann, M. (2003) Rapid filtration analysis of nucleotide binding to Na,K -ATPase. *Biochemistry* **42**, 3536-3543.
70. Hardwicke, P. M., and Green, N. M. (1974) The effect of delipidation on the adenosine triphosphatase of sarcoplasmic reticulum. Electron microscopy and physical properties. *Eur J Biochem* **42**, 183-193.
71. Boulanger, P., le Maire, M., Bonhivers, M., Dubois, S., Desmadril, M., and Letellier, L. (1996) Purification and structural and functional characterization of FhuA, a transporter of the *Escherichia coli* outer membrane. *Biochemistry* **35**, 14216-14224.
72. Esmann, M. (1988) ATPase and phosphatase activity of Na^+,K^+ -ATPase: molar and specific activity, protein determination. *Methods Enzymol* **156**, 105-115.
73. Ayorinde, F. O., Gelain, S. V., Johnson, J. H., and Wan, L. W. (2000) Analysis of some commercial polysorbate formulations using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Rapid Commun. Mass Spectrom.* **14**, 2116-2124.
74. Picard, M., Dahmane, T., Garrigos, M., Gauron, C., Giusti, F., le Maire, M., Popot, J. L., and Champeil, P. (2006) Protective and inhibitory effects of various types of amphipols on the Ca^{2+} -ATPase from sarcoplasmic reticulum: a comparative study. *Biochemistry* **45**, 1861-1869.
75. le Maire, M., Champeil, P., and Møller, J. V. (2000) Interaction of membrane proteins and lipids with solubilizing detergents. *Biochim. Biophys. Acta* **1508**, 86-111.
76. Dahout-Gonzalez, C., Brandolin, G., and Pebay-Peyroula, E. (2003) Crystallization of the bovine ADP/ATP carrier is critically dependent upon the detergent-to-protein ratio. *Acta Crystallogr. D Biol. Crystallogr.* **59**, 2353-2355.
77. Jidenko, M., Nielsen, R. C., Sorensen, T. L., Møller, J. V., le Maire, M., Nissen, P., and Jaxel, C. (2005) Crystallization of a mammalian membrane protein overexpressed in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 11687-11691.
78. Schwarz, F., and Aebi, M. (2011) Mechanisms and principles of N-linked protein glycosylation. *Curr. Opin. Struct. Biol.* **21**, 576-582.
79. Dennis, J. W., Nabi, I. R., and Demetriou, M. (2009) Metabolism, cell surface organization, and disease. *Cell* **139**, 1229-1241.
80. Durr, K. L., Tavraz, N. N., Zimmermann, D., Bamberg, E., and Friedrich, T. (2008) Characterization of Na,K -ATPase and H,K -ATPase enzymes with glycosylation-deficient beta-subunit variants by voltage-clamp fluorometry in *Xenopus* oocytes. *Biochemistry* **47**, 4288-4297.
81. Musatov, A., and Robinson, N. C. (2002) Cholate-induced dimerization of detergent- or phospholipid-solubilized bovine cytochrome C oxidase. *Biochemistry* **41**, 4371-4376.

82. Ravaud, S., Do Cao, M. A., Jidenko, M., Ebel, C., Le Maire, M., Jault, J. M., Di Pietro, A., Haser, R., and Aghajari, N. (2006) The ABC transporter BmrA from *Bacillus subtilis* is a functional dimer when in a detergent-solubilized state. *Biochem. J.* **395**, 345-353.
83. Galian, C., Manon, F., Dezi, M., Torres, C., Ebel, C., Levy, D., and Jault, J. M. (2011) Optimized Purification of a Heterodimeric ABC Transporter in a Highly Stable Form Amenable to 2-D Crystallization. *PLoS One* **6**, e19677.
84. Hayashi, Y., Matsui, H., and Takagi, T. (1989) Membrane protein molecular weight determined by low-angle laser light-scattering photometry coupled with high-performance gel chromatography. *Methods Enzymol.* **172**, 514-528.
85. Philo, J., Talvenheimo, J., Wen, J., Rosenfeld, R., Welcher, A., and Arakawa, T. (1994) Interactions of neurotrophin-3 (NT-3), brain-derived neurotrophic factor (BDNF), and the NT-3.BDNF heterodimer with the extracellular domains of the TrkB and TrkC receptors. *J. Biol. Chem.* **269**, 27840-27846.
86. Beri, R. G., Walker, J., Reese, E. T., and Rollings, J. E. (1993) Characterization of chitosans via coupled size-exclusion chromatography and multiple-angle laser light-scattering technique. *Carbohydr. Res.* **238**, 11-26.
87. Tanford, C., Nozaki, Y., Reynolds, J. A., and Makino, S. (1974) Molecular characterization of proteins in detergent solutions. *Biochemistry* **13**, 2369-2376.
88. Riske, K. A., Politi, M. J., F. Reed, W., and Lamy-Freund, M. T. (1997) Temperature and ionic strength dependent light scattering of DMPG dispersions. *Chemistry and Physics of Lipids* **89**, 31-44.

Acknowledgments. Thanks are due to Hugues Lortat-Jacob, Aline Le Roy and Mathilde Lethier (Institut de Biologie Structurale Grenoble France) for helpful discussions, help in Analytical Ultracentrifugation experiments and in protein purification, respectively. We thank the 'Partnership for Structural Biology' in Grenoble for the use of the mass spectroscopy, AUC and PAOL platforms at the IBS. We thank the Danish Crown Slaughterhouse, Horsens, for supplying us with fresh material for preparation of H^+ , K^+ -ATPase.

FOOTNOTES

* This work was supported by the CNRS, by the Commissariat à l'Energie Atomique, by the Université Joseph Fourier – Grenoble 1, by the Université Paris-Sud and by grants from the E.U. (FP7 Grant agreement N°226507 - NMI3).

¹The abbreviations used are: SV, sedimentation velocity.

FIGURE LEGENDS

Fig. 1: Preparation of H^+,K^+ -ATPase, solubilized by $C_{12}E_8$, and purified by size exclusion chromatography with Tween 20.

2.4 mg of the tubulovesicular preparation of H^+,K^+ -ATPase, suspended in 0.7 ml in a buffer containing 40 mM MES (pH 6.0), 40 mM KCl, 0.25 M sucrose, 3 mM Mg^{2+} , and 1 mM EDTA was solubilized with 14 mg $C_{12}E_8$, and non-solubilized residues removed by centrifugation for 35 min at 100,000 g at 4°C. 0.5 ml of the supernatant was loaded onto a 10 x 300 mm Superose 6 column, equilibrated and eluted at the rate of 0.25 ml/min at 4°C in the above buffer, not containing $C_{12}E_8$ but with Tween 20 at 2 mg/ml (1A), or with $C_{12}E_8$ at 2 mg/ml but without Tween 20 (1B). Protein concentration, crosses; phospholipid concentration, red lines with diamonds; activity, continuous blue line, measured at 23°C as described in the Methods. The three arrows numbered as 1, 2, and 3 correspond to regions of the peak analyzed in sedimentation velocity and shown in Fig.3. V_0 , Void volume of the column. Fig 1C: activities in top fraction kept at 4°C measured as a function of time after column elution. Fig. 1D: Column calibration with the following water-soluble globular proteins: Thy, thyroglobulin; Fer, ferritin; Cat, catalase; HSA

human serum albumin; CA, carbonic anhydrase; TrpI, trypsin inhibitor. Fig. 1E: SDS-PAGE electrophoresis (11 % tricine gels) prepared from fractions collected from an HPLC experiment similar to that shown in Fig.1A. Lanes # 1-5 are from fractions eluting at 12- to 14.5 ml, corresponding to the peak of enzymatically active H^+,K^+ -ATPase. Lane MR contains the protein standard (phosphorylase *b* (92 kDa), BSA (67 kDa), ovalbumin (42 kDa), carbonic anhydrase (32 kDa), Trypsin inhibitor (22 kDa), α -lactalbumin (12 kDa). Lanes # 6-10 are from the third protein peak, corresponding to elution at 15 - 17 ml. The lane labelled TV represents the $C_{12}E_8$ -solubilized tubulovesicular membranes.

Fig. 2. Mass spectroscopy of H^+, K^+ -ATPase

A: MALDI TOF spectrum providing molecular masses of 42488 Da and 114781 Da, corresponding to the glycosylated β -subunit and to the α -subunit ; **B,C:** LC-ESI TOF MS spectra providing molecular masses of 41662 Da corresponding to the glycosylated β -subunit (**B**), and of 114112 Da corresponding to the α -subunit (**C**).

Fig. 3: Sedimentation velocity of H^+, K^+ -ATPase.

Experiments were done at 42,000 rpm and 7°C, for different fractions of the main H^+,K^+ -ATPase peak from gel filtration. The solvent was 40 mM KCl, 40 mM Mes buffer (pH 6.0), 3 mM Mg^{2+} , 1 mM EDTA, and 0.25 M sucrose 2 mg/ml Tween 20. A-B: Superposition, for the central fraction, of selected experimental and fitted profiles (top panels) and of their differences (bottom panels) obtained at 280 nm (**A**) and using interference optics (**B**). The last profiles correspond to 14 h of sedimentation. **C:** Superposition of the $c(s)$ distributions obtained in the range of 0.1–10 S from the two optics for, from top to bottom, the first, central and last fractions of the main peak. **D:** Evolution of the content of the fractions from gel filtration along the main peak of H^+,K^+ -ATPase, evaluated from the area under the $c(s)$ curve from the data at 280 nm. In this last panel, the sedimentation coefficients are given as $s_{20,w}$.

Fig. 4. Graphical analysis of the sedimentation velocity of H^+, K^+ -ATPase in hydrogenated and deuterated solvents

The analysis is made in terms of a complex composed of two pseudo-components: the glycosylated protein and bound (detergent + lipid). Common input values are: $\bar{v}_{PG}=0.728$ ml/g; $\eta_H=1.834$ cP; $\rho_H=1.0354$ g/ml; $\eta_D=2.2383$ cP; $\rho_D=1.1257$ g/ml; $(M_D/M_H)_{PG}=1.011$; $(M_D/M_H)_{D+L}=1.016$. On each panel, the four drawn curves give the ensemble of solutions (δ_{D+L} ; \bar{v}_{D+L}), with δ_{D+L} the amount in g/g of glycosylated protein and \bar{v}_{D+L} the partial specific volume for bound detergent plus lipid. Lines are drawn for the limiting experimental values of the sedimentation coefficients determined from the $c(s)$ analysis: $s_H - \Delta s_H = 3.2S$ (khaki line, gold point); $s_H + \Delta s_H = 3.26S$ (green line, red point); $s_D - \Delta s_D = 1.5S$ (cyan line, navy point); $s_D + \Delta s_D = 1.65S$ (aquamarine line, blackpoint). Indexes PG and D+L, H and D, correspond to glycosylated protein, detergent and lipid, hydrogenated solvent and deuterated solvent, respectively. For each panel, the ff_{min} and assembly state hypothesis are written. Mathematical solutions are defined by the area encircled by the four curves.

Fig. 5. Molar protein mass of Tween 20-solubilized H^+, K^+ -ATPase after size exclusion chromatography with registration of light absorbance at 280 nm, refractive index and light scattering

Superposition of A_{280} (green, thin), differential refractive index (blue, thin), light scattering (red) signals, in arbitrary units (left axis), and molar masses for the glycosylated protein (blue, thick), detergent + lipid (green, thick) and complex, *i.e.* their sum (purple) (right axis). The calculation considers $(\partial n/\partial c)$ for the glycosylated protein to be 0.185 ml/g (with 9 kDa of carbohydrate for a monomeric α,β protein), and $(\partial n/\partial c)$ for detergent + lipid, to be 0.082 ml/g, with extinction coefficients of 0.975 and 0.0115 ml/(mg cm), respectively. H^+,K^+ -ATPase after solubilization was subjected to size exclusion chromatography on a Superose 6 column; fractions corresponding to the central part of the main protein peak were pooled, and 30 μ l were injected onto a KW 804 column equilibrated at 4°C, with a flow rate of

0.5 ml/min, with an eluent filtered at 0.1 μm and containing 40 mM KCl, 40 mM MES (pH 6), 0.25 M sucrose, 3 mM MgCl₂, 1mM EDTA, 2 mg/ml Tween 20.

Fig. 6. Model of the α,β monomer H⁺,K⁺-ATPase.

Model of the H⁺,K⁺-ATPase α,β monomer complex as represented by the structure of the pig kidney Na⁺,K⁺-ATPase $\alpha 1\beta 1$ complex in the potassium-occluded state trapped with MgF₄²⁻ (PDB 3KDP (37)). The α -subunit is in blue and the β -subunit in wheat. The structure spans approximately 160 \AA in the long dimension, excluding glycosylations on the β -subunit.

	M	δ	\bar{v}	$\partial n/\partial c$	M_D/M_H	$E_{0.1\%, 280\text{nm}}$
	kDa	g per g of protein	ml/g	ml/g		ml/(mg cm) ^o
α, β H^+, K^+ -ATPase	147.346 ^a		0.734 ^a	0.187 ^b	1.011 ^a	1.035 ^a
carbohydrate	9 ^c	0.061 ^c	0.63 ^d	0.15 ^e	1.019 ^a	0 ^f
glycosylated protomer ^g	156.346		0.728	0.185	1.011	0.975
Tween 20	(see text)	(see text)	0.869 ^h , 0.87 ^c	0.082 ^c	1.016 ^a	0.0115 ^c
Phospholipid	36.8 ^c	0.25 ^c	0.975 ⁱ	0.12 ^j	1.016 ^f	0 ^f
Cholesterol	3.5 ^c	0.024 ^c	0.975 ⁱ			

Table 1. Values characterizing the components of the solubilized and chromatographed H^+,K^+ -ATPase complex

^a Calculated from amino acid sequence or estimated from chemical composition, α - and β - subunits have molecular masses of 114.288 and 33.058 kDa, respectively; \bar{v} calculated using the Sednterp program (<http://www.jphilo.mailway.com>) is given at 7°C ($\bar{v} = 0.739$ ml/g at 20°C); M_D/M_H values are derived considering 80% of maximum H-D exchange for the protein, and full exchange for the other components. ^b from (84). ^c This work; results concerning Tween 20 are described in the Supplementary Fig. S2 and used in this paper. ^d from (85). ^e value for dextran from (86). ^f considered values. ^g derived from the values given in the two lines above. ^h from (75). ⁱ calculated according to the lipid composition of the membrane (Table 1S) and the partial specific volumes given for individual lipids by (87); \bar{v} for cholesterol is ill-defined (see (87)). ^j given for dimyristoyl phosphatidylglycerol by (88).

Methods	Results	Hypothesis
Chromatography	Phospholipids: 0.25 g/g	
Chromatography	Cholesterol: 0.024 g/g	
Chromatography	Stokes radius = 6.3 nm	column calibration with globular soluble proteins
MALDI-TOF and ESI Mass Spectrometry	Non glycosylated α subunit β subunit: $\delta_G = 0.06$ g/g (9 kDa)	
SV	$s = 3.23$ S, $s_{20,w} = 7.5$ S (main species)	
SV : J/A_{280} -value	$\delta_D = 1.6$ g/g	Assuming a glycosylation of 0.06 g/g and 0.274 g lipid/g protein and based on $E_{0.1\%, 280nm}$ and $\partial n/\partial c$ from Table1
SV + chromatography: s - and R_s -values	1: Buoyant molecular mass $M_b = 42.3$ kDa 2: α, β -protomer, however δ_D is ill-defined.	1: Without hypothesis 2: Assuming a glycosylation of 0.06 g/g and 0.274 g lipid/g protein.
s_H/s_D	$s_D = 1.6 \pm 0.1$ S 1: $\bar{v}_{complex} = 0.818 \pm 0.015$ ml/g 2: $\delta_D = 1.0 \pm 0.7$ g/g $R_s = 7.9 \pm 1$ nm ($f/f_{min} = 1.6 \pm 0.2$) 3: α, β -protomer with $f/f_{min} = 1.5 \pm 0.1$ dimer $\alpha_2\beta_2$ is rejected	1: with $M_D/M_{H complex} = 1.014$ 2: also based on glycosylation of 0.06 g/g and 0.274 g lipid/g protein.. 3: with $M_D/M_{H complex} = 1.014$
Chromatography coupled to light scattering	$M_{PG} = 133$ kDa $\delta_{D+L} = 0.85-1.3$ g/g α, β -protomer	Based on $E_{0.1\%, 280nm}$ and $\partial n/\partial c$ from Table1

Table 2. Summary of the composition and hydrodynamic data of the Tween 20-solubilized H^+,K^+ -ATPase obtained by various procedures in the present study.
Values in g/g are given in g per non-glycosylated protein.




Fig. 1




Fig. 2




Fig. 3




Fig.4




Fig. 5

H⁺,K⁺-ATPase is a monomer




Fig. 6