AARHUS UNIVERSITY

Coversheet

This is the accepted manuscript (post-print version) of the article.

Contentwise, the accepted manuscript version is identical to the final published version, but there may be differences in typography and layout.

How to cite this publication

Please cite the final published version:

J. H. Jeppesen, R. H. Jacobsen, R. N. Jørgensen, A. Halberg and T. S. Toftegaard (2017). Identification of High-Variation Fields based on Open Satellite Imagery. In *Advances in Animal Biosciences*, 8 (SI2), (Papers presented at the 11th European Conference on Precision Agriculture (ECPA 2017), John McIntyre Centre, Edinburgh, UK, July 16–20 2017), pp 388-393.

DOI: <https://doi.org/10.3168/jds.2016-11533>

Publication metadata

Title:	Identification of High-Variation Fields based on Open Satellite Imagery
Author(s):	J. H. Jeppesen, R. H. Jacobsen, R. N. Jørgensen, A. Halberg and T. S. Toftegaard
Journal:	<i>Advances in Animal Biosciences</i>
DOI/Link:	https://doi.org/10.3168/jds.2016-11533
Document version:	Accepted manuscript (post-print)

This article has been published in a revised form in *Advances in Animal Biosciences* <https://doi.org/10.3168/jds.2016-11533>. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © The authors and The Animal Consortium 2017.

General Rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

If the document is published under a Creative Commons license, this applies instead of the general rights.

Identification of High-Variation Fields based on Open Satellite Imagery

J. H. Jeppesen^{1†}, R. H. Jacobsen¹, R. N. Jørgensen¹, A. Halberg² and T. S. Toftegaard¹

¹Department of Engineering, Aarhus University, Finlandsgade 22, 8000 Aarhus C, Denmark; ²Airinov Denmark, Vejstrupgaards Alle 5882 Vejstrup, Denmark

This paper proposes a simple method for categorizing fields on a regional level, with respect to intra-field variations. It aims to identify fields where the potential benefits of applying precision agricultural practices are highest from an economic and environmental perspective. The categorization is based on vegetation indices derived from Sentinel-2 satellite imagery. A case study on 7678 winter wheat fields is presented, which employs open data and open source software to analyze the satellite imagery. Furthermore, the method can be automated to deliver categorizations at every update of satellite imagery, hence coupling the geospatial data analysis to direct improvements for the farmers, contractors, and consultants.

Keywords: Remote sensing, vegetation index, zonal statistics, open data, decision support

Introduction

Data-driven precision agriculture has the potential to improve the yield while minimizing the environmental impact, leading to an economic and environmental benefit (Skjødtt *et al.*, 2003, Mamo *et al.*, 2003, Kindred *et al.*, 2014). This is achieved by considering the intra-field variations while employing farming practices. A prerequisite of precision agriculture is an increasing practice of advanced machinery and obtainment of application maps. Consequently, it is desirable to use an automated process of identifying fields where the potential benefits of employing precision agriculture are highest. It has been shown how the profit of employing precision agriculture increases as the intra-field variations increases (Nilsson, 2010; Delin *et al.*, 2015). This calls for a method for identification, on a regional or national scale, of fields exhibiting high degrees of variations. Hereby, the benefit of employing precision agriculture is maximized regarding economic and environmental gain.

Satellite imagery has become readily available following the opening of access to Landsat data in 2008 (Wulder *et al.*, 2012) by NASA and the open access to all Sentinel satellites by ESA. This is complemented by significant improvements in the open source software for handling and processing the data (Jeppesen *et al.*, 2015). Multi-spectral Sentinel-2 satellite imagery with resolutions of 10 m × 10 m is freely available from open online data sources with a current global revisit time of 10 days, which will be halved to 5 days when Sentinel-2B will launch (Drusch *et al.*, 2012). One possible application of such satellite imagery is to monitor and track crop development via calculation of vegetation indices. This paper investigates and presents a

simple method for categorizing fields on a regional level according to their intra-field variations based on Sentinel-2 satellite imagery, which results in a possibility for identifying fields with the highest potential for further action with precision agricultural practices (Nilsson, 2010; Delin *et al.*, 2015). This can be used for calculating e.g. variable rate nitrogen application maps based on the satellite imagery (Seelan *et al.*, 2003), or for selecting fields for further investigation using e.g. drones. The method can be implemented using open source software. It can be automated to continuously provide categorizations as new satellite images become available.

Materials and methods

The area under study is Funen in Denmark. The island has an area of 3100 km² and counts presently 7678 winter wheat fields. Funen has historically been recognized for its rich farmland.

An overview of the processing workflow of the proposed method is illustrated in Figure 1. Initially, satellite images are downloaded and used to create a vegetation index from combinations of measured reflectance in the different spectral bands. The vegetation index is subsequently used along with field polygons to calculate zonal statistics, including the standard deviation of the pixel values of the vegetation index inside each field polygon. The field-level standard deviation is used as a measure of the intra-field variation within the specific field. All fields within the region of interest are then compared and ranked based on the standard deviation. This results in a categorization based on the intra-field variations. The method can be implemented using open source processing software, such as QGIS, or in custom applications using libraries such as GDAL and R.

[†] E-mail: jhj@eng.au.dk

Figure 1 Overview of the software and data workflow.

QGIS was used for the data processing, which is an open source GIS application for data processing and visualization. To carry out the zonal statistics, the required field polygons for Denmark were downloaded and the winter-wheat polygons were extracted. The Sentinel-2 images were downloaded from Amazon AWS, and orthophotos from spring 2016 were used as base maps. Orthophotos of Denmark are produced on a yearly basis and provided freely by The Danish Geodata Agency.

The ESA Sentinel-2 satellite provides 13-band multi-spectral imagery, with a spatial resolution down to 10 m x 10 m, at no cost. The imagery is divided in 100 km x 100 km tiles, which is provided by several online platforms (e.g. ESA Scientific Hub, Amazon AWS, Google Cloud Platform). In the proposed method, the required Sentinel-2 bands are downloaded in the 100 km x 100 km tiles in jpeg2000 format. The data is then converted to GeoTIFF and merged to produce a single image for each spectral band for the entire region of interest.

A vegetation index is used in remote sensing to estimate properties of vegetation. For instance, NDVI (Normalized Difference Vegetation Index) can be used to estimate biomass (Rouse *et al.*, 1973). Two or more spectral bands from a multi-spectral sensor are used to calculate vegetation indices. Table 1 gives an overview of the center wavelengths and spatial resolution of 7 bands in the Sentinel-2 satellite, along with the center wavelengths of a 4 band multi-spectral drone (Multispec 4C), such as the ones used by Airinov, for comparison. Note that the altitude of flight determines the spatial resolution of the drone.

In the past, a wide variety of vegetation indices have been developed for different purposes, such as estimation of biomass (Rouse *et al.*, 1973) and chlorophyll concentration (Gitelson *et al.*, 1996). Table 2 shows an overview of some of the most widespread vegetation indices as defined for Sentinel-2 imagery, alongside with the achievable spatial resolution.

The choice of vegetation index is flexible due to the many spectral bands provided by the Sentinel-2 satellite. However, it should be considered thoroughly, for instance, NDVI is simple to calculate and provides an easy way to identify vegetation, but it saturates at high chlorophyll concentrations, directing one towards GNDVI to alleviate this problem

(Gitelson *et al.*, 1996). REIP (Red Edge Inflection Point) is considered to be less sensitive to soil reflectance (Raper *et al.*, 2015), but is only valid where vegetation is present (Clevers *et al.*, 2001), resulting in inconsistencies when measuring e.g. bare soil or water. Combined vegetation indices have also been developed, such as the SCCCI (Simplified Canopy Chlorophyll Content Index) which provides good correlation with crop nitrogen status (Raper *et al.*, 2015).

The zonal statistics (cf. Figure 1) calculates the mean and standard deviation of the vegetation index pixel values within specified field polygons. The first step is to obtain the field polygons, which should be openly available within the EU. In Denmark, field polygons are provided by The Danish Agrifish Agency and are updated each spring for the new growth season. A specific crop type is queried, and the field polygons are reduced in size to avoid inclusion of mixed pixels in the zonal statistics. Mixed pixels are present due to the geometric precision and the resolution of the satellite, where e.g. a road next to the field is sensed as part of the field. The zonal statistics are calculated, and the intra-field standard deviation is then used to categorize fields, such that a field with a low and high standard deviation corresponds to a field exhibiting low and high intra-field variation respectively.

Results

To validate our proposed method we provide a case study on an NDVI based categorization of 7678 winter wheat fields on Funen, Denmark, on 4th of June 2016. However, the method is applicable to all crop types with meaningful vegetation indices.

The vegetation index used for the case study was chosen to be NDVI, as it is well known and the use of it is widespread in the agricultural domain. The Sentinel-2 images were converted to GeoTIFF and merged using the raster translate and merge tools in QGIS. The winter wheat field polygons were selected, and the fixed buffer size tool was used to reduce the polygons in size by 20 m, to avoid mixed pixels. The calculations of the zonal statistics were carried out in QGIS on the reduced size

polygons, where the mean and standard deviation of the NDVI values within individual fields were found. The standard deviations were then compared on a regional level using all 7678 fields and used to categorize the fields into five categories defined by equal numbered quantiles. The five categories are found according to the statistics and then used for categorization of the original size polygons. The categorized winter-wheat field polygons can then be used to highlight the fields exhibiting the highest intra-field variation.

To determine the validity of the Sentinel-2 NDVI map, a multi-spectral drone and satellite imagery of a field on Funen were captured on 20th of May 2016 and 4th of June 2016 respectively. The drone NDVI maps were provided by Airinov and captured using a Multispec 4C sensor. Figure 2a and 2b, which provide an overview of the resulting NDVI maps of the field, show good agreement with each other. Due to the higher resolution of the NDVI map resulting from the Multispec 4C sensor, the drone NDVI image (Figure 2a) can reveal the tracks from the machinery that has been operating on the field. These details cannot be distinguished on the satellite NDVI image (Figure 2b) because of the lower spatial resolution.

The drone image is interpreted as a ground truth with respect to the NDVI calculated from the satellite image. It should be noted that the NDVI absolute values differ on the two measurements. This is caused by a range of factors, such as different wavelengths of the multi-spectral cameras, atmospheric disturbances, different dates, and sun orientation. Nevertheless, the important factor in the proposed method are the relative values for the same measurement. Figure 2c shows the reduced size polygon used for statistics, where the outermost 20 m have been cropped out to circumvent the mixed pixel effects.

Table 1 Center wavelength and spatial resolution of sensor bands (Drusch *et al.*, 2012, Haghighattalab *et al.*, 2016).

Band	Sentinel-2 satellite	Multispec 4C (drone)
Blue (B)	490 nm, 10 m x 10 m (B2)	-
Green (G)	560 nm, 10 m x 10 m (B3)	550 nm
Red (VIR)	665 nm, 10 m x 10 m (B4)	660 nm
Red edge 1 (RE1)	705 nm, 20 m x 20 m (B5)	-
Red edge 2 (RE2)	740 nm, 20 m x 20 m (B6)	735 nm
Near infrared 1 (NIR1)	783 nm, 20 m x 20 m (B7)	790 nm
Near infrared 2 (NIR2)	842 nm, 10 m x 10 m (B8)	-

Table 2 Vegetation indices as defined for the Sentinel-2 satellite

Vegetation index	Expression	Spatial res.	Reference
GNDVI	$\frac{NIR2 - G}{NIR2 + G}$	10 m x 10 m	(Gitelson <i>et al.</i> , 1996)
NDVI	$\frac{NIR2 - VIR}{NIR2 + VIR}$	10 m x 10 m	(Rouse <i>et al.</i> , 1973)
NDRE	$\frac{NIR1 - RE1}{NIR2 + RE1}$	20 m x 20 m	(Barnes <i>et al.</i> , 2000)
REIP	$705 + 35 \left(\frac{(VIR + NIR) - RE1}{RE2 - RE1} \right)$	20 m x 20 m	(Herrmann <i>et al.</i> , 2011)
SCCCI (combined index)	NDRE/NDVI	20 m x 20 m	(Barnes <i>et al.</i> , 2000)

The standard deviation of the pixels within the fields provides information on the intra-field variation as shown in the histograms in Figure 3. The categorization illustrates the fields with high standard deviation values, and hence the fields with high intra-field variations. These fields are expected to provide the highest potential for further action through precision agricultural practices. Furthermore, it is possible to track the mean values within the fields to quantify the status of individual fields compared on a regional or national scale. Examples of the categorized fields are seen in Figure 4.

Figure 5 shows examples of two winter-wheat fields from the lowest and highest variation categories. An important benefit of open satellite imagery is the availability of historic data, which grants the possibility of investigating the development of the fields in the months preceding the categorization date. NDVI maps of the two fields were generated for 8th of April and 5th of May 2016, along with the NDVI map used for the categorization on 4th of June. Note that a direct comparison between the two fields is less meaningful due to unknown differences in the early crop cycle conditions, such as the sowing time.

The top row in Figure 5 shows the low-variation field, where the benefits of variable rate nitrogen application are limited due to the homogeneity of the field. Looking at the high-variation field in the bottom row, significant benefits can be achieved from applications maps based on satellite imagery (Seelan *et al.*, 2003), or even larger benefits might be achieved through further investigation and creation of application maps based on high-resolution drone imagery.

Discussion

The proposed method relies on the application of open data and open source software, which have been made possible by the recent increase in publicly available data and the improvements in open source software for advanced geospatial analysis (Jeppesen *et al.*, 2016). This contributes to the overall cost-efficiency of our approach. The method can be automated, continuously providing updates for every satellite image throughout the growth season. Satellite imagery does have limits regarding cloud coverage, resulting in uncertain measurement times. Although cloud coverage is a problem, one can expect to have several cloud-free images of a field throughout the growing season. Moreover, cloud masks are supplied when downloading the satellite data, hence the modifications needed

Figure 2 Comparisons of a single field using a) drone NDVI on 20th of May, b) Sentinel-2 satellite NDVI on 4th of June, and in c) a highlight of the reduced size polygon used for zonal statistics.

Figure 3 Histograms showing a) mean NDVI values and b) categorized NDVI standard deviation values.

to perform cloud masking before statistical analysis are minimal, and the categorization can be done based on the cloud-free field polygons. Finally, Landsat 8 can also be used, however, it does not measure the RE1, RE2, and NIR1 bands, and it has a spatial resolution of 30 m × 30 m (Roy *et al.*, 2014).

The proposed method is agnostic to the choice of vegetation index, which can be chosen to map certain vegetation properties, for instance, biomass (e.g. NDVI) or chlorophyll density (e.g. NDRE). Due to the short revisit time of the satellite, the method allows us to track the development of intra-field variations over a crop cycle in a simple manner. This provides a valuable tool for farmers and consultants when planning which fields to visit. Secondly, machine pool services and drone services may offer site specific services to new customers based on the intra-field variations. Thirdly, coarse nitrogen application maps can be generated in an automated manner for all fields visible by the satellite (Seelan *et al.*, 2003). Drone operations quickly become impractical and overwhelmingly expensive with frequent imagery, hence the method complements the use of drone imagery for an in-depth investigation of high-variation fields through its improved spatial resolution.

To publish the results to a web-platform in a seamless manner, GeoNode can be used. It is a web-based spatial data

infrastructure providing a simple interface for upload, download, and visualization of geospatial data online (Jeppesen *et al.*, 2016). It provides a back-end system, based on GeoServer and a database (e.g. PostGIS), which can also be used along with OGC web services, such as WMS and WFS, to create customer-friendly web pages to show the results.

Conclusions

In this paper, we proposed and validated a simple method for categorization of fields according to their intra-field variations of vegetation indices calculated from multispectral satellite imagery. The study categorized 7678 winter wheat fields on Funen, Denmark, where drone imagery was used to establish ground truth. The proposed method showed good potential for identifying fields where additional conceivable agricultural practices, such as variable rate nitrogen application, would provide the highest potential for an economic benefit. The presented method is based on open data and can be implemented using open source software, such that the cost of employing the method is minimal. Furthermore, it can be automated, such that categorization is updated when new satellite imagery is acquired. Future work includes

Figure 4 Map of categorized winter wheat fields, with base maps showing a) RGB orthophoto and b) Sentinel-2 NDVI map from 4th of June 2016.

Figure 5 Examples of NDVI maps from a field from the lowest variation category on a) 8th of April, b) 5th of May, and c) 4th of June, along with NDVI maps of a field from the highest variation category on d) 8th of April, e) 5th of May, and f) 4th of June. All dates from year 2016.

determination of nitrogen application maps based on satellite imagery.

Acknowledgements

This work was funded by Innovation Fund Denmark as part of the Future Cropping project (J. nr. 5107-00002B), a large Danish research project with the aim of increasing agricultural productivity while minimizing the environmental impact beyond the current state of the art.

References

- Barnes E, Clarke T, Richards S, Colaizzi PD, Haberland J, Kostrzewski M, *et al.* 2000. Coincident Detection of Crop Water Stress, Nitrogen Status and Canopy Density Using Ground-Based Multispectral Data. Proceedings of the Fifth International Conference on Precision Agriculture.
- Clevers JGPW, de Jong SM, Epema GF, van der Meer F, Bakker WH, Skidmore AK and Addink EA 2001. MERIS and the red-edge position. International Journal of Applied Earth Observation and Geoinformation 3 (4), 313–320.
- Delin S, Gruvæus I, Wetterlind J, Stenberg M, Frostgård G, Börling K, *et al.* 2015. Fertilisation for optimised yield can minimise nitrate leaching in grain production. Proceedings of the International Fertiliser Society 774.

- Drusch M, Del Bello U, Carlier S, Colin O, Fernandez V, Gascon F, *et al.* 2012. Sentinel-2: ESA's Optical High-Resolution Mission for GMES Operational Services. *Remote Sensing of Environment* 120, 25–36.
- Gitelson AA, Kaufman YJ and Merzlyak MN 1996. Use of a green channel in remote sensing of global vegetation from EOS-MODIS. *Remote Sensing of Environment* 58 (3), 289–298.
- Haghighattalab A, González Pérez L, Mondal S, Singh D, Schinstock D, Rutkoski J, *et al.* 2016. Application of unmanned aerial systems for high throughput phenotyping of large wheat breeding nurseries. *Plant Methods* 12 (1), 35.
- Herrmann I, Pimstein A, Karnieli A, Cohen Y, Alchanatis V and Bonfil DJ 2011. LAI assessment of wheat and potato crops by VENμS and Sentinel-2 bands. *Remote Sensing of Environment* 115 (8), 2141–2151.
- Jeppesen JH, Jacobsen RH, Jørgensen RN and Toftegaard TS 2016. Towards Data-Driven Precision Agriculture using Open Data and Open Source Software. CIGR-AgEng conference. <http://conferences.au.dk/cigr-2016/full-papers/> (retrieved 07/12/16).
- Kindred DR, Milne AE, Webster R, Marchant BP and Sylvester-Bradley R 2014. Exploring the spatial variation in the fertilizer-nitrogen requirement of wheat within fields. *The Journal of Agricultural Science* 153 (01), 25–41.
- Mamo M, Malzer GL, Mulla DJ, Huggins DR and Strock J 2003. Spatial and Temporal Variation in Economically Optimum Nitrogen Rate for Corn. *Agronomy Journal* 95 (4), 958–964.
- Nilsson C 2010. Möjligheter att minska kväveutlakningen genom att anpassa kvävegödslingen till variationer inom stråsädesfält. http://stud.epsilon.slu.se/1531/1/nilsson_c_100701.pdf (retrieved 05/12/16).
- Raper TB and Varco JJ 2015. Canopy-scale wavelength and vegetative index sensitivities to cotton growth parameters and nitrogen status. *Precision Agriculture* 16, 62–76.
- Rouse JW, Hass RH, Schell JA and Deering DW 1973. Monitoring vegetation systems in the great plains with ERTS. Third Earth Resources Technology Satellite (ERTS) Symposium 1, 309–317.
- Roy DP, Wulder MA, Loveland TR, Woodcock CE, Allen RG, Anderson MC, *et al.* 2014. Landsat-8: Science and product vision for terrestrial global change research. *Remote Sensing of Environment* 145, 154–172.
- Seelan SK, Laguet S, Casady GM and Seielstad GA 2003. Remote sensing applications for precision agriculture: A learning community approach. *Remote Sensing of Environment* 88 (1-2), 157–169.
- Skjød P, Hansen PM and Nyholm RN 2003. Sensor Based Nitrogen Fertilization Increasing Grain Protein Yield in Winter Wheat. <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.193.135&rep=rep1&type=pdf> (retrieved 01/12/16).
- Wulder MA, Masek JG, Cohen WB, Loveland TR and Woodcock CE 2012. Opening the archive: How free data has enabled the science and monitoring promise of Landsat. *Remote Sensing of Environment* 122, 2–10.