AARHUS UNIVERSITY

Coversheet

This is the accepted manuscript (post-print version) of the article.

Contentwise, the accepted manuscript version is identical to the final published version, but there may be differences in typography and layout.

How to cite this publication

Please cite the final published version:

Dias, G. F. (2017). The time-varying GARCH-in-mean model. *Economics Letters*, 157, 129-132.
<https://doi.org/10.1016/j.econlet.2017.06.005>

Publication metadata

Title:	The time-varying GARCH-in-mean model
Author(s):	Dias, G. F.
Journal:	<i>Economics Letters</i>
DOI/Link:	10.1016/j.econlet.2017.06.005
Document version:	Accepted manuscript (post-print)
Document license:	<i>[If the document is published under a Creative Commons, enter link to the license here]</i>

General Rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

If the document is published under a Creative Commons license, this applies instead of the general rights.

The time-varying GARCH-in-mean model

Gustavo Fruet Dias^{a,1}

^a*CREATES and Aarhus University, Fuglesangs Allé 4, 8210 Aarhus V, Denmark.
E-mail address: gdias@econ.au.dk*

Abstract

1 I propose an estimation strategy for the stochastic time-varying risk premium
2 parameter in the context of a time-varying GARCH-in-mean (TVGARCH-in-
3 mean) model. A Monte Carlo study shows that the proposed algorithm has good
4 finite sample properties. Using monthly excess returns on the CRSP index, I
5 document that the risk premium parameter is indeed time-varying and shows
6 high degree of persistence.

JEL classification numbers: C13, C15, C22, G12

Keywords: risk-return tradeoff, time-varying coefficients, iterative estimators,
GARCH-type models

¹I thank George Kapetanios, Cristina Scherrer and an anonymous referee for constructive suggestions. I acknowledge support from CREATES - Center for Research in Econometric Analysis of Time Series (DNRF78), funded by the Danish National Research Foundation.

7 1. Introduction

8 Asset pricing theories suggest that riskier assets should demand higher ex-
 9 pected returns. Using Merton's (1973) theoretical framework, the conditional
 10 expectation of the market excess returns reads

$$\mathbb{E}(r_{t+1}^m | \mathcal{F}_t) - r_t^f = \lambda_t \text{Var}(r_{t+1}^m | \mathcal{F}_t), \quad (1)$$

11 where r_{t+1}^m and r_t^f are the returns on the market portfolio and risk-free asset,
 12 $\mathcal{F}_t$ is the market-wide information available at time t , and λ_t is the coefficient
 13 of relative risk aversion defined as the elasticity of marginal value with respect
 14 to wealth. Most studies assume the risk-return trade-off is constant over time
 15 and linear in the variance, which is usually associated with the reasons behind
 16 mixed empirical evidences when estimating the risk-return trade-off (Linton
 17 and Perron (2003), Brandt and Wang (2010), Christensen, Dahl, and Iglesias
 18 (2012), among others). To address this issue, I adopt the time-varying GARCH-
 19 in-mean (TVGARCH-in-mean) model in the spirit of Anyfantaki and Demos
 20 (2016) which allows λ_t to be a time-varying stochastic process and put forward
 21 a feasible estimation strategy for λ_t (see references in Anyfantaki and Demos
 22 (2016) for variants of the TVGARCH-in-mean models). Specifically, I com-
 23 bine Giraitis, Kapetanios, and Yates's (2013) time-varying kernel least squares
 24 estimator with Linton and Perron's (2003) semiparametric iterative approach
 25 to estimate the time-varying risk premium coefficient. A Monte Carlo study
 26 shows that the proposed algorithm has good finite sample properties. Using the
 27 excess returns of the Center for Research on Security Prices (CRSP) index, I
 28 document that the risk premium parameter is indeed time-varying, alternating
 29 positive (statistically significant) and nonsignificant values over time.

30 2. The time-varying GARCH-in-mean

31 The generic TVGARCH-in-mean(p,q) is defined as:

$$r_t = \lambda_t \sigma_t + \epsilon_t, \quad (2)$$

$$\epsilon_t = \sigma_t \eta_t, \quad (3)$$

$$\sigma_t^2 = \omega + \sum_{i=1}^p \alpha_i \epsilon_{t-i}^2 + \sum_{i=1}^q \beta_i \sigma_{t-i}^2, \quad (4)$$

$$\epsilon_t^2 = \psi_0 + u_t + \sum_{i=1}^{\infty} \psi_i u_{t-i}, \quad (5)$$

32 where η_t is an independent and identically distributed (*iid*) zero mean process
 33 with unit variance; σ_t is a latent conditional standard deviation; (5) is the
 34 $MA(\infty)$ representation of the conditional variance equation; $u_t = \epsilon_t^2 - \sigma_t^2$ is a
 35 martingale difference sequence process; $\phi = (\omega, \alpha_1, \dots, \alpha_p, \beta_1, \dots, \beta_q)'$ collects the
 36 free parameters in (4); and $\psi_i := \varrho_i(\phi)$ $i = 1, 2, \dots$ are deterministic functions

of the elements in ϕ . Similarly as in Giraitis, Kapetanios, and Yates (2013), the time-varying risk premium parameters are assumed to evolve smoothly over time, so that it satisfies a local stability condition in the form of $\sup_{s: \|s-t\| \leq h} \|\lambda_t + \lambda_s\|_2 = O_p(h/t)$.

Estimating the free parameters in (2) and (4) by maximum-likelihood is not a feasible alternative, as the class of TVGARCH-in-mean(p,q) models involves two unobserved processes: λ_t and ϵ_t . Anyfantaki and Demos (2016) address this issue in the context of the time-varying EGARCH(1,1)-in-mean model. Specifically, their work differs from mine in two ways. First, they parameterize the conditional variance as an EGARCH(1,1) model and, most importantly, λ_t as a stationary AR(1) process. By contrast, λ_t in (2) is assumed to satisfy $\sup_{s: \|s-t\| \leq h} \|\lambda_t + \lambda_s\|_2 = O_p(h/t)$, which encompasses the case of the driftless random walk process considered in Chou, Engle, and Kane (1992). Second, while I propose a kernel-based nonparametric method to estimate the time-varying risk premium parameter, Anyfantaki and Demos's (2016) estimation strategy is based on Bayesian methods (Markov chain Monte Carlo (MCMC) likelihood based estimation procedure).

I combine Linton and Perron's (2003) iterative semiparametric estimator with Giraitis, Kapetanios, and Yates's (2013) kernel-based least squares framework to estimate the free parameters $\theta = (\underline{\lambda}, \phi)'$, where $\underline{\lambda} = (\lambda_1, \dots, \lambda_T)'$. This method consists of recursively updating estimates of σ_t and u_t on each iteration, and then computing estimates of $\underline{\lambda}$ and ϕ . To this end, consider moment conditions based on (2) and (5),

$$\mathbb{E}[\sigma_t(r_t - \lambda_t \sigma_t)] = 0, \quad \text{for each } t = 1, 2, \dots, T, \quad (6)$$

$$\mathbb{E}[z_t u_t] = 0, \quad \text{with } z_t := \frac{\partial \left(\psi_0 + \sum_{i=1}^{\bar{q}} \psi_i u_{t-i} \right)}{\partial \phi}, \quad (7)$$

where (7) is truncated at some lag-order $\bar{q}$ with $\bar{q} > p + q + 1$. Notably, (7) holds because u_t is a martingale difference sequence and z_t is a function of lagged values of u_t . It follows that estimating θ by the standard generalized method of moments (GMM) using the moments defined in (6) and (7) is not operational, as z_t and σ_t are latent variables. Using Linton and Perron's (2003) approach, rewrite (6) and (7) using estimates of σ_t and u_t obtained at some j iteration,

$$\mathbb{E}[\sigma_{j,t}(r_t - \lambda_{j+1,t} \sigma_{j,t})] = 0, \quad \text{for each } t = 1, 2, \dots, T, \quad (8)$$

$$\mathbb{E}[z_{j,t} u_{j+1,t}] = 0, \quad (9)$$

where $\sigma_{j,t}$ and $z_{j,t}$ denote the filtered estimates of σ_t and z_t based on $\hat{\theta}_j$, and $u_{j+1,t} = \epsilon_{j,t}^2 - \psi_{j+1,0} - \sum_{i=1}^{\bar{q}} \psi_{j+1,i} u_{j,t-i}$ with $\epsilon_{j,t}^2 = (r_t - \lambda_{j+1,t} \sigma_{j,t})^2$. While the finite sample counterpart of (9) is given by the usual sample mean, computing the sample counterpart of (8) is less obvious. The work of Giraitis, Kapetanios, and Yates (2013) suggests the use of local kernels to construct operational

sample counterparts of (8). In turn, a feasible moment condition based on (8) reads

$$K_t^{-1} \sum_{\tau=1}^T k_{t,\tau} \sigma_{j,\tau} \left(r_\tau - \hat{\lambda}_{j+1,t} \sigma_{j,\tau} \right) = 0, \quad \text{for each } t = 1, 2, \dots, T, \quad (10)$$

where $k_{t,\tau} = K((t-\tau)/H)$ denotes a kernel function such that $K(x) \geq 0$ for any $x \in \mathbb{R}$ is a continuous bounded function with a bounded first derivative and $\int K(x)dx = 1$; H is the bandwidth parameter satisfying $H = o(T/\ln(T))$ as $H \rightarrow \infty$; and $K_t = \sum_{\tau=1}^T k_{t,\tau}$. Notably, writing the moment conditions as in (10) is consistent with previous studies in the time-varying parameter literature which maximizes kernel weighted log-likelihood functions (see Robinson (1989), Giraitis, Kapetanios, Wetherilt, and Žikeš (2016), among others).

I use the fact that (10) is exactly identified for each t , and hence estimates of λ_t can be obtained independently of ϕ . In turn, estimates of θ are computed iteratively by a two-step procedure. The first step consists of solving (10) for each t , while the second step mimics the work of Linton and Perron (2003) and consists of estimating ϕ using the sample counterpart of (9). In practise, the kernel-based iterative estimator is as follows:

Step 1: Choose starting values $\hat{\lambda}_0$ and $\hat{\phi}_0$, such that $\hat{\phi}_0$ satisfies the second-order stationarity conditions of the GARCH(1,1) model. Using $\hat{\theta}_{0,t} = (\hat{\lambda}_0, \hat{\phi}_0)'$, compute recursively $\{\sigma_{0,t}^2\}_{t=1}^T$, and $\{u_{0,t}\}_{t=1}^T$ from (2)-(5).

Step 2: Given $\{\sigma_{0,t}^2\}_{t=1}^T$, calculate

$$\hat{\lambda}_{1,t} = \left(\sum_{\tau=1}^T k_{t,\tau} \sigma_{0,\tau}^2 \right)^{-1} \sum_{\tau=1}^T k_{t,\tau} \sigma_{0,\tau} r_\tau, \quad \text{for each } t = 1, 2, \dots, T. \quad (11)$$

Step 3: Solving the sample counterpart of (9) is equivalent to estimate $\hat{\phi}_1$ by nonlinear least squares. Calculate

$$\hat{\phi}_1 = \arg \min_{\hat{\phi}_1} \sum_{t=1}^T \left\{ \left(r_t - \hat{\lambda}_{1,t} \sigma_{0,t} \right)^2 - \hat{\psi}_{1,0} - \sum_{i=0}^{\bar{q}} \hat{\psi}_{1,i} u_{0,t-1-i} \right\}^2. \quad (12)$$

Step 4: Update recursively $\{\sigma_{1,t}^2\}_{t=1}^T$ and $\{u_{1,t}\}_{t=1}^T$ based on $\hat{\theta}_1$.

Repeat steps 2-4 j times until $\hat{\theta}_j$ converges. Convergence occurs when $\|\hat{\lambda}_j - \hat{\lambda}_{j-1}\|_2 \leq \varepsilon$ and $\|\hat{\phi}_j - \hat{\phi}_{j-1}\|_2 \leq \varepsilon$, with ε set to 10^{-5} . Parameters on

the j^{th} iteration are given by:

$$\hat{\lambda}_{j,t} = \left[\sum_{\tau=1}^T k_{t,\tau} \sigma_{j-1,\tau}^2 \right]^{-1} \sum_{\tau=1}^T k_{t,\tau} \sigma_{j-1,\tau} r_{\tau}, \quad \text{for each } t = 1, 2, \dots, T, \quad (13)$$

$$\hat{\phi}_j = \arg \min_{\hat{\phi}_j} \sum_{t=1}^T \left[r_t - \hat{\lambda}_{j,t} \sigma_{j-1,t} \right]^2 - \hat{\psi}_{j,0} - \sum_{i=0}^{\bar{q}} \hat{\psi}_{j,i} u_{j-1,t-1-i} \Bigg]^2. \quad (14)$$

Finally, three inputs are still necessary to implement the above algorithm: the kernel function, the bandwidth parameter H , and the truncation lag $\bar{q}$. As in Giraitis, Kapetanios, and Yates (2013), three kernel functions are used: the Epanechnikov, Gaussian, and flat kernels. The choice of H follows from the Monte Carlo study conducted in Section 3.1, while $\bar{q}$ is chosen to be proportional to $\ln(T)$, (Dufour and Jouini (2005)).

Asymptotic theory for the Quasi-Maximum Likelihood (QMLE) estimator in the GARCH-in-mean models is yet to be fully established. Conrad and Mammen (2016) give an important step forward and prove the asymptotic distribution of the QMLE estimator for the simple GARCH(1,1)-in-mean. As discussed in Linton and Perron (2003), the semiparametric GARCH-in-mean models offer additional complications compared to the standard GARCH-in-mean models, and hence rigorous inference is still not available. Similar difficulties arise in the TVGARCH-in-mean specification. In turn, this note follows Linton and Perron's (2003) approach as it briefly discusses the general conditions required for consistency and asymptotic normality; uses the wild bootstrap to conduct inference; and adopts a Monte Carlo study to assess the finite sample properties of the proposed iterative estimator.

The concept of asymptotic contraction mapping (ACM) developed in Dominitz and Sherman (2005) is useful to guide the discussion on the asymptotic properties of the kernel iterative estimator. If a collection is an ACM, then it will have a unique fixed point that depends on the sample characteristics and hence the iterative procedure converges.² While the two-step procedure given in (13) and (14) is seen as the sample mapping, (8) and (9) are their population counterpart. Consistency and asymptotic normality require the population mapping to be an ACM, which implies, under some uniform convergence conditions, that the sample mapping is also an ACM and hence has an unique fixed point (regardless of the initial values). Combining Theorem 4 in Dominitz and Sherman (2005) with Giraitis, Kapetanios, and Yates's (2013) results, estimates of λ_t are expected to be $\sqrt{H}$ consistent and asymptotically normally distributed, and estimates of ϕ are expected to be consistent and asymptotically normally distributed at the usual $\sqrt{T}$ rate.

²See Dominitz and Sherman, 2005, p. 841 for a formal definition of the ACM.

128 3. Numerical illustrations

129 3.1. Monte Carlo

130 I simulate data from (2)-(4) where $p = q = 1$, η_t is normally distributed with
131 zero mean and variance equal to one, and λ_t follows a bounded random walk
132 process (see detailed discussion in the online Supplement). The sample size and
133 the number of replications are set to 2,000 and 1,000, respectively.

134 Table 1 displays the results for the kernel-based iterative estimator computed
135 with alternative bandwidth choices and the Epanechnikov, Gaussian, and flat
136 kernels. Results are reported in terms of the root mean squared error (RMSE)
137 and pointwise correlation between λ_t and the kernel-based estimates. The best
138 choices of bandwidth parameters, in terms of minimizing the RMSE, are $H =$
139 $T^{1/2}$ and $H = T^{6/10}$. These are also the bandwidths that deliver the highest
140 pointwise correlation (about 0.85) between the kernel-based estimates and the
141 true latent time-varying risk premium parameter. All combinations of kernel
142 methods and bandwidth parameters deliver unbiased estimates of $\phi = (\omega, \alpha, \beta)'$
143 (apart from $H = T^{2/10}$). Finally, convergence rates are greater than 98% for all
144 specifications, suggesting that (8) and (9) are ACMs.

145 3.2. Empirical results

146 I use excess returns of the CRSP value-weighted index aggregated on a
147 monthly basis. Figure 1 plots monthly estimates of λ_t and their 90% wild
148 bootstrap confidence bands from a TVGARCH(1,1)-in-mean model with band-
149 width $H = T^{6/10}$ (see the Supplement material for a Monte Carlo study showing
150 that the wild bootstrap produces valid inference). Not surprisingly, likewise the
151 semiparametric GARCH-in-mean models, the empirical confidence bands are
152 relatively wide, which reflects the difficulties associated with estimating the risk-
153 return trade-off (Linton and Perron (2003) and Christensen, Dahl, and Iglesias
154 (2012)). I find that the risk premium parameter is indeed time-varying, with $\hat{\lambda}_t$
155 assuming both positive (generally significant) and negative (insignificant) val-
156 ues. This finding sheds light on the mixed evidence on the risk-return literature
157 regarding the sign and significance of the risk premium parameter. Addition-
158 ally, in periods where $\hat{\lambda}_t$ is statistically significant, market volatility is low. On
159 contrary, when $\hat{\lambda}_t$ is not statistically significant, market volatility is high. This
160 indicates that identification of the risk premium parameter is problematic in
161 periods of high volatility (Rossi and Timmermann (2010)).

162 4. Conclusion

163 I introduce a kernel-based iterative estimator that combines the estimators in
164 Giraitis, Kapetanios, and Yates (2013) and Linton and Perron (2003) to estimate
165 the stochastic time-varying risk premium parameter in the TVGARCH(1,1)-in-
166 mean model. The Monte Carlo study shows that the kernel-based estimator
167 presents a good finite sample performance. I investigate the time-varying risk
168 premium for the CRSP index and find strong evidence that λ_t is indeed time-
169 varying.

- 170 ANYFANTAKI, S., AND A. DEMOS (2016): “Estimation and Properties of a
171 Time-Varying EGARCH(1,1) in Mean Model,” *Econometric Reviews*, 35(2),
172 293–310.
- 173 BRANDT, M. W., AND L. WANG (2010): “Measuring the Time-Varying Risk-
174 Return Relation from the Cross-Section of Equity Returns,” Discussion paper,
175 Duke University.
- 176 CHOU, R., R. F. ENGLE, AND A. KANE (1992): “Measuring risk aversion from
177 excess returns on a stock index,” *Journal of Econometrics*, 52(1), 201 – 224.
- 178 CHRISTENSEN, B. J., C. M. DAHL, AND E. M. IGLESIAS (2012): “Semipara-
179 metric inference in a GARCH-in-mean model,” *Journal of Econometrics*, 167,
180 458–472.
- 181 CONRAD, C., AND E. MAMMEN (2016): “Asymptotics for parametric GARCH-
182 in-Mean models,” *Journal of Econometrics*, 194(2), 319 – 329, Financial
183 Statistics and Risk Management.
- 184 DOMINITZ, J., AND R. SHERMAN (2005): “Some Convergence Theory for Iter-
185 ative Estimation Procedures with an Application to Semiparametric Estima-
186 tion,” *Econometric Theory*, 21, 838–863.
- 187 DUFOUR, J.-M., AND T. JOUINI (2005): “Asymptotic Distribution of a Simple
188 Linear Estimator for Varma Models in Echelon Form,” in *Statistical Mod-
189 eling and Analysis for Complex Data Problems*, ed. by P. Duchesne, and
190 B. Rémillard, pp. 209–240. Springer US.
- 191 GIRAITIS, L., G. KAPETANIOS, A. WETHERILT, AND F. ŽIKEŠ (2016): “Es-
192 timating the Dynamics and Persistence of Financial Networks, with an Ap-
193 plication to the Sterling Money Market,” *Journal of Applied Econometrics*,
194 31(1), 58–84, jae.2457.
- 195 GIRAITIS, L., G. KAPETANIOS, AND T. YATES (2013): “Inference on stochastic
196 time-varying coefficient models,” *Journal of Econometrics*.
- 197 LINTON, O., AND B. PERRON (2003): “The Shape of the Risk Premium: Ev-
198 idence From a Semiparametric Generalised Autoregressive Conditional Het-
199 eroscedasticity Model,” *Journal of Business & Economic Statistics*, 21, 354–
200 367.
- 201 MERTON, R. C. (1973): “An Intertemporal Capital Asset Pricing Model,”
202 *Econometrica*, 41(5), 867–887.
- 203 ROBINSON, P. (1989): “Nonparametric Estimation of Time-Varying Param-
204 eters,” in *Statistics Analysis and Forecasting of Economic Structural Change*,
205 ed. by P. Hackl, chap. 15, pp. 253–264. Springer, Berlin.
- 206 ROSSI, A., AND A. TIMMERMANN (2010): “What is the Shape of the Risk-
207 Return Relation?,” Discussion paper, The Rady School of Management, Uni-
208 versity of California, San Diego.

Table 1: Finite sample performance of the kernel-based iterative estimator

Bandwidth - H	Kernel	λ_t		RMSE			Mean		
		RMSE	Corr	ω	α	β	ω	α	β
$T^{2/10}$	Epanechnikov	0.39	0.60	0.01	0.03	0.10	0.02	0.07	0.84
$T^{2/10}$	Gaussian	0.25	0.71	0.01	0.02	0.06	0.01	0.05	0.88
$T^{2/10}$	Flat	0.36	0.61	0.01	0.02	0.07	0.01	0.06	0.86
$T^{3/10}$	Epanechnikov	0.26	0.72	0.01	0.02	0.05	0.01	0.05	0.88
$T^{3/10}$	Gaussian	0.18	0.81	0.01	0.02	0.05	0.01	0.05	0.89
$T^{3/10}$	Flat	0.24	0.73	0.01	0.02	0.05	0.01	0.05	0.89
$T^{4/10}$	Epanechnikov	0.18	0.80	0.01	0.02	0.05	0.01	0.05	0.90
$T^{4/10}$	Gaussian	0.13	0.86	0.01	0.02	0.06	0.01	0.05	0.90
$T^{4/10}$	Flat	0.17	0.81	0.01	0.02	0.05	0.01	0.05	0.90
$T^{5/10}$	Epanechnikov	0.13	0.86	0.01	0.02	0.05	0.01	0.05	0.90
$T^{5/10}$	Gaussian	0.11	0.88	0.01	0.02	0.05	0.01	0.05	0.90
$T^{5/10}$	Flat	0.13	0.86	0.01	0.02	0.05	0.01	0.05	0.90
$T^{6/10}$	Epanechnikov	0.12	0.88	0.01	0.02	0.06	0.01	0.05	0.89
$T^{6/10}$	Gaussian	0.13	0.86	0.01	0.02	0.06	0.01	0.05	0.89
$T^{6/10}$	Flat	0.12	0.86	0.01	0.02	0.06	0.01	0.05	0.89
$T^{7/10}$	Epanechnikov	0.13	0.85	0.01	0.02	0.05	0.01	0.05	0.90
$T^{7/10}$	Gaussian	0.16	0.79	0.01	0.02	0.05	0.01	0.05	0.90
$T^{7/10}$	Flat	0.15	0.81	0.01	0.02	0.05	0.01	0.05	0.90
$T^{8/10}$	Epanechnikov	0.17	0.77	0.01	0.02	0.05	0.01	0.05	0.89
$T^{8/10}$	Gaussian	0.21	0.68	0.01	0.02	0.05	0.01	0.05	0.89
$T^{8/10}$	Flat	0.19	0.67	0.01	0.02	0.05	0.01	0.05	0.89

RMSE accounts for root mean squared error; Corr is the pointwise correlation between λ_t and $\hat{\lambda}_t$; λ_t is defined as a bounded random walk process with upper and lower bounds given by 0.90 and -0.90, respectively; and $\phi = (0.01, 0.05, 0.9)'$.

Figure 1: Time-varying risk premium estimation and conditional standard deviation

The left-hand-side, center and right-hand-side graphs display estimates of λ_t with the Epanechnikov, Gaussian, and the flat kernel functions, respectively. Estimates of λ_t and the 90% confidence intervals are on the left axis. The bandwidth parameter is equal to $H = T^{6/10}$. The conditional standard deviation is in light blue on the right axis. I perform 1000 replications in the bootstrap algorithm. The shaded areas account for the recession periods from the National Bureau of Economic Research (NBER).