Organic Aerosols in Anoxic and Oxidic Atmospheres of Earth-like Exoplanets: VUV-MIR Spectroscopy of CHON Tholins

Liseth Gavilan¹, Nathalie Carrasco¹, Søren Vrønning Hoffmann², Nykola C. Jones², and Nigel J. Mason³

¹ LATMOS, Université Versailles St. Quentin, UPMC Université Paris 06, CNRS, 11 Blvd. d'Alembert, F-78280 Guyancourt, France; liseth.gavilan@gmail.com

² ISA, Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, DK-8000 Aarhus C, Denmark

³ Department of Physical Sciences, The Open University, Milton Keynes MK7 6AA, UK

Received 2018 February 5; revised 2018 May 4; accepted 2018 May 28; published 2018 July 11

Abstract

Organic hazes can drastically impact the characterization and detection of exoplanet atmospheres. We explore the effects of the transition from an anoxic to oxidic atmosphere on the molecular structure of organic aerosols, by producing aerosols in N₂:CO₂:CH₄-rich plasmas with increasing CO₂/CH₄ gas ratios. We performed a spectroscopic study on the resulting aerosol analogs, CHON-rich tholins, from the vacuum-ultraviolet (VUV) to the mid-infrared (MIR), i.e., the 0.13–10 μm range. VUV spectroscopy revealed the presence of π - π^* and n - π^* transitions in the 200–500 nm range in all samples. These are attributed to electronic transitions in amine groups. As the CO₂/CH₄ ratio increases, new bands emerge that can be attributed to electronic transitions in hydroxyl and carboxyl functional groups. MIR spectroscopy showed that the molecular structure of oxidized aerosols is dominated by the CO₂/CH₄ ratio. Band deconvolution revealed the oxygenation of the organic matrix via the formation of oxygenated functional groups, including amide, hydroxyl, and carbonyl groups. For the most oxidized aerosols, absorption is greatest in the 0.13–0.3 μm and 6–10 μm regions. Some of the oxidized samples were further irradiated with VUV photons, resulting in the formation of π - π^* bands in the 200–250 nm region and a decrease in the band intensities below 200 nm, attributed to amine and nitrile losses. We discuss how oxidized aerosols could provide a detectable (bio-)signature in exoplanet atmospheres.

Key words: planets and satellites: – fundamental parameters – methods: laboratory: solid state – planets and satellites: atmospheres – techniques: spectroscopic

1. Introduction

Photochemical aerosols are present across the atmospheres of several bodies in the solar system. These can be observed as hazes from the stratosphere to the ionosphere. As such, and depending on their optical properties, morphology and size distribution, they could heavily impact the radiative transfer of solar radiation and planetary surface temperature. One of the most notable examples of an atmosphere dominated by photochemical aerosols is Titan. Its quasi-reducing atmospheric chemistry is the result of photon and energetic particles (protons, cosmic rays) impact on N₂ and CH₄, an efficient gas mixture for the generation of organic photochemical aerosols (Wilson & Atreya 2004; Lavvas et al. 2013).

On Earth, organic aerosols are presently one of the largest drivers of radiative forcing, affecting the Earth's thermal balance and providing a mechanism for climate change (Jacobson et al. 2000). In the Archean Earth (~4–2.5 Gyr ago), the presence of photochemical aerosols could have influenced early planetary habitability via radiative shielding and surface cooling (Pavlov et al. 2001). Unlike Titan's anoxic atmosphere, CO₂ is expected to have been present in the atmosphere of Earth from its early stages following the emergence of life (Kasting 1993; von Paris et al. 2008; Feulner 2012). Atmospheric oxidation therefore probably impacted the formation and optical properties of early organic hazes (Haqq-Misra et al. 2008; Wolf & Toon 2010).

Aerosols are likely to be ubiquitous in the atmospheres of extrasolar planets (Marley et al. 2013) and their composition can be as widely varying as that of the exoplanetary atmospheres in which they are formed (Lavvas & Koskinen 2017). Methane was detected in the atmosphere of exoplanet HD 189733b

(Swain et al. 2008) with CO, CO₂, and H₂O (Swain et al. 2009b). Swain et al. (2009a) detected H₂O, CH₄, and CO₂ in the near-infrared spectrum of exoplanet HD 209458b. CO₂ photo-dissociation is relevant to the radiative transfer in exoplanet atmospheres (Venot et al. 2018). Its presence could also affect the photochemical production of organic hazes in Earth-like exoplanets as shown by models (Arney et al. 2017) and experiments (Trainer et al. 2006; Fleury et al. 2017; Gavilan et al. 2017).

While the optical properties of organic aerosols are needed to refine models of the Archean Earth and exoplanet atmospheres, their molecular properties are necessary to understand the spectral variations as a function of the degree of oxidation. The aim of this work is to experimentally simulate the composition of increasingly oxidizing atmospheres to understand the impact of CO₂ in the optical properties of the generated organic aerosols covering a wide spectral range, from the vacuum-ultraviolet to the mid-infrared. This study is a continuation of our first study on the UV–visible complex refractive indices of oxidized organic aerosols (Gavilan et al. 2017) but here we focus our study on the molecular, electronic, and absorption properties of these aerosols.

2. Experiments

Tholins were produced using the PAMPRE⁴ experimental facility, a low-pressure (0.95 mbar) radiofrequency (RF) plasma reactor (Szopa et al. 2006). This apparatus allows the simulation of the complex organic chemistry initiated by energetic particles in planetary ionospheres. The 13.56 MHz

⁴ PAMPRE: Production d'Aérosols en Microgravité par Plasmas REactifs.

Table 1Synthesis Parameters for Tholins Prepared at Increasing CO₂/CH₄ Ratios, Thickness, and Tauc Gap Energies

Gas mixture N ₂ :CO ₂ :CH ₄	CO ₂ /CH ₄	Deposit (s)	Thickness (nm)	E_g (eV)
95:0:5	0	3	36 ± 7	2.35 ± 0.20
95:2.5:2.5	1	15	79 ± 16	2.54 ± 0.40
90:5:5	1	20	79 ± 16	2.49 ± 0.40
95:3:2	1.5	30	195 ± 15	3.06 ± 0.40
90:6:4	1.5	60	74 ± 40	2.66 ± 0.40
95:4:1	4	600	43 ± 21	3.20 ± 0.20
90:8:2	4	480	33 ± 16	3.23 ± 0.20

RF source, set to 30 W, generates a capacitively coupled plasma fed by a gas mixture at 0.95 mbar. The aerosol analogs presented in this study were prepared in a cold plasma where neutrals have a temperature of ~ 340 K. However, the most observed exoplanets to date are hot giants (~ 800 – 3000 K) and are particularly conducive to atmospheric observations (Madhusudhan et al. 2016). Future laboratory simulations under different temperatures and pressures should explore how these parameters could further impact the formation, composition, and optical properties of aerosols. This paper is a first step in the investigation of the molecular properties of oxidized organic aerosols.

Initially, a gas mixture was chosen to simulate the reducing composition of Titan’s atmosphere, i.e., N₂:CH₄ with a 95:5 ratio (e.g., Mahjoub et al. 2012; Sciamma-O’Brien et al. 2012). We then simulated increasingly oxidizing atmospheres by adding CO₂ to the N₂–CH₄ gas mixture. The resulting tholins, solids rich in C, H, O, and N (CHON-rich), are produced on substrates placed on the bottom electrode of the capacitor producing the plasma. One set of samples was prepared with a gas mixture, keeping the N₂ partial pressure at 90%, while a second set of samples was prepared keeping the N₂ partial pressure at 95%. For both series, gas mixtures with CO₂:CH₄ ratios equal to 0, 1, 1.5, and 4 were used. We can translate the CO₂:CH₄ ratios into C/O ratios, used to characterize exoplanet atmospheres in terms of their carbon content (e.g., Venot et al. 2015),

$$\frac{C}{O} = \frac{1 \times N_{CO_2} + 1 \times N_{CH_4}}{2 \times N_{CO_2}} = 0.5 + \frac{0.5}{CO_2:CH_4}. \quad (1)$$

Using this conversion, we find C/O ratios equal to ∞ , 1, 0.83, and 0.68. According to Venot et al. (2015), this C/O range includes C-rich (C/O = ∞ and 1) and near-solar (C/O = 0.68) type atmospheres.

The gas mixtures and deposition times of tholins are shown in Table 1. Although the deposition timescales differ by a factor of 100 between experiments, we have striven to obtain a film thickness leading to sufficient S/N in transmission over the VUV–MIR range. Additional effects like sputtering and electron impact from longer exposures to the plasma could further affect the properties of such films, and this will be the object of further studies. During deposition, bare silicon (10 × 10 mm) and MgF₂ windows (20 mm diameter) were placed on the bottom electrode within the PAMPRE chamber. The thickness of the deposited films is estimated from ellipsometry on the Si substrates, and the method is described in Gavilan et al. (2017). In this paper, we present the result of transmission measurements performed on the films deposited

on MgF₂ windows. Synchrotron and benchtop spectrophotometers were used to measure the transmission spectra of these films in the vacuum-ultraviolet (VUV; 130–250 nm), ultraviolet-visible (UV–Vis; 210–1000 nm), near-infrared (NIR; 1–3 μ m), and mid-infrared (MIR; 1.43–10 μ m) spectral ranges. These are described in more detail below.

Although the film thickness can vary from one substrate to the other during plasma deposition, we estimate that this variation does not exceed 30% of the thickness measured on the films deposited on Si. The estimated thickness of each film is reported in Table 1. According to Arney et al. (2017), the production of organic hazes decreases when CO₂:CH₄ > 10. In our study, the aerosol production rate is drastically quenched at a CO₂:CH₄ ratio of 4.

VUV transmission spectra were measured on the AU-UV beam line at the ASTRID⁵ synchrotron at Aarhus University, Denmark (Palmer et al. 2015). Thin films deposited on MgF₂ windows were placed in the *lollipop* sample holder, able to mount up to four samples. UV–Vis transmission spectra were measured with a bench-top UV–Vis spectrometer (*Thermo Scientific Evolution 300*). The overlapping spectral range was used to combine the VUV and UV–Vis data sets. One sample, the fully reduced tholin, had previously been measured in the APEX⁶ chamber of the DISCO⁷ beam line of the SOLEIL⁸ synchrotron. We confirmed that VUV spectra are reproducible between beam lines at both facilities.

Near-infrared transmission spectra were measured at the Planetary Emissivity Laboratory (DLR) using a vacuum Fourier transform infrared (FTIR) spectrometer (Bruker Vertex 80V) with a liquid nitrogen cooled HgCdTe (mercury cadmium telluride, MCT) detector at 1 cm^{−1} spectral resolution. A Bruker A480 parallel beam accessory was mounted in front of the sample holder within the Vertex 80V to reduce artifacts due to beam reflections on the sample. This accessory helps collimate the beam when measuring highly refractive/reflective samples in transmission mode.

Mid-infrared transmission spectra were measured with a vacuum FTIR spectrometer (Bruker Vertex 80V) available at ASTRID2. The reduced tholin mid-infrared spectrum was measured on the iN10 infrared microscope available at SOLEIL. These were recorded at 4 cm^{−1} spectral resolution and were corrected with a polynomial baseline to subtract the continuum absorption.

3. VUV and MIR Spectroscopy

Our spectroscopic analysis involves deconvolution of the main electronic bands for the VUV–UV spectra and of the main vibrational modes in the MIR spectra. This is done via the Levenberg–Marquardt nonlinear least-squares minimization method using the IDL MPFIT routine (Markwardt 2009). We have chosen Gaussian band shapes, suitable for characterizing molecular transitions in solids. In fitting each band, the peak position and intensity are left as free parameters, while the width is varied up to 10% between spectra. Band positions and widths may shift depending on the bonding environment for each sample.

⁵ ASTRID: Aarhus STorage RIng in Denmark, <http://www.isa.au.dk>.

⁶ APEX: Atmospheric Pressure Experiment.

⁷ DISCO: Dichroism, Imaging and Spectroscopy.

⁸ SOLEIL: Source Optimisée de Lumière d’Énergie Intermédiaire du LURE.

Figure 1. Vis-VUV spectra of tholins (normalized to the maximum optical depth in this range), prepared in plasmas of increasing CO_2/CH_4 ratios. Due to the large thickness ($d \sim 195$ nm), the VUV spectrum of the 95:3:2 tholin (green curve) is saturated for wavenumbers larger than $6 \mu\text{m}^{-1}$.

3.1. VUV-Visible Spectroscopy

The VUV-visible spectra of the oxidized tholins have been normalized to the maximum optical depth in the VUV. These are shown in Figure 1. We notice that increasing the CO_2/CH_4 ratio has a direct effect on the electronic absorption bands of these tholins. This evolution is best represented by the appearance of a narrow but intense peak at ~ 180 nm and a decrease in the intensity of the far-UV (FUV) band. Varying the N_2 partial pressure from 90% to 95% has a negligible effect on the VUV spectra of tholins prepared with similar CO_2/CH_4 ratios.

We used the visible-near-UV spectral range (250–1000 nm) to determine the optical band gap of these materials, also known as the Tauc gap (Tauc et al. 1966). The gap values were obtained by extrapolating a regression from the linear region in a $(\alpha \times E)^{1/2}$ versus E plot, where α is the absorption coefficient and E is in eV. The Tauc extrapolation is found using the linear region between the visible and near-ultraviolet range, i.e., 3.8–04.5 eV (see Figure 2). This region avoids the exponential absorption tail appearing in the visible energy range (1.2–3.8 eV), characteristic of disorder in amorphous materials, known as the Urbach tail (Urbach 1953; Jones 2012). We varied the endpoints of this range to determine the average systematic error in the gap determination. The gap values are listed in Table 1. We note that, as the CO_2/CH_4 ratio increases, so does the gap value, i.e., E_g increases from 2.35 to 3.20 eV. These gap values are consistent (within the error bars) with those found via the ellipsometry on similar films deposited on Si substrates (Gavilan et al. 2017), validating the coherence of both methods.

As seen in Figure 1, the VUV-visible spectral profile for samples prepared with the same CO_2/CH_4 ratio are similar, e.g., 95:4:1 and 90:8:2 both with a ratio of 4 and 95:2.5:2.5 and 90:5:5 both with a ratio of 1. Because the spectra of the 95:3:2 sample is optically saturated in the VUV, we chose the 90% N_2 series to deconvolve the VUV-vis spectra for increasing CO_2/CH_4 ratios = 0, 1, 1.5, 4. The resulting deconvolution is shown in Figure 3.

VUV-visible spectra reveal transitions of electrons between the molecular orbitals of organic materials. Identification of possible carriers of the electronic bands in our tholins is done

Figure 2. Tauc optical gap determination from the visible to near-ultraviolet spectra for tholins prepared in plasmas with increasing CO_2/CH_4 ratios in $\text{N}_2:\text{CO}_2:\text{CH}_4$ mixtures.

by comparison to UV-Vis gas phase spectra of common chromophores present in organic molecules (e.g., Hornback 2005; Workman 2016). These are listed in Table 2. The major electronic transitions in organic compounds are $\sigma \rightarrow \sigma^*$, $\pi \rightarrow \pi^*$, $n \rightarrow \sigma^*$, and $n \rightarrow \pi^*$, and associated bands. Generally π -electronic transitions are found at $1.6 < \lambda^{-1} < 5 \mu\text{m}^{-1}$ ($200 < \lambda < 600$ nm), while σ -electronic transitions are found at higher energies, $\lambda^{-1} > 3.8 \mu\text{m}^{-1}$ ($\lambda < 260$ nm) (Gadallah et al. 2011; Gavilan et al. 2016).

For our sample spectra, each VUV-Vis spectrum can be deconvolved into three main bands. However, as the oxidation degree increases, band and peak positions are modified. For the tholin with no CO_2 , two bands, which we call G1 and G2, are attributed to transitions in amine functional groups. G1 and G2 are assigned to electronic transitions from the lone-pair electrons on nitrogen to nitrogen atomic-like orbitals (Hubin-Franskin et al. 2002; Duarte et al. 2013). The largest band, called G3, found in the far-UV, is generally attributed to $\sigma-\sigma^*$ and $n-\sigma^*$ type transitions and its peak position is found at $8 < \lambda^{-1} < 8.5 \mu\text{m}^{-1}$ (118–125 nm).

The VUV spectra evolve as the oxidation degree increases. The spectra for samples with $\text{CO}_2/\text{CH}_4 = 1$ and 1.5 are remarkably similar. The smaller UV bands have shifted, which signals the molecular evolution of the solid. G4 is then attributed to low energy $\pi-\pi^*$ and $n-\pi^*$ transitions arising from structural disorder (Gadallah et al. 2011). G5 is a new VUV band centered at $\lambda^{-1} \sim 5.4 \mu\text{m}^{-1}$ (185 nm), whose intensity increases with increasing oxidation degree, and is attributed to electronic transitions in hydroxyl groups due to the promotion of an electron from the highest filled molecular orbital to an antibonding orbital (σ_{OH}^*) (Duarte et al. 2013). This band usually appears at $5 < \lambda^{-1} < 6.1 \mu\text{m}^{-1}$ (165–200 nm) (Xu et al. 1999).

Finally, for the most oxidized sample, with $\text{CO}_2/\text{CH}_4 = 4$, the G5 band is most prominent as a consequence of the larger number of oxygenated bonds. G6 is a new band appearing at $\lambda^{-1} \sim 4.2 \mu\text{m}^{-1}$ (240 nm) and is assigned to the $n_{\text{O}} \rightarrow \pi_{\text{CO}}^*$ transitions from the lone-pair orbital on the carbonyl oxygen to the antibonding π_{CO} valence orbital (Barnes & Simpson 1963; Ari & Güven 2000) or to the valence shell excitations of hydroxyl groups (Duarte et al. 2013). For this sample, the

Figure 3. Spectral deconvolution of the Vis-VUV of increasingly oxidized organic tholins prepared in plasmas of increasing CO_2/CH_4 ratios.

position of G3 is shifted to lower energies and its relative intensity decreases, indicating a loss of σ -bonds.

3.2. Mid-infrared Spectroscopy

The mid-infrared spectra of our samples were split into three spectral ranges corresponding to the most intense vibrational

Table 2
VUV Electronic Band Assignments

Label	Peak Pos. nm/eV	Literature Position (eV)	Electronic Transition	Functional Group
G1	215/5.77	5.70, ^a 5.77 ^b	$n_N \rightarrow 3sa$	Amine
G2	166/7.47	7.14, ^b 7.93 ^c	$n_N \rightarrow 3pa$	Amine
G3	118/10.51	10.16 ^{d-12}	$\sigma \rightarrow \sigma^*$, $n \rightarrow \sigma^*$	Methyl
G4	300/4.13	3.50 ^f –4.75 ^g	$\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$	Carbonyl
G5	185/6.7	6.2–7.5 ^h	$n_O \rightarrow \sigma^*$	Hydroxyl
G6	241/5.15	5.80 ⁱ	$n_O \rightarrow \pi_{CO}^*$	Carboxyl

Notes.

^a Hubin-Franskin et al. (2002).

^b Tannenbaum et al. (1953).

^c Harshbarger (1971).

^d Workman (2016).

^e Gavilan et al. (2016).

^f Hornback (2005).

^g Gadallah et al. (2011); Robertson & O'Reilly (1987).

^h Xu et al. (1999).

ⁱ Barnes & Simpson (1963); Ari & Güven (2000).

Table 3
Mid-infrared Band Assignments for Tholins Prepared at
Increasing CO_2/CH_4 Ratios

Peak (cm^{-1})	Range (cm^{-1})	Band Assignments ^{a,b}
3320	3500–3100	NH/NH ₂ stretch (Amine I)
3200	...	NH stretch (Amine I)
3175	...	NH stretch (Amide I)
2960	3000–2800	CH stretch (ali. CH_2/CH_3 , arom. =CH)
2846	2900–2800	aldehyde CH stretch ($\text{H}-(\text{R})\text{C}=\text{O}$)
2240	2260–2230	R–C≡N, nitrile stretch
2180	2230–2150	C=C–CN conjugated nitrile stretch ^c
2145	2180–2120	R–N≡C, isonitrile stretch ^d
2110	2120–2100	aromatic –N≡C stretch
1700	1730–1680	C=O stretch Also: carbonyl (aldehyde, ketone), carboxyl
1625	1650–1620	C=N stretch Also: C=O, C=C stretch
1560	1570–1540	C=C stretch Also: C=N, quinones, carboxylates, NO
1460	1480–1440	$\text{CH}_{2,3}$, NH ₂ , C–OH bending
1350	1370–1340	$\text{CH}_{2,3}$ bending Also: aromatic C–N bending, N–O stretch
1230	1250–1220	C–O stretch Also: C–OH stretch, ester

Notes.

^a <http://webbook.nist.gov/chemistry/>

^b <https://webspectra.chem.ucla.edu/irttable.html>

^c Imanaka et al. (2004).

^d Mutsukura (1999).

bands in the mid-infrared: (a) amines ($3800\text{--}2400\text{cm}^{-1}$), (b) nitriles ($2300\text{--}2050\text{cm}^{-1}$), and a (c) hetero-aromatics band ($1900\text{--}1000\text{cm}^{-1}$) (Mutsukura 1999; Imanaka et al. 2004). The assignment of these modes are provide in Table 3. We have deconvolved the main mid-infrared spectral bands of all samples prepared with a 95% N_2 concentration, corresponding to the following $\text{N}_2:\text{CO}_2:\text{CH}_4$ mixtures: 95:0:5, 95:2.5:2.5,

Figure 4. Spectral deconvolution of the main mid-infrared spectral bands (amines, nitriles, hetero-aromatics spanning 3800–1000 cm^{-1}) of increasingly oxidized organic tholins. To illustrate the effect of the increasing CO_2/CH_4 ratio on the molecular structure, only tholins from the same series (95% N_2) and varying CO_2/CH_4 ratios = 0, 1, 1.5, and 4, are used.

95:3:2, and 95:4:1, plotted in Figure 4. Such deconvolution provides further insight into the evolution of the molecular structure of organic aerosols as a function of their oxidation degree. In this first-order analysis, we use Gaussian functions to fit the most intense vibrational modes due to broad bonding distributions in the solid network that result in statistical bell curves, and neglect any perturbations to the line shape due to secondary resonances. Fitting vibrational bands is done by using the number of Gaussians needed to characterize the vibrational spectra of the $\text{CO}_2/\text{CH}_4 = 0$ tholin, which has the largest number of peaks. To fit the tholin spectra at different oxidation degrees, we have assumed that (1) the organic network of tholins may be similar to first order and (2) oxidation will produce new bonds that may shift the energies and abundances of functional groups.

For tholins prepared with increasing CO_2/CH_4 ratios, two main features appear in the amine band. These are attributed to the stretching modes of NH functions in amide groups at

$\sim 3180 \text{ cm}^{-1}$ and to aldehyde CH stretching modes at $\sim 2800 \text{ cm}^{-1}$. In the nitrile band, increasing oxidation translates into a decrease of singly bonded and aromatic isonitriles at $\sim 2140 \text{ cm}^{-1}$ and $\sim 2105 \text{ cm}^{-1}$, respectively, and to the decrease of singly bonded nitrile modes at $\sim 2240 \text{ cm}^{-1}$. In contrast at CO_2/CH_4 ratios greater than 1, we note the effect of increasing oxidation on the heteroaromatic band: the unsaturated $\text{C}=\text{C}$ and $\text{C}=\text{N}$ modes decrease while the intensity of the carbonyl ($\text{C}=\text{O}$) band at 1700 cm^{-1} increases. This is confirmed by the formation of a new band in the region between 1550 and 1300 cm^{-1} , attributed to bending modes in NH_2 , $\text{CH}_{2,3}$, and $\text{C}-\text{OH}$ functional groups.

From the deconvolution of the amines band an increasing CO_2/CH_4 ratio results in the rapid evolution of the NH stretching mode associated with amides and to the aldehyde CH stretch. Although the primary NH bonds associated with amines are present and common for all samples, in samples with a $\text{CO}_2/\text{CH}_4 = 4$ ratio, the NH band associated with

Figure 5. Imaginary refractive index k spanning the VUV–Vis, NIR, and MIR ranges for tholins prepared at increasing CO_2/CH_4 or C/O ratios. As oxidation increases, the absorption coefficient increases steeply in the ultraviolet range (130–300 nm), linearly in the mid-infrared, while the k inversion region is blueshifted.

amides is relatively more intense. At maximum oxidation there is also a loss of the aliphatic CH stretching band, suggesting a decreased C/N ratio in the most oxidized materials. The deconvolution of the nitrile band shows the progressive loss of isonitrile modes with increasing oxidation. Together with the loss of singly bonded nitriles, this suggests a gain in disorder in the structure of oxidized tholins. Mass loss through nitriles with respect to the amine and hetero-aromatic bands is a significant effect of oxidation.

Due to the large number of overlapping stretching and bending modes, deconvolution of the hetero-aromatic band is performed by simplifying the number of participant modes. The region between 1500 and 1300 cm^{-1} is attributed to a large peak, including the overlapping contribution of bending modes from CH, NH, and OH groups. As the CO_2/CH_4 ratio increases, the C=O stretching mode at $\sim 1700 \text{ cm}^{-1}$ increases, while the relative intensities of the C=N and aromatic C=C stretching modes decrease. In the plasma, oxidation is due to O atoms and OH radicals produced in the dissociation of CO_2 and CH_4 molecules. As the oxidation degree increases, the formation of NH and C–N bonds are favored relative to C=N bonds. We also note that, even without CO_2 in the gas mixture, the C=O mode is present in the reduced tholin. This could be due to surface oxidation by O_2 in air while transporting the sample for analysis or to trace oxygen species present in the plasma.

3.3. Absolute Imaginary Indices

Transmission spectra in the VUV–visible, near-infrared, and mid-infrared ranges for each sample and the thicknesses listed in Table 1 are used to estimate the wavelength-dependent imaginary index $k(\nu)$, over this spectral region, by means of the relation,

$$k(\nu) = \frac{\tau(\nu)}{4\pi\nu d}, \quad (2)$$

where $\tau(\nu)$ is the optical depth, ν is the frequency, and d is the sample thickness. The result is shown in Figure 5. The progressive increase of oxidation in these organic materials directly results in the increase of the absolute absorption over

Figure 6. Vis–VUV spectra of oxidized tholins irradiated with VUV photons. The $\tau_{\text{irr}} - \tau_o$ lines around 0 show the new and lost band intensities (scaled by a factor of 3).

Figure 7. Mid-infrared spectra (normalized) of VUV-irradiated tholins showing the evolution of the (a) amine (3600–2600 cm^{-1}) and (b) nitrile (2300–2050 cm^{-1}) bands.

the entire wavelength range (from 0.13 to 9 μm). For the most oxidized tholin, C/O = 0.68, the absolute absorption increases most prominently in the the VUV, 0.13–0.2 μm , and UV,

0.2–0.3 μm , in agreement with previous results reported in Gavilan et al. (2017).

In the mid-infrared the most absorptive region corresponds to the hetero-aromatic band, which increases linearly with the C/O ratio. We also note a significant drop of k over two orders of magnitude from the VUV to the visible region (0.13–1 μm) for the most reduced sample, the Titan type tholin with C/O = ∞ . McKay et al. (2001) showed that this drop is essential for reproducing the shape of Titan’s geometric albedo.

The uncertainty in near-infrared absorption is larger due to the lower absorption indices in this region, where $k \leq 0.2$. The absorption in the near-infrared increases with oxidation due to the presence of second and third overtones of NH, NH₂, and OH vibrations. This region is also characterized by absorption minima. An inversion of k is present at 1.8–2 μm for the C/O = ∞ tholin, and the position of this inversion feature blueshifts toward 0.9–1.1 μm for the tholin with C/O = 1 and to 0.5–0.6 μm for the tholin with C/O = 4. However, this k inversion is absent from the tholin corresponding to C/O = 0.83 (90:6:4). Although this sample is more oxidized, as seen by the decrease in nitrile intensity and the oxidation of the amine and hetero-aromatic bands, the lack of a minimum imaginary absorption index is surprising. This may be explained by the peculiar plasma conditions produced at a ratio of CO₂/CH₄ = 1.5. At this ratio the plasma attains a peculiar (dis-)equilibrium where tholin production is enhanced rather than quenched. Further experiments with this particular gas mixture should unveil the gas kinetics of this plasma and its effects on aerosol production.

4. VUV Irradiation of Tholins

Two tholins prepared with a ratio of CO₂/CH₄ = 1 were irradiated with a monochromatic beam of 160 nm photons. The VUV–visible spectra of the samples were measured before, τ_o , and after irradiation, τ_{irr} .

In all cases, a slight decrease in the spectral intensity in the 5.5–7 μm^{-1} (200–150 nm) region is attributed to the VUV photolysis of aromatic C = O structures. In contrast, we note the intensity growth of the 3.5–4.5 μm^{-1} (200–300 nm) region, attributed to π -bond formation of the organic matrix. The photon flux that the beam line produces was measured using an AXUV100G photodiode. Using the known responsivity of this diode, at $\lambda = 160$ nm there are 6.7×10^{11} photons/s/100 mA. During the irradiation, the beam current was recorded every second and the total fluence was calculated based on the average beam current (~ 180 mA). The 90:5:5 sample received a dose of $\sim 5 \times 10^{17}$ eV cm⁻², while the 90:6:4 sample received a dose of $\sim 8 \times 10^{17}$ eV cm⁻². The VUV spectra before and after irradiation are shown in Figure 6.

These results are consistent with previous VUV irradiation experiments performed by Gadallah et al. (2011) on hydrogenated amorphous carbon (HAC) materials. They used a deuterium lamp, whose emission peaks at 160 nm, and irradiated carbonaceous thin films with doses of at least 10^{22} eV cm⁻². UV spectra of the irradiated HAC samples showed an increase of the sp² carbon fraction, giving rise to π – π^* electronic transitions peaking at $\lambda^{-1} \sim 4.6 \mu\text{m}^{-1}$ (217 nm). Our irradiation experiments confirm that structural modifications can arise at lower UV fluences in organic materials. During irradiation, the transmission of the MgF₂ window could also evolve due to VUV photon absorption. However, as MgF₂ is a nonorganic crystal, the growth of new bands in the 200–300 nm range is attributed to the evolution of the tholin.

The mid-infrared spectra of the VUV irradiated samples were measured before and after irradiation as shown in Figure 7. Following irradiation, the amine band shows the greatest loss in intensity. This loss is most marked for amines in the 3400–3200 cm⁻¹ region, with a maximum loss at ~ 3300 cm⁻¹. The lower frequency modes in the amines region, including the CH and OH bands around 3000–2600 cm⁻¹, are little affected. Following VUV irradiation, the nitrile band shows a general depletion with a maximum loss of conjugated nitriles. Both mass loss (desorption) and photolysis (inducing fragmentation and/or bond creation) could occur during the VUV irradiation of tholins. Further experiments are required to unveil which mechanism is responsible for the photo-degradation effects.

To compare the laboratory UV doses to the radiation from an exoplanet host star, we use the UV spectra of GJ876, an M dwarf observed by the *International Ultraviolet Explorer* (Linsky 2014). In this study, the spectral flux is rescaled to the distance of its habitable zone (0.21 au). We consider the flux of GJ876 at $\lambda = 160$ nm, of ~ 8 erg cm⁻² s⁻¹, corresponding to the monochromatic photons used at the synchrotron. The laboratory doses are then equivalent to exposure times of ~ 28 hr (for the 90:5:5 tholin) and ~ 45 hr (for the 95:2.5:2.5 tholin) at 0.21 au of GJ876. Considering the low formation yields (and slow accumulation) of oxidized organic aerosols, it is likely that these will undergo photoprocessing during their lifetime in exoplanet atmospheres. The laboratory VUV-MIR spectra of photo-degraded tholins could be relevant to future observations of exoplanetary hazes.

5. Discussion

Future space telescopes like ARIEL⁹ and LUVOIR¹⁰ will enable the investigation of exoplanets ranging from Jupiter- and Neptune-size down to super-Earth and Earth-size. ARIEL instruments are expected to cover the visible and infrared 0.5–7.8 μm range (Tinetti et al. 2016), while LUVOIR will enable multiwavelength observations covering the ultraviolet to near-infrared from 0.2 to 2.5 μm .

Our spectroscopic VUV-MIR study covering the 0.13–10 μm range can thus provide relevant laboratory constraints for the future observations of exoplanet atmospheres. The spectra of oxidized aerosols signals a distinct large absorption in the 0.13–0.4 μm range and a flatter but strong absorption band in the 6–7.5 μm range relative to reduced aerosols. This high absorption coefficient may increase the detectability of exoplanet atmospheres where hazes form under oxidizing conditions. Together with a weaker nitrile band and a minimum absorption window in the visible range, these constitute potential observables of atmospheric oxidation. Organic aerosols constitute potential exoplanet atmospheric biosignatures, as organic hazes can be derived from biogenic methane (Catling et al. 2018), although abiotic sources are not discarded. While the simultaneous detection of CO₂ and CH₄ in a habitable exoplanet atmosphere has been recently proposed as a potential biosignature (Krissansen-Totton et al. 2018), ensuing oxidized organic hazes could also constitute observable biosignatures.

⁹ ARIEL: Atmospheric Remote-Sensing Infrared Exoplanet Large-survey.

¹⁰ LUVOIR: Large UV/Optical/IR Surveyor.

6. Conclusions

We have produced laboratory analogs to oxidized organic aerosols. These were prepared in increasingly oxidizing plasmas from mixtures of $\text{N}_2:\text{CO}_2:\text{CH}_4$, characterized by an increasing CO_2/CH_4 ratio from 0 (fully reducing) to 4. These correspond to atmospheres with C/O ratios from 0.68, 0.83, 1, and ∞ , i.e., representative of carbon-poor, solar-like to carbon-rich atmospheres. The samples were characterized via transmission spectroscopy from the vacuum-ultraviolet to the mid-infrared.

1. The molecular and electronic structure of these materials is dominated by the CO_2/CH_4 ratio in the $\text{N}_2:\text{CO}_2:\text{CH}_4$ gas mixture. Ultraviolet spectroscopy revealed the presence of $\pi-\pi^*$ and $n-\pi^*$ transitions in the $2 < \lambda^{-1} < 5 \mu\text{m}^{-1}$ (200–500 nm) range in all samples. These are attributed to electronic transitions in amine bonds for tholins prepared in fully reducing conditions. As the CO_2/CH_4 ratio increases new bands emerge in this range, attributed to electronic transitions in hydroxyl and carboxyl bonds. For all cases, a large absorption band in the far-ultraviolet is present and is attributed to $\sigma-\sigma^*$ and $n-\sigma^*$ electronic transitions.
2. Infrared spectroscopy revealed the formation of oxygenated bonds with an increasing CO_2/CH_4 ratio including stretching modes from NH, OH, C=O, and C=N and the formation of a new band in the $\sim 1400\text{--}1300 \text{ cm}^{-1}$ region, which could be attributed to NH, C–N, OH, and C–O bonds, as expected in CHON-rich organic matter. The decomposition of the nitriles band revealed the loss of aromatic nitriles and singly bonded R–N=C=N modes with oxidation.
3. The absolute absorption index increases linearly for all samples with increasing degrees of oxidation. The absorption is greatest in the VUV–UV range from 0.13 to $3 \mu\text{m}$ for the most oxidized samples, which are also the most opaque in the $6\text{--}10 \mu\text{m}$ hetero-aromatic band region. Oxidized aerosols show a weaker nitrile band and a blueshifted absorption minimum compared to reduced aerosols.
4. VUV irradiation of oxygenated aerosols revealed the formation of new $\pi-\pi^*$ bands in the $4 < \lambda^{-1} < 5 \mu\text{m}^{-1}$ (200–250 nm) region and the loss of electronic bands below 200 nm, corresponding to the formation of NH, C–N, and OH bonds and the loss of amines and isonitriles observed in their MIR spectra.
5. Our results suggest that oxidative plasma chemistry can result in solids with complex oxygenated bonds leading to solids that are more opaque in the VUV–MIR range. Because of their high absorption properties, oxidized organic aerosols could improve the detectability of exoplanet atmospheres. This spectral signature is indicative of the presence of oxic atmospheric conditions and could offer a potential biosignature for photosynthetic life.

We thank the staff at the ASTRID2 synchrotron for their help and access to beam time. We thank A. Giuliani for the preliminary VUV measurement of the reduced tholin sample at the DISCO beam line of the SOLEIL synchrotron (project No. 20160171) and E. Dartois for kindly supplying the DISCO-adapted sample holder. We thank A. Maturilli for his help with the near-infrared measurements at the Planetary Emissivity Laboratory (DLR). L.G. recognizes support for this visit from the Europlanet Research Infrastructure funded by the European

Union's Horizon 2020 research and innovation programme under grant agreement No. 654208. N.J.M. recognizes support from the COST Action TD1308 *Origins*. N.C. and L.G. thank the European Research Council for funding via the ERC *PrimChem* project (grant agreement No. 636829.).

Software: IDL MPFIT (Markwardt 2009).

ORCID iDs

Liseth Gavilan <https://orcid.org/0000-0001-8645-8415>
Nathalie Carrasco <https://orcid.org/0000-0002-0596-6336>

References

- Ari, T., & Güven, M. H. 2000, *JESRP*, 106, 29
- Arney, G. N., Meadows, V. S., Domagal-Goldman, S. D., et al. 2017, *ApJ*, 836, 49
- Barnes, E. E., & Simpson, W. T. 1963, *JChPh*, 39, 670
- Catling, D. C., Krissansen-Totton, J., Kiang, N. Y., et al. 2018, *AsBio*, 18, 709
- Duarte, A. A., Gomes, P. J., Ribeiro, J. H. F., et al. 2013, *EPJE*, 36, 98
- Feulner, G. 2012, *RvGeo*, 50, RG2006
- Fleury, B., Carrasco, N., Millan, M., Vettier, L., & Szopa, C. 2017, *E&PSL*, 479, 34
- Gadallah, K. A. K., Mutschke, H., & Jäger, C. 2011, *A&A*, 528, A56
- Gavilan, L., Alata, I., Le, K. C., et al. 2016, *A&A*, 586, A106
- Gavilan, L., Broch, L., Carrasco, N., Fleury, B., & Vettier, L. 2017, *ApJL*, 848, L5
- Haqq-Misra, J. D., Domagal-Goldman, S. D., Kasting, P. J., & Kasting, J. F. 2008, *AsBio*, 8, 1127
- Harshbarger, W. R. 1971, *JChPh*, 54, 2504
- Hornback, J. 2005, *Organic Chemistry* (2nd ed.; Boston, MA: Cengage Learning)
- Hubin-Franskin, M.-J., Delwiche, J., Giuliani, A., et al. 2002, *JChPh*, 116, 9261
- Imanaka, H., Khare, B. N., Elsila, J. E., et al. 2004, *Icar*, 168, 344
- Jacobson, M. C., Hansson, H. C., Noone, K. J., & Charlson, R. J. 2000, *RvGeo*, 38, 267
- Jones, A. P. 2012, *A&A*, 540, A2
- Kasting, J. F. 1993, *Sci*, 259, 920
- Krissansen-Totton, J., Olson, S., & Catling, D. C. 2018, *SciA*, 4, eaao5747
- Lavvas, P., & Koskinen, T. 2017, *ApJ*, in press (arXiv:1708.09257)
- Lavvas, P., Yelle, R. V., Koskinen, T., et al. 2013, *PNAS*, 110, 2729
- Linsky, J. 2014, *Chall*, 5, 351
- Madhusudhan, N., Agúndez, M., Moses, J. I., & Hu, Y. 2016, *SSRv*, 205, 285
- Mahjoub, A., Carrasco, N., Dahoo, P.-R., et al. 2012, *Icar*, 221, 670
- Markwardt, C. B. 2009, *adass XVIII*, 411, 251
- Marley, M. S., Ackerman, A. S., Cuzzi, J. N., & Kitzmann, D. 2013, in *Clouds and Hazes in Exoplanet Atmospheres*, ed. S. J. Mackwell et al. (Tucson, AZ: Univ. Arizona Press), 367
- McKay, C. P., Coustenis, A., Samuelson, R. E., et al. 2001, *P&SS*, 49, 79
- Mutsukura, N. 1999, *TSF*, 349, 115
- Palmer, M. H., Ridley, T., Hoffmann, S. V., et al. 2015, *JChPh*, 142, 134302
- Pavlov, A. A., Brown, L. L., & Kasting, J. F. 2001, *JGR*, 106, 23267
- Robertson, J., & O'Reilly, E. P. 1987, *PhRvB*, 35, 2946
- Sciamma-O'Brien, E., Dahoo, P.-R., Hadamcik, E., et al. 2012, *Icar*, 218, 356
- Swain, M. R., Tinetti, G., Vasisht, G., et al. 2009a, *ApJ*, 704, 1616
- Swain, M. R., Vasisht, G., & Tinetti, G. 2008, *Natur*, 452, 329
- Swain, M. R., Vasisht, G., Tinetti, G., et al. 2009b, *ApJL*, 690, L114
- Szopa, C., Cernogora, G., Boufendi, L., Correia, J. J., & Coll, P. 2006, *P&SS*, 54, 394
- Tannenbaum, E., Coffin, E. M., & Harrison, A. J. 1953, *JChPh*, 21, 311
- Tauc, J., Grigorovici, R., & Vancu, A. 1966, *PSSBR*, 15, 627
- Tinetti, G., Drossart, P., Eccleston, P., et al. 2016, *Proc. SPIE*, 9904, 99041X
- Trainer, M. G., Pavlov, A. A., Dewitt, H. L., et al. 2006, *PNAS*, 103, 18035
- Urbach, F. 1953, *PhRv*, 92, 1324
- Venot, O., Bénilan, Y., Fray, N., et al. 2018, *A&A*, 609, A34
- Venot, O., Hébrard, E., Agúndez, M., Decin, L., & Bounaceur, R. 2015, *A&A*, 577, A33
- von Paris, P., Rauer, H., Lee Grenfell, J., et al. 2008, *P&SS*, 56, 1244
- Wilson, E. H., & Atreya, S. K. 2004, *JGRE*, 109, E06002
- Wolf, E. T., & Toon, O. B. 2010, *Sci*, 328, 1266
- Workman, J. 2016, *Concise Handbook Of Analytical Spectroscopy, The: Theory, Applications, And Reference Materials* (Singapore: World Scientific)
- Xu, K., Amaral, G., & Zhang, J. 1999, *JChPh*, 111, 6271