AARHUS UNIVERSITY

Coversheet

This is the accepted manuscript (post-print version) of the article.

Contentwise, the accepted manuscript version is identical to the final published version, but there may be differences in typography and layout.

How to cite this publication

Please cite the final published version:

Philippsen, B. (2018). Reservoir Effects in a Stone Age Fjord on Lolland, Denmark. *Radiocarbon*, 60(2), 653-665. doi:10.1017/RDC.2018.6

Publication metadata

Title:	Reservoir Effects in a Stone Age Fjord on Lolland, Denmark
Author(s):	Philippsen, B.
Journal:	<i>Radiocarbon</i>
DOI/Link:	https://doi.org/10.1017/RDC.2018.6
Document version:	Accepted manuscript (post-print)

This article has been published in a revised form in Radiocarbon, <https://doi.org/10.1017/RDC.2018.6>. This version is free to view and download for private research and study only. Not for re-distribution or re-use. © 2018 by the Arizona Board of Regents on behalf of the University of Arizona

General Rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

If the document is published under a Creative Commons license, this applies instead of the general rights.

RADIOCARBON

CAMBRIDGE
UNIVERSITY PRESS**Reservoir effects in a Stone Age fjord on Lolland, Denmark**

Journal:	<i>Radiocarbon</i>
Manuscript ID	RDC-RES-2017-0027.R2
Manuscript Type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Philippsen, Bente; Museum Lolland-Falster, Femern project; Aarhus University, Aarhus AMS Centre, Department of Physics and Astronomy
Keywords:	Radiocarbon AMS dating, Coastal, Reservoir effects, Mesolithic, Neolithic

SCHOLARONE™
Manuscripts

Reservoir effects in a Stone Age fjord on Lolland, Denmark

*Bente Philippsen, Museum Lolland-Falster, Denmark (current address:
Aarhus AMS Centre, Department of Physics and Astronomy, Aarhus
University. Ny Munkegade 120, 8000 Aarhus C, Denmark.
bphilipp@phys.au.dk)*

Abstract

On the island of Lolland, southeast Denmark, an area of almost 300 ha is currently under archaeological investigation prior to the planned construction of a tunnel between Denmark and Germany under the Femern belt. The area investigated in the context of the "Femern project" includes a former fjord or lagoon, which was used both as an economic resource and as background for ritual activities in the Neolithic. The wetland conditions give excellent preservation conditions for organic material. A yet unsolved issue, however, is the question of reservoir effects. The local reservoir effect needs to be known for accurate radiocarbon dating of samples with possible aquatic carbon sources, such as human bones or food residues on pottery. Therefore, this paper attempts at calculating the local reservoir effect for the study area. I will discuss the possibilities and limitations when analysing radiocarbon dates from a rescue excavation. When applying the estimated reservoir corrections to a hoard of jaws and other bones, an interesting change in ritual activity at c. 4000 cal BC can be observed. Furthermore, I examined

radiocarbon dates on bulk organic sediment and will discuss their implications for building chronologies and for reconstructing the environment of the Stone Age fjord. Finally, I will discuss the pitfalls and uncertainties associated with radiocarbon dates for sea level reconstruction.

Introduction

The aim of this paper is to estimate the reservoir effect in a Stone Age fjord and how it influences the dating of materials such as bones and sediments, and samples used for sea level reconstruction. The paper demonstrates how the scarce data available from a rescue excavation can be used to obtain valuable information about the reservoir effect.

Each reservoir effect, e.g. the marine or the freshwater reservoir effect, can be a complex issue on its own (e.g. Olsson 1976; Heier-Nielsen et al. 1995; Ingram and Southon 1996; Lanting and van der Plicht 1998; Eiríksson et al. 2004; Ascough et al. 2005; Cage et al. 2006; Olsen et al. 2009; Keaveney and Reimer 2012; Loughheed et al. 2013; Philippsen et al. 2013; Heron and Craig 2015; Holmquist et al. 2015). In estuaries, such as the fjord of the study, these two combine. As the relative proportions of marine and fresh water, terrestrial runoff, and groundwater can vary on short and long timescales, reservoir effects are potentially highly variable in coastal waters (Philippsen et al. 2013). The mechanisms leading to freshwater reservoir effects have been explained elsewhere (Deevey et al. 1954; Broecker and Walton 1959; Olsson et al. 1983; Philippsen 2012; Philippsen 2013; Philippsen and Heinemeier 2013) and will not be described in detail in this paper. One important aspect is the fact that high freshwater reservoir ages can be caused by carbonate-rich water, the so-called

hardwater effect (e.g. Deevey et al. 1954; Philippsen et al. 2010). As the groundwater in the study area on the island of Lolland (see below) contains significant amounts of carbonate, a high freshwater reservoir effect can be expected. The total hardness of the groundwater on Lolland has been measured to 17.6°dH in Rødby (the town closest to the excavation area), and to 20.4, 19.8 and 21.5°dH on three other stations distributed across the island (<http://www.lollandforsyning.dk/vand/vandets-hardhedsgrad>, retrieved on October 5, 2015). In general, the average hardness of the water on Lolland lies between 18 and 24°dH (weighted average of all water works, weighted with their output; <http://data.geus.dk/geusmap/?mapname=drikkevand>, retrieved on October 5, 2015).

The reservoir age of the Baltic Sea is smaller or of the same order of magnitude as the oceans' reservoir age of about 400 years, but can be substantially higher, e.g. in fjords with freshwater influence (Heier-Nielsen et al. 1995; Hedenström and Possnert 2001; Lougheed et al. 2013).

It is vital to quantify reservoir effects in order to be able to correct radiocarbon dates of various materials. Most important in this context are human bones, food residues on pottery, and sediment samples. Therefore, the aim of this study is to quantify the relative contributions of the marine and freshwater reservoir effect to radiocarbon dates. It will be shown how the limited data retrieved during a rescue excavation can be used to estimate reservoir effects, and how radiocarbon ages of bulk sediment deviate from those of artefacts embedded in the same layers. Different fractions of terrestrial sediment were dated as well to get a better understanding of how the radiocarbon dates of different components

of the sediment. The estimated reservoir age will be applied to a small case study of a deposit of animal bones and to samples used for a sea level curve.

Study area

In the south of the island of Lolland, Denmark, extensive archaeological surveys precede the planned construction of a tunnel between Lolland and the German island of Fehmarn through the Femern Bælt/Fehmarnbelt (Figure 1, Sørensen (2017)).

The excavations (Figure 2) have so far yielded evidence from a large time-span, from around 10.000 calBC to the Iron Age. The period, which is represented best by our material, is the Early Neolithic, the time of the introduction of agriculture in Denmark. An entire economic and ritual landscape has been preserved by marine gyttja in the former Stone Age fjords (Sørensen 2017). However, even though there is evidence for animal husbandry, hunting, gathering and especially fishing continued to be of great importance. Large systems of stationary fishing devices have been found in the former Stone Age fjord. The rich estuarine environment attracted people throughout prehistory and caused large amounts of well preserved artefacts. At the same time, the heavy dependence on aquatic resources introduces a risk of reservoir effects in several different sample types.

Archaeological background

Reservoir corrections will enable radiocarbon dating of marine or partly marine samples, such as human bones or food residues on pottery. The direct dating of pottery will be important to understand the dynamics of the Mesolithic-Neolithic

transition, which is accompanied by a change in pottery style, from Ertebølle to Funnel Beaker pottery. The cultural transition accompanies the very gradual economic transition. Different aspects of the Danish Neolithic occur roughly contemporaneously, including domesticated animals, cereal agriculture, and Funnel Beakers. However, the exact relationship between those aspects has not been understood in detail. Did they arrive together as one “Neolithic package”, or were they adopted independently from each other? Precise and accurate radiocarbon dates of all Neolithic aspects are crucial to answer this question, and may lead to a better understanding for the reasons of the transition to the Neolithic.

Materials

At the time of writing, about 150 samples have been radiocarbon dated for the entire project, and more dates are continuously being produced. Most of them are wooden artefacts and constructions, and thus not applicable for reservoir effect studies.

No pairs of unequivocally contemporaneous terrestrial and marine samples have been found yet, neither in the excavations nor in special cores taken for scientific analyses. We can thus only compare terrestrial samples with the sediment from the same level, and marine samples with sediment from their level. Therefore, we can only indirectly estimate the reservoir age for the study region.

Radiocarbon dates of the following samples will be used to estimate the reservoir effect:

1. A wooden artefact on the sea floor and a sample of the sediment in which it was deposited

2. From a sediment core obtained in the former fjord:
 - Plant remains and sediment samples from the same level
 - Marine shells and sediment samples from the same level
3. Different chemical fractions of sediment samples from a terrestrial site in order to understand their origin – a discussion of sediment dates is necessary as we compare terrestrial and marine dates to sediment dates in 2.

The reservoir age estimates will be used in later phases of the project to correct e.g. radiocarbon dates of food residues on pottery. In this paper, the corrections will be applied to ritually deposited bones and to samples dated for a sea level curve of the study area.

Methods

Wood samples were pretreated with 1M HCl for one hour, 1M NaOH for three hours, both at 80°C, and lastly 1M HCl overnight at room temperature.

Sediment samples were pretreated using different methods. In the beginning of the project, sediment samples had been pretreated as “bulk” samples in the following way: 1M HCl for 1hr, 1M NaOH for 3 hr in total, changed up to two times, and lastly 1M HCl overnight, all at 20°C. The other pretreatment method resulted in base-soluble (“humic”) and base-insoluble (“residual”) fractions of the sediment: 1M HCl at 80°C for 1hr, repeated if necessary, 0.5M HCl NaOH at 80°C, repeated until no colour change was observed. Adding 6M HCl and slowly heating to 80°C precipitated the base-soluble fraction. Bulk sediment and the “residual” fraction thus represent similar extraction methods.

Shells were pretreated by mechanical and/or ultrasound cleaning, etching of the outer 10-20% with HCl, and removal of organic contamination by KMnO_4 at 80°C. They were converted to CO_2 using 80% H_3PO_4 in sealed evacuated containers.

Radiocarbon dates and stable isotope ratios were measured at the Aarhus AMS Centre, Department of Physics and Astronomy, Aarhus University.

Results

Table 1 displays the radiocarbon dates that can be used to discuss the order of magnitude of the reservoir effect, and the accuracy of sediment dates. The excavations from which these samples derive are mapped in Figure 2.

Radiocarbon dates for the sea level curve are presented in Table 2.

Reservoir effect calculation

A horizontal and a vertical stake from a woven mat of hazel branches, which was lying on the former sea floor, were dated (Table 1). It had most probably been part of a stationary fishing device, such as a fish weir or fence, before it was deposited on the sea floor. A sediment sample of the sea floor was dated as well. Not surprisingly, the radiocarbon dates of the two wood samples agree with each other. The sediment is more than 400 ^{14}C years older than the wood. As it has a marine $\delta^{13}\text{C}$ value, part of this age difference must be ascribed to the marine reservoir effect.

Four pairs of samples from a sediment core from the excavation MLF00902-I were dated (Table 1). Two shell samples were almost 200 ^{14}C years younger than

the surrounding sediment. This indicates that the sediment holds other age offset sources than just the reservoir effect.

Two pairs of unidentified plant and sediment were dated as well (Table 1). The plants are probably wetland plants, and are expected to have low reservoir effects due to heavy reliance on atmospheric CO₂. This is supported by the fact that the age difference between the plants and the sediments is greater than the age difference between the (marine) shells and the sediments.

If we assume that the ~200 year “sediment offset” also applies to the first sediment sample, AAR-19330, a reservoir effect can be calculated: $R = (\text{diff. sediment-wood}) - (\text{diff. sediment-shell}) = 434 \pm 40 - 187 \pm 31 = 247 \pm 51$ ¹⁴C years. However, this is only a rough estimate, and should in the future be replaced by a larger dataset of paired samples of terrestrial and marine origin, or by many terrestrial and marine samples from a long sediment sequence.

Radiocarbon dating of sediment samples

Sediment samples from four features (pits and postholes) from the excavation MLF00902-II have been dated using different fractions (AAR-24316 – 24319 in Table 1). Firstly, half of the sample was pretreated as “bulk sediment”, i.e. pretreated and dated in the same way as the many samples from the core drillings in the beginning of the project. Secondly, the base-soluble (“humic”) and base-insoluble (“residual”) fractions of the other half of the sample were extracted and dated. Therefore, we can compare three dates of the same sediment sample. In all four cases, the date on the humic fraction is the youngest. In three out of the four cases, the residual fraction is the oldest and the “bulk” sample represents an intermediate age. This is to be expected, as bulk sediment

and residual fraction are prepared with similar methods. The main difference is the fact that the complete removal of the base-soluble fraction was attempted for the residual fraction. Apart from these general tendencies, the age differences between the fractions are not consistent enough to allow the calculation of correction factors, which could be applied to compare “bulk” sediment dates from the early phases of the project with sediment dates from other studies. The $\delta^{13}\text{C}$ values of those samples indicate the terrestrial origin, which agrees with their origin from features such as pits and postholes. The temporal relation between the formation of the organic material in the sediment and the construction of the pits and postholes, however, is unknown.

Discussion

Reservoir effects

The reservoir effect in the Syltholm fjord is estimated to be about 250 ± 50 ^{14}C yr. This value is not unusual for the Western Baltic: a mollusc shell collected in 1888 at a water depth of 19m off Lolland had a reservoir age of 302 ± 50 ^{14}C years (Lougheed et al. 2013). Reservoir ages between 200 and 270 ^{14}C years were suggested for the Kattegat prior to 4500 BP (Olsen et al. 2009). Freshwater influence can be neglected, as the carbonate-rich freshwater in the streams on Lolland would increase the reservoir age (Lougheed et al. 2013). This is useful information when later dating other sample types, such as food residues on pottery or human bones. Already now, taking the marine reservoir effect into account helps to precisely date the bones of omnivores and carnivores, as the example in Figure 3 illustrates. The samples originate from an

area of about 25m², where about 50 mandibles, numerous artefacts and other bones had been deposited (Sørensen, 2017). Radiocarbon dates on mandibles from different animals have been corrected for the marine reservoir effect. The reservoir age of the marine food component consumed by omnivorous animals is expected to be greater than the 250 ¹⁴C years we found for the Syltholm fjord: the dogs would not have consumed molluscs from the fjord, but fish caught within the fjord and further out in the open Baltic.

It has therefore been assumed that the marine reservoir effect is about 400 years and that the endpoints for 100% terrestrial and 100% marine diet are -21 ‰ and -10 ‰, respectively. Later, when more data is available, those assumptions will of course be replaced by the actual reservoir effect and isotope baseline measured in the project. However, already now a general tendency can be observed: after correction, the omnivorous animals are shifted towards younger ages. There appears to be a shift in the deposition of mandibles from deer to omnivorous animals, before or at about 4000 calBC. Not regarding the young date of a wild boar, one could conclude that there is a shift from animals hunted for meat to those hunted (or kept, in the case of the dog) for fur.

Also for the reconstruction of a sea level curve (Table 2), reservoir effects are important. The following discussion will highlight some of the challenges with the use of radiocarbon dates for sea level reconstruction.

Fjord sediments and terrestrial sediments

All sediment samples from aquatic contexts of this project have marine $\delta^{13}\text{C}$ values. Therefore, the marine reservoir effect must be present in those cases, due to decayed marine organic matter from marine plants and animals. Influx of

terrestrial material into the fjord appears to be of lesser importance. However, the sediment samples had even larger radiocarbon ages than marine samples from the same layer. Therefore, sediment dates must contain another offset source than only the marine reservoir effect. For example, parts of the seafloor could have been eroded away before the shells were deposited, or organic material could have been redeposited centuries after its initial deposition (Winn et al. 1998; Rößler et al. 2011). In a sediment sequence from the coast of the Littorina Sea, for example, the age difference between sediment and terrestrial macrofossils was even up to 1100 and 700 ^{14}C years (Hedenström & Possnert, 2001).

When comparing radiocarbon dates of the sediment samples from terrestrial features with each other, one has to take the deposition environment into account – terrestrial sediments and marine sediments from our excavations cannot be compared directly. The former can have accumulated during soil formation processes, while the latter represent marine and terrestrial material in different proportions, which were deposited under water in the shallow fjords.

Irrespective of the depositional environment, however, this small case study highlights the uncertainties associated with sediment dating (see e.g. Olsson 1979). The different ages of the base-soluble and insoluble fractions indicate the different carbon sources for the sediment. The same effect was observed in sediments from the Baltic, which were used to date the Littorina transgression (Rößler et al. 2011). Sediment dating should therefore be avoided in those cases where human activities are to be dated. In the literature, for sediments containing too low amounts of terrestrial macrofossils, dating of organism-

specific compounds or biomarkers is suggested (Eglinton et al. 1996; Hatté and Jull 2013). In the context of the Femern project, radiocarbon dating paired samples of sediment and terrestrial remains is important in order to calculate correction factors. This would enable us to make use of the many dates on bulk sediment from the early phase of the project, where core drillings yielded no other datable material than bulk sediment.

Samples for a sea level curve

Tree stubs

Tree stubs are commonly used as sea-level indicators (Dörfler et al. 2009). Different tree species die at different water levels and can thus be used to reconstruct the sea level, provided that the cause of death for the tree was the rising water table. Some trees die already when the groundwater table rises, some others survive in wetlands and only die due to permanent inundation or flooding by the sea. The reliability of the radiocarbon date of a tree stub relies mainly on the sample selection process. Besides ensuring that the tree died due to the rising sea level, one should select wood from the outer, youngest, tree rings and avoid bark, which could be significantly older.

Wetland plants

Generally, aquatic plants should be avoided for radiocarbon dating. Even floating plants, which are expected to assimilate atmospheric CO₂, can have high reservoir effects (Philippsen and Heinemeier 2013). This can be due to internal air flow through the plant to the upper surface of the leaves (Dacey 1980), or due to DIC degassing (Hatté and Jull 2013). Also sphagnum macrofossils can be affected by reservoir effects due to the mineralization of old organic matter

(Hatté and Jull 2013). Reed is found in great numbers in the sediments of the study area. It is expected to rely largely on atmospheric photosynthesis and to have only a small risk of reservoir effects (Heikkinen and Äikää 1977). However, other carbon sources can be assimilated as well, including DIC, sedimentary CO₂ and nutrients from previous growing seasons that had been stored in the rhizomes. As reed thrives both in the (wet) terrestrial environment and at water depths of up to four metres (Haslam 2010), and as its roots can extend to up to four metres below ground level (Haslam 2010), this species adds an extra depths uncertainty to the sea level curve (Table 2).

Seeds and other terrestrial macrofossils

Generally, the most fragile terrestrial macrofossils should be chosen for dating, because they are most probable to show indications of bioturbation or redeposition, if present. Highly resistant seeds, such as those of *Carex*, can also be intact in reworked sediments and should therefore be avoided (Hatté and Jull 2013). Unidentified seeds could originate from aquatic or wetland species and should therefore be avoided as well.

Sediment compaction

Layers can be eroded or compacted, so level measurements can be problematic (Jakobsen et al. 2004; Dörfler et al. 2009; Baeteman et al. 2011). On the Femern excavations, this compaction can be observed with stakes and other wooden artefacts, which show a zigzag-profile today. The compaction is not uniform and varies considerably between sediment types. For example, on one site, a peat layer of about 20cm had compacted by 5cm, while the underlying moraine clay and the overlying marine gyttja showed almost no compaction.

Draft of a sea level curve

Taking into account all uncertainties, reservoir effects and level indicators, the shape of the sea level curve changes. Figure 4 exemplifies this effect with some of the dates (Table 2) that will be used to reconstruct the sea level curve for the study area. Radiocarbon ages and depth measurements are displayed as blue symbols. The blue curve is a sea level curve only based on these measurements. Additionally, shaded areas above, below, or around the dates display the range of possible sea levels and ages for each data point. For example, artefacts that have been deposited horizontally on the former sea floor indicate that the sea level must have been higher than the artefacts' level. A fireplace, in contrast, must have been above sea level, although it is uncertain how much. The black curve is drawn through those shaded areas and thus represents the most probable sea level at a given time. The resulting sea level curve is overall steeper than the blue curve and does not show the fluctuation that is present in the blue curve. However, at least at the present level of resolution, uncertainties in height are more important than uncertainties in radiocarbon age due to e.g. reservoir effects. Another challenge is the fact that the preliminary dataset lacks samples from the marine environment for the early phases of the project, and samples from the terrestrial environment for the later phases. However, as more radiocarbon dates are being measured and excavations are expected to continue in the future, improved sea level reconstructions will be possible.

Conclusion

Although systematic sampling for reservoir effect studies had not been an option in the study presented here, it was still possible to extract some research results from analyses made during the rescue excavation.

The marine reservoir effect in the study area appears to be approximately 250 ^{14}C years. Freshwater reservoir effects due to carbonate-rich freshwater influence have not been observed. Further research is necessary to follow the development of the marine reservoir effect through time. The reservoir effect in the shallow fjords of the study area might be lower than the reservoir effect of fish or marine mammals that lived in deeper parts of the Baltic. Therefore, not only shells, but also fish bones should be radiocarbon dated in comparison with contemporaneous terrestrial material. The average reservoir effect of fish and marine mammals caught by the people is expected to be larger than the 250 ^{14}C years of the fjord.

Radiocarbon ages of marine sediments are almost 200 ^{14}C years older than marine shells from the same layer. Therefore, erosion and redeposition of sediments have to be taken into account when dating marine sediments. From earlier phases of the project, large numbers of sediment dates are available. They had been used to roughly estimate the age of the organic sediments. The results of this study show that the ages of the sediment samples have to be corrected by 450-600 years to obtain the correct time of deposition. Further research will show whether this applies to the entire fjord system during all periods.

For the sea level curve, reservoir effects will be relevant when including shells or aquatic plants, although uncertainties in depth/height are an important issue as

well. Taking the different corrections into account, the sea level curve becomes significantly steeper than the one based on uncorrected depth/height relations.

Acknowledgments

I would like to thank all archaeologists and technical staff from Museum Lolland-Falster for their support in the field and magazines, for their constant helpfulness and interest in scientific archaeology. I had many good discussions about the archaeological part of this study with Søren A. Sørensen. He also made useful comments on an earlier version of this paper.

References

- Ascough P, Cook G, Dugmore A. 2005. Methodological approaches to determining the marine radiocarbon reservoir effect. *Progress in Physical Geography* 29(4):532-47.
- Baeteman C, Waller M, Kiden P. 2011. Reconstructing middle to late Holocene sea-level change: A methodological review with particular reference to 'A new Holocene sea-level curve for the southern North Sea' presented by K.-E. Behre. *Boreas* 40(4):557-72.
- Broecker WS, Walton A. 1959. The geochemistry of C^{14} in fresh-water systems. *Geochimica et Cosmochimica Acta* 16:15-38.
- Bronk Ramsey C. 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon* 51(1):337-60.
- Cage AG, Heinemeier J, Austin WEN. 2006. Marine radiocarbon reservoir ages in Scottish coastal and fjordic waters. *Radiocarbon* 48:31-43.

- Dacey JWH. 1980. Internal Winds in Water Lilies: An Adaptation for Life in Anaerobic Sediments. *Science* 210(4473):1017-9.
- Deevey ES, Gross MS, Hutchinson GE, Kraybill HL. 1954. The Natural C14 Contents of Materials from Hard-Water Lakes. *PNAS - Proceedings of the National Academy of Sciences of the United States of America* 40:285-8.
- Dörfler W, Jakobsen O, Kloof S. 2009. Indikatoren des nacheiszeitlichen Meeresspiegelanstiegs der Ostsee. Eine methodische Diskussion am Beispiel der Ostseeförde Schlei, Schleswig-Holstein. In: Müller J, al. e, editors. *Zwischen Nord- und Ostsee 1997-2007*. Bonn. p 177ff.
- Eglinton TI, Aluwihare LI, Bauer JE, Druffel ERM, McNichol AP. 1996. Gas chromatographic isolation of individual compounds from complex matrices for radiocarbon dating. *Analytical Chemistry* 68:904-12.
- Eiríksson J, Larsen G, Knudsen KL, Heinemeier J, Símonarson LA. 2004. Marine reservoir age variability and water mass distribution in the Iceland Sea. *Quaternary Science Reviews* 23:2247-68.
- Haslam SM. 2010. A Book of Reed (*Phragmites australis* (Cav.) Trin. ex Steudel, *Phragmites communis* Trin.): Forrest Text.
- Hatté C, Jull AJT. 2013. 14C of Plant Macrofossils. In: Elias SA, Mock CJ, editors. *Encyclopedia of Quaternary Science*. Amsterdam: Elsevier. p 361-7.
- Hedenström A, Possnert G. 2001. Reservoir ages in Baltic Sea sediment - a case study from the Litorina stage. *Quaternary Science Reviews* 20:1779-85.
- Heier-Nielsen S, Heinemeier J, Nielsen HL, Rud N. 1995. Recent Reservoir Ages for Danish Fjords and Marine Waters. *Radiocarbon* 37(3):875-82.
- Heikkinen A, Äikää O. 1977. Geological Survey of Finland radiocarbon measurements VII. *Radiocarbon* 19(2):263-79.

- Heron C, Craig OE. 2015. Aquatic Resources in Foodcrusts: Identification and Implication.
- Holmquist JR, Reynolds L, Brown LN, Southon JR, Simms AR, MacDonald GM. 2015. Marine radiocarbon reservoir values in Southern California estuaries: interspecies, latitudinal, and interannual variability. *Radiocarbon* 57(3):449-58.
- Ingram BL, Southon JR. 1996. Reservoir Ages in Eastern Pacific Coastal and Estuarine Waters. *Radiocarbon* 38(3):573-82.
- Jakobsen O, Meurers-Balke J, Hoffmann-Wieck G, Thiede J. 2004. Postglazialer Meeresspiegelanstieg in der südwestlichen Ostsee. In: Schernewski G, Dolch T, editors. Ergebnisse der 22. Jahrestagung des Arbeitskreises "Geographie der Meere und Küsten" Coastline Report 1. Warnemünde. p 9-21.
- Keaveney EM, Reimer PJ. 2012. Understanding the variability in freshwater radiocarbon reservoir offsets: a cautionary tale. *Journal of Archaeological Science* 39(5):1306-16.
- Lanting JN, van der Plicht J. 1998. Reservoir effects and apparent ^{14}C ages. *The Journal of Irish Archaeology* 9:151-65.
- Lougheed BC, Filipsson HL, Snowball I. 2013. Large spatial variations in coastal ^{14}C reservoir age - a case study from the Baltic Sea. *Clim. Past* 9(3):1015-28.
- Nehring S, Leuchs H. 2000. Wiederfund der 'verschollenen' Netzreusenschnecke *Nassarius reticulatus* (Linnaeus, 1758) in der Mecklenburger Bucht. *Archiv der Freunde der Naturgeschichte in Mecklenburg* 39:105-14.

- Olsen J, Rasmussen P, Heinemeier J. 2009. Holocene temporal and spatial variation in the radiocarbon reservoir age of three Danish fjords. *Boreas* 38(3):458-70.
- Olsson IU. The radiocarbon content of various reservoirs. In: Berger R, Suess HE, editors; 1976; Los Angeles and La Jolla. University of California Press. p 613-8.
- Olsson IU. 1979. A warning against radiocarbon dating of samples containing little carbon. *Boreas* 8(2):203-7.
- Olsson IU, El-Daoushy F, Vasari Y. 1983. Säynäjälampi and the difficulties inherent in the dating of sediments in a hard-water lake. *Hydrobiologia* 103(1):5-14.
- Philippsen B, Kjeldsen H, Hartz S, Paulsen H, Clausen I, Heinemeier J. 2010. The hardwater effect in AMS ^{14}C dating of food crusts on pottery. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* 268(7-8):995-8.
- Philippsen B. 2012. Variability of freshwater reservoir effects: Implications for radiocarbon dating of prehistoric pottery and organisms from estuarine environments: Aarhus University.
- Philippsen B. 2013. The freshwater reservoir effect in radiocarbon dating. *Heritage Science* 1:24.
- Philippsen B, Heinemeier J. 2013. Freshwater reservoir effect variability in Northern Germany. *Radiocarbon* 55(2-3):1085-101.
- Philippsen B, Olsen J, Lewis JP, Rasmussen P, Ryves DB, Knudsen KL. 2013. Mid- to late-Holocene reservoir age variability and isotope-based

palaeoenvironmental reconstruction in the Limfjord, Denmark. *The Holocene* 23(7):1015 - 25.

Reimer PJ, Bard E, Bayliss A, Beck JW, Blackwell PG, Bronk Ramsey C, Buck CE, Cheng H, Edwards RL, Friedrich M, Grootes PM, Guilderson TP, Hafflidason H, Hajdas I, Hatté C, Heaton TJ, Hoffmann DL, Hogg AG, Hughen KA, Kaiser KF, Kromer B, Manning SW, Niu M, Reimer RW, Richards DA, Scott EM, Southon JR, Staff RA, Turney CSM, van der Plicht J. 2013. IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP. *Radiocarbon* 55(4):1869-87.

Rößler D, Moros M, Lemke W. 2011. The Littorina transgression in the southwestern Baltic Sea: new insights based on proxy methods and radiocarbon dating of sediment cores. *Boreas* 40(2):231-41.

Sørensen SA. 2017. Denmark's largest Stone Age excavation. Mesolithic Miscellany submitted.

Winn K, Erlenkeuser H, Nordberg K, Gustafsson M. 1998. Paleohydrography of the Great Belt, Denmark, during the Littorina transgression: the isotope signal. *Meyniana* 50:237-51.

Statement

This manuscript has not been published elsewhere and has not been submitted simultaneously for publication elsewhere.

Figures and tables

Figure 1: Location map of the excavation area of the Femern project. Today a straight coastline protected by a dike, the area comprised an ever-changing fjord landscape during the Stone Age. Background map in the public domain (via commons.wikimedia.org), map of Denmark by krak.dk, aerial photograph by google.com.

Figure 2: Location of the different excavations of the Femern project. The excavations have numbers such as MLF01078 or MLF00906-I, on this map abbreviated as e.g. 1078 or 906-I.

Figure 3: Radiocarbon dates with and without reservoir correction from an area where artefacts and animal bones, especially mandibles, had been deposited in the shallow water of the fjord. Calibrated with OxCal 4.2, IntCal 13 (Bronk Ramsey 2009; Reimer et al. 2013).

Figure 4: Examples for the effects of uncertainties in depth and radiocarbon ages on sea level reconstruction. Radiocarbon dates and levels (below present sea level) of samples that can be used for sea level reconstruction are marked with blue symbols. Shaded areas represent the possible ranges of the past sea level.

Table 1: Paired radiocarbon dates on wood, sediment and shells, which can be used to estimate reservoir effects.

AAR	Name	^{14}C age	$\delta^{13}\text{C}$ (‰ VPDB)		
19328	MLF00906-I P29, horizontal branch	3932±28	-31.95±0.05	Weighted mean, wood: 3951±26 ^{14}C yr	Difference sediment – wood: 434±40 ^{14}C yr
19329	MLF00906-I P30, vertical stake	3969±27	-30.84±0.05		
19330	MLF00906-I P31, bulk sediment	4385±30	-16.88±0.05		

20429	MLF00902-I P1- Lag 5 - A, plant, maybe rush/reed	3307±32	-25.04±0.05	Difference plant – sediment: 351±44 ¹⁴ C yr	Average: 360±33 ¹⁴ C yr
20433	MLF00902-I P1- Lag 5 – B, bulk sediment	3658±30	-17.41±0.05		
20430	MLF00902-I P1- Lag 7 – A, unident. plant	3679±39	-20.20±0.05	Difference plant – sediment:	
20434	MLF00902-I P1- Lag 7 – B, bulk sediment	4050±30	-18.11±0.05	371±49 ¹⁴ C yr	
20431	MLF00902-I P1- Lag 10 – A, cockle shell	4250±32	-2.08±0.05	Difference shell – sediment:	Average: 187±31 ¹⁴ C yr
20435	MLF00902-I P1- Lag 10 – B, bulk sediment	4434±31	-18.97±0.05	184±45 ¹⁴ C yr	
20432	MLF00902-I P1- Lag 13 – A, unident. shells, maybe Acrididae	4381±30	0.65±0.05	Difference shell – sediment:	
20436	MLF00902-I P1- Lag 13 – B, bulk sediment	4571±29	-20.07±0.05	190±42 ¹⁴ C yr	

24312	MLF00902-II A9 (pit) P11 Bulk sediment	8080±34	-29.68 ± 0.31	Difference bulk- residual: -208±48	
24316.1	MLF00902-II A9 (pit) P11 Sediment, residual	8288±34	-37 ± 0.31	Difference bulk-humic: 425±53	
24316.2	MLF00902-II A9 (pit) P11 Sediment, humic	7655±41	-26.47 ± 0.31	Difference residual-humic: 633±53 ¹⁴ C yr	
24313	MLF00902-II A17 (posthole) P12 Bulk sediment	6457±32	-29.03 ± 0.31	Difference bulk-residual: -84±45	
24317.1	MLF00902-II A17 (posthole) P12 Sediment, residual	6541±32	-29.91 ± 0.31	Difference bulk-humic: 441±45	
24317.2	MLF00902-II A17 (posthole) P12 Sediment, humic	6016±31	-28.16 ± 0.31	Difference residual-humic: 525±45 ¹⁴ C yr	
24314	MLF00902-II A15 (pit) P17 Bulk sediment	6228±32	-31.67 ± 0.31	Difference bulk-residual: 70±45	
24318.1	MLF00902-II A15	6158±31	-27.25 ± 0.31	Difference bulk-humic: 610±54	

	(pit) P17 Sediment, residual			
24318.2	MLF00902-II A15 (pit) P17 Sediment, humic	5618±43	-28.02 ± 0.31	Difference residual-humic: 540±53 ¹⁴ C yr
24315	MLF00902-II A21 (posthole) P22 Bulk sediment	6770±32	-32.41 ± 0.31	Difference bulk- residual: -260±47
24319.1	MLF00902-II A21 (posthole) P22 Sediment, residual	7030±34	-32.44 ± 0.31	Difference bulk-humic: 834±45
24319.1	MLF00902-II A21 (posthole) P22 Sediment, humic	5936±31	-27.3 ± 0.31	Difference residual-humic: 1094±46 ¹⁴ C yr

Appendix

Table 2: Samples used to exemplify the effect of age and level uncertainties on a sea level curve. Age uncertainties comprise for example the old wood effect and especially the marine reservoir effect. Furthermore, there are uncertainties in the relation between a sample's level and the inferred sea level.

Lab code	Material and species	^{14}C age	Age span/ uncertainty	Level bpsl	Level span/ uncertainty
AAR-21521	<i>Phragmites</i>	4280±32	Reservoir	-1.44	From sea level or several meters below: can grow in water depths of up to 4 meters, and the roots can grow up to 5 m deep.
AAR-22307	<i>australis</i> (common	5983±28	effect between	-3.25	
AAR-22308	reed), stalk,	5921±29	0 and 400	-2.33	
AAR-21519	leaves, roots or	5998±28	years –	-2.59	
	rhizomes (not specified)		probably little		
AAR-21531	<i>Phragmites australis</i> roots	5053±33	reservoir effect (Heikkinen & Äikää, 1977)	-1.15	
AAR 21522	<i>Suaeda</i>	3864±29	Unknown	-1.40	Probably none – the plants grow at sea level, and the seeds/plant remains were probably deposited at that level
AAR-21527	<i>maritima</i> (sea	4172±36	reservoir	-1.51	
AAR-21528	blite)	4522±32	effect, but	-1.57	
AAR-21515	Seeds of <i>Atriplex</i> and <i>Suaeda</i>	1163±27	probably mainly atmospheric	-0.81	
AAR-21518	Seeds of <i>Atriplex</i> from an organic-rich layer in a	1903±25	photosynthesis	-1.32	

	former channel				
AAR-21523	<i>Cladium mariscus</i> (saw grass)	7396±35		-13.47	Probably none – grows on wet soil behind the reed belt, probably at or around sea level
AAR-21516	Piece of wood from an organic layer	1097±25	Probably redeposited	-0.67	Probably redeposited
AAR-22067	Root of an oak trunk	5984±33	No old wood effect (ring 7+8 of 10-year-old root dated)	-2.33	Depth of root in relation to former soil surface or sea level unknown. Oaks die, before the water table reaches the roots.
AAR-22087		5440±26	Old wood effect small (root 12 years old)	-1.27	
AAR-21525	<i>Quercus</i> (oak)	4954±33	No old wood effect – outer 3 rings dated	-1.17	Oaks die, before the water table reaches the roots.
AAR-21526	trunk	6019±31		-2.43	
AAR-21532		6049±26		-2.43	
AAR-21514		6115±28		-2.59	
AAR-22066	<i>Quercus</i> (oak) trunk	6036±29	Old wood effect ≥ 25 years	-2.22	
AAR-21533	<i>Alnus</i> (alder) trunk	6321±31	No old wood effect – outer 3 rings dated	-2.86	Alder can tolerate a very high water table

AAR-21044, 21045, 21046	Charcoal from fireplace (weighted mean of three dates)	6321±17	Probably no old wood effect – outer rings of twigs dated	-2.31	The fireplace must have been on dry land – but how high, is unknown.
MLF00902-II P3a	Charcoal from cooking pit	5863±31	Possible old wood effect (although the risk and order of magnitude are expected to be low)	-1.66	Must have been above sea level, but unknown how much
AAR-22281, 22282	Burnt bones from an undisturbed activity layer (weighted mean of two dates)	10008±32	The carbon originates from the fuel and the bones/animal – small possibility of old wood effect	-2.55	Deposited on dry land; the bone fragments could have moved down through the soil
AAR-21529	<i>Ostrea edulis</i> (oyster) shell	4584±28	Reservoir age around 250-400 years	-1.75	Assuming the shells have not been redeposited: they typically occur in water depths of 1-10m
AAR-21530		4616±29		-1.63	
AAR-21520		5675±28		-2.24	

AAR-21517	<i>Nassarius reticulatus</i> (netted dog-whelk) shell	4388±29	Reservoir age around 250-400 years	-1.80	Occurs at water depths up to 50 m; buries a few cm into the sediment (Nehring and Leuchs 2000)
AAR-21390 A7	Mats (A7 and A9) of woven	4188±27	None – young branches	-1.39	The mats were found lying on the former sea floor. The sea level is expected to have been up to one or two meters above the mats.
AAR-21394 A7	hazel twigs, probably from	4233±25			
AAR-21395 A9	fish fence	4202±25		-1.62

Figure 1: Location map of the excavation area of the Femern project. Today a straight coastline protected by a dike, the area comprised an ever-changing fjord landscape during the Stone Age. Background map in the public domain (via commons.wikimedia.org), map of Denmark by krak.dk, aerial photograph by google.com.

Figure 2: Location of the different excavations of the Femern project. The excavations have numbers such as MLF01078 or MLF00906-I, on this map abbreviated as e.g. 1078 or 906-I.