AARHUS UNIVERSITY

Coversheet

This is the accepted manuscript (post-print version) of the article.

Contentwise, the accepted manuscript version is identical to the final published version, but there may be differences in typography and layout.

How to cite this publication

Please cite the final published version:

He, F., Bhoobalan-Chitty, Y., Van, L.B. *et al.* Anti-CRISPR proteins encoded by archaeal lytic viruses inhibit subtype I-D immunity. *Nat Microbiol* **3**, 461–469 (2018). <https://doi.org/10.1038/s41564-018-0120-z>

Publication metadata

Title:	Anti-CRISPR proteins encoded by archaeal lytic viruses inhibit subtype I-D immunity
Author(s):	F. He, Y. Bhoobalan-Chitty, L.B. Van <i>et al</i>
Journal:	Nature Microbiology
DOI/Link:	https://doi.org/10.1038/s41564-018-0120-z
Document version:	Accepted manuscript (post-print)

This is the peer reviewed version of the following article: He, F., Bhoobalan-Chitty, Y., Van, L.B. et al. Anti-CRISPR proteins encoded by archaeal lytic viruses inhibit subtype I-D immunity. Nat Microbiol 3, 461–469 (2018), which has been published in final form at <https://doi.org/10.1038/s41564-018-0120-z>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

General Rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

If the document is published under a Creative Commons license, this applies instead of the general rights.

1 **Anti-CRISPR proteins encoded by archaeal lytic viruses inhibit**
2 **subtype I-D immunity**

3 Fei He^{1#}, Yuvaraj Bhoobalan-Chitty^{1#}, Lan B Van², Anders L Kjeldsen¹, Matteo Dedola¹, Kira S
4 Makarova³, Eugene V Koonin³, Ditlev E Brodersen², Xu Peng^{1*}

¹Danish Archaea Centre, Department of Biology, University of Copenhagen, Ole Maaløes Vej 5,
DK-2200 Copenhagen N, Denmark.

²Centre for Bacterial Stress Response and Persistence, Department of Molecular Biology and
Genetics, Aarhus University, Gustav Wieds Vej 10c, DK-8000 Aarhus C, Denmark.

5 ³National Center for Biotechnology Information, National Library of Medicine, NIH, Bethesda,
6 USA.

7 # These authors contributed equally to the work

8 *Correspondence to: peng@bio.ku.dk

9

10 **Abstract**

11 Viruses employ a range of strategies to counteract the prokaryotic adaptive immune system,
12 CRISPR-Cas, including mutational escape and physical blocking of enzymatic function using anti-
13 CRISPR proteins (Acrs). Acrs have been found in many bacteriophages but so far not in archaeal
14 viruses despite the near ubiquity of CRISPR-Cas systems in archaea. Here, we report the functional
15 and structural characterization of two archaeal Acrs from the lytic rudiviruses, SIRV2 and SIRV3.
16 We show that a 4-kb deletion in the SIRV2 genome dramatically reduces infectivity in *Sulfolobus*
17 *islandicus* LAL14/1 that carries functional CRISPR-Cas subtypes I-A, I-D, and III-B. Subsequent
18 insertion of a single gene from SIRV3, gp02 (acrID1), which is conserved in the deleted fragment,
19 successfully restored infectivity. We demonstrate that AcrID1 protein inhibits the CRISPR-Cas
20 subtype I-D system by interacting directly with Cas10d protein, which is required for the
21 interference stage. Sequence and structural analysis of AcrID1 show that it belongs to a conserved
22 family of compact, dimeric $\alpha\beta$ -sandwich proteins characterized by extreme pH and temperature
23 stability and a tendency to form protein fibers. We identify about 50 homologs of AcrID1 in four
24 archaeal viral families demonstrating the broad distribution of this group of anti-CRISPR proteins.

25

26

27 **Introduction**

28 Prokaryotes have evolved multiple strategies to combat mobile genetic elements (MGEs) such as
29 viruses and plasmids¹. Of particular interest are the CRISPR-Cas (Clustered Regularly Interspaced
30 Short Palindromic Repeats and CRISPR-associated proteins) adaptive immune systems, which are
31 present in many bacteria and almost all archaea²⁻⁴. Short fragments of DNA originating from MGEs,
32 known as spacers, are incorporated into CRISPR arrays between repeat sequences (a process
33 dubbed "adaptation") and subsequently transcribed and processed into mature CRISPR RNAs
34 (crRNAs)⁵⁻⁸. During the "interference" stage, the mature crRNA and a set of Cas proteins form a
35 complex that recognizes and inactivates MGE-containing sequences complementary to the spacer
36 portion of the crRNA by cleaving the MGE genome^{2,9}. On the basis of distinct architectures of the
37 effector complexes that are involved in interference, the CRISPR-Cas systems have been divided
38 into Class 1 and Class 2, each encompassing three types (Class 1: types I, III and IV; Class 2: types
39 II, V and VI), with several subtypes for each type^{2,4}.

40 In the arms race between viruses and their hosts, viruses evade CRISPR-Cas immunity via
41 mutational escape. In addition, some bacteriophages have been shown to encode anti-CRISPR (Acr)
42 proteins that inhibit CRISPR-Cas by binding directly to and inhibiting specific subunits of the
43 effector complexes¹⁰⁻¹². The Acrs identified to date originate primarily from temperate
44 bacteriophages, prophages, and genomic islands but also from virulent phages associated with some
45 bacterial phyla. The Acrs form 21 families of small proteins that appear to evolve fast, presumably
46 as a result of the arms race with CRISPR-Cas systems, and have no detectable homologs in the
47 current protein databases¹³. Such proteins have so far been shown to inhibit bacterial CRISPR-Cas
48 systems of subtypes I-E, I-F, II-A, and II-C^{11,13-18}. Although CRISPR-Cas systems are encoded in
49 almost all archaeal genomes, no Acrs have so far been identified in the genomes of the numerous
50 known archaeal viruses or other MGEs¹⁹.

51 The crenarchaeon *Sulfolobus islandicus* LAL14/1 is a laboratory host for the lytic rod-shaped virus
52 SIRV2₂₀. The LAL14/1 genome encodes five CRISPR arrays, four interference *cas* gene cassettes
53 (one subtype I-A, one subtype I-D, and two subtype III-B loci) and two adaptation *cas* gene
54 cassettes (one subtype I-A and one subtype I-D). The five repeat arrays carry 13 spacers matching
55 the SIRV2 genome, each with a cognate Protospacer Adjacent Motif (PAM) and 2 to 6 bp
56 mismatches. Despite imperfect complementarity to the viral target, at least some of these spacers
57 are predicted to be functional because the mismatches are located outside the seed sequence that is
58 essential for interference²¹⁻²⁴. Nevertheless, SIRV2 is able to propagate in LAL14/1 with an average
59 viral yield of 30 virions/cell and causes host chromosome degradation, eventually leading to cell
60 lysis²⁵. The ability of SIRV2 to propagate in the presence of active CRISPR-Cas systems carrying
61 matching spacers suggests that the virus could encode anti-CRISPR proteins.

62 Here, we report the functional and structural characterization of two Acrs encoded by SIRV2 and
63 SIRV3. We initially identify a mutant of SIRV2 (SIRV2M) that arises after successive transfers of
64 the virus in a CRISPR-deficient host. SIRV2M is characterized by a 4-kb deletion near one of the
65 termini of the linear viral genome and has lost the capacity to infect the original host LAL14/1. By
66 inserting SIRV3 genes that are conserved in the deleted 4 kb fragment into the SIRV2M genome,
67 we further identify a single gene capable of restoring the SIRV2M infectivity in LAL14/1. Finally,
68 we show that the product of this gene, AcrID1, is a conserved and compact dimeric $\alpha\beta$ -sandwich
69 protein conferring anti-CRISPR activity by interacting directly with Cas10d, the large subunit of the
70 subtype I-D CRISPR-Cas effector complex in *Sulfolobus islandicus*.

71

72 **Results**

73 **Isolation of a SIRV2 mutant that is non-infectious in *S. islandicus* LAL14/1**

74 *Sulfolobus solfataricus* strain 5E6 lacks functional CRISPR-Cas systems and is highly susceptible
75 to SIRV2 infection²⁶. During our earlier transcriptomic²⁶, genetic and biochemical studies²⁷⁻³¹ of
76 SIRV2 in this host, we occasionally observed deletions at the left genomic terminus. These
77 observations prompted us to monitor the genomic stability of this site during successive transfers of
78 SIRV2 infected culture supernatant to fresh cells of this CRISPR-Cas deficient host. After several
79 transfers, SIRV2 was observed to gradually lose a near-terminal 4-kb fragment spanning the *gp02*
80 to *gp09* genes, as shown by PCR amplification of this genomic region (Fig. 1a and 1b). However,
81 the presence of longer PCR products suggests that smaller deletions also occurred and a low amount
82 of wild-type SIRV2 was still present in the viral mixture. A mutant isolate that we denote SIRV2M
83 was purified from the mixture by a plaque assay, and subsequent PCR amplification and sequencing
84 demonstrated that the deletion corresponds to positions 1491 to 5474 bp (ORFs *gp02* through *gp09*)
85 of the SIRV2 genome, and likely result from recombination between two adjacent 13 bp repeats
86 (Fig. 1a and 1c). The importance of this 3984-bp fragment for viral reproduction in CRISPR-Cas
87 carrying hosts was further reinforced by the inability of SIRV2M to inhibit growth of *S. islandicus*
88 LAL14/1 (Fig. 1d). Subsequent infection of *S. islandicus* LAL14/1 with the SIRV2 deletion mixture
89 favored propagation of wild-type SIRV2 as suggested by the increased intensity of the
90 corresponding PCR band (Fig. 1b, wt) and concomitant decrease of the lower band (Fig. 1b, m).
91 Together, these results suggest that the 4-kb region lost in SIRV2M contains one or more genes that
92 are important for SIRV2 propagation in a host with functional CRISPR-Cas immunity.

93 **SIRV3 *gp02* is necessary and sufficient to restore SIRV2M infectivity in LAL14/1**

94 Next, we tested if reintroduction of the deleted fragment into the SIRV2M genome through
95 homologous recombination would restore viral infectivity. To this end, a PCR-amplified DNA

96 fragment containing SIRV2 *gp02-gp09* flanked by the repeat sequence was electroporated into *S.*
97 *islandicus* LAL14/1 cells, and SIRV2M was immediately added. The PCR analysis showed that this
98 fragment successfully recombined with the SIRV2M genome and enabled resultant recombinant
99 viral genome to propagate in the culture (Supplementary Fig. 1). To facilitate the search for genes
100 required for SIRV2M propagation in LAL14/1, we employed another *Sulfolobus* rudivirus, SIRV3,
101 which, like SIRV2, is infectious in LAL14/1 but carries no readily detectable homologs of SIRV2
102 *gp02*, *gp04*, *gp05*, *gp08*, or *gp09*³². Accordingly, we inferred that the gene(s) required for SIRV2M
103 propagation in LAL14/1 were among the remaining three genes, SIRV3 *gp02*, *gp04* and *gp05* which
104 share between 70% and 80% nucleotide sequence identity with SIRV2 *gp03*, *gp06* and *gp07*,
105 respectively (Fig. 2a).

106

107 Fragments of various lengths, containing up to four of the SIRV3 genes *gp02* to *gp05*, were
108 amplified by PCR and inserted into the bacterial plasmid pJET1.2, which does not replicate in *S.*
109 *islandicus*. The cloned region was flanked by two fragments, each of about 500 bp, derived from
110 the region covering SIRV2 *gp22* and *gp23* to facilitate recombination between the clones and
111 SIRV2M resulting in insertion of the SIRV3 genes between SIRV2 *gp22* and *gp23* (Fig. 2b). The
112 plasmids were electroporated into LAL14/1 and SIRV2M was added immediately, followed by
113 incubation at 78°C. Potential recombination events and propagation of the recombinant viral
114 genome were monitored by subsequent PCR amplification of the SIRV2 *gp22-gp23* region (Fig. 2b,
115 2c). PCR amplification using culture supernatants, before dilution and further viral propagation in a
116 second culture, revealed a band derived from SIRV2M in all cultures, suggesting that DNA of the
117 initially added SIRV2M was still present after 2 days of incubation following electroporation.
118 Importantly, some samples showed extra bands that were larger than the SIRV2M band, suggesting
119 that recombination between the clones and SIRV2M genome indeed occurred in these samples (Fig.

120 2c). The culture supernatants were then diluted 1000 times into fresh LAL14/1 cells (50 ml) and
121 PCR performed again after two days of incubation (Fig. 2d). The DNA band derived from SIRV2M
122 disappeared in most samples, possibly due to dilution and/or DNA degradation of the initially added
123 SIRV2M. In contrast, larger PCR products were obtained from samples that were initially
124 electroporated with clones containing SIRV3 *gp02* alone or together with other genes. Thus, SIRV3
125 *gp02* integration into the SIRV2M genome apparently restored viral infectivity and allowed the
126 recombinant viruses to spread in the culture. Consistently, all clones that lacked an intact copy of
127 SIRV3 *gp02* failed to restore SIRV2M infectivity (Fig. 2d).

128 The recombinant virus derived from SIRV2M, with only SIRV3 *gp02* inserted (denoted
129 SIRV2Mgp02), was purified and shown to substantially retard LAL14/1 growth (Fig. 3A). Spot
130 assays demonstrated similar infection efficiencies of SIRV2Mgp02 and wt SIRV2, whereas
131 SIRV2M exhibited a 100 to 1000-fold lower infectivity (Fig. 3b). Furthermore, SIRV2Mgp02
132 propagated efficiently in LAL14/1 cells, whereas SIRV2M was completely eliminated from this
133 culture (Fig. 3c). Finally, overexpression of the plasmid-borne SIRV3 *gp02* in LAL14/1 led to host
134 growth retardation after SIRV2M infection (Fig. 3d). Together, these results indicate that *gp02*
135 alone is necessary and sufficient to restore SIRV2M infectivity in *S. islandicus* LAL14/1 possibly
136 through inhibiting the CRISPR-Cas immunity.

137 **SIRV3 *gp02* (AcrID1) inhibits CRISPR-Cas subtype I-D immunity**

138 The *S. islandicus* LAL14/1 genome carries two *cas6* genes which are responsible for crRNA
139 processing and thus essential for immunity, one in the subtype I-A and another in the subtype I-D
140 system²⁰. To determine whether SIRV3 *gp02* inhibits any one or both of these CRISPR-Cas systems,
141 we deleted the two *cas6* genes individually in *S. islandicus* LAL14/1 (Supplementary Fig. 2) and
142 subjected the deletion mutants to viral infection. Notably, SIRV2M gained infectivity in the I-D

143 *cas6* knockout mutant $\Delta cas6(I-D)$, but not in the I-A *cas6* knockout mutant $\Delta cas6(I-A)$ (Fig. 4a, left
144 and middle). In contrast, SIRV2Mgp02 was infectious in both mutants (Fig. 4a, left and middle).
145 Complementation of $\Delta cas6(I-D)$ with a plasmid-borne I-D *cas6* restored the resistance to SIRV2M
146 but not to SIRV2Mgp02 (Fig. 4a, right). Together, these results suggest that SIRV3 gp02
147 specifically inhibits subtype I-D CRISPR-Cas activity.

148 To validate the anti-CRISPR subtype I-D activity of SIRV3 gp02, we compared the SIRV2M and
149 SIRV2Mgp02 infectivity in a LAL14/1 CRISPR array deletion mutant ($\Delta arrays$) carrying a
150 plasmid-borne mini-array (a single spacer flanked by two repeats). In this setup, the plasmid
151 T_{gp18}GTG carries a subtype I-D mini-array targeting the template strand of *gp18* and the spacer
152 matches the viral protospacer perfectly with a functional subtype I-D PAM (GTN)₃₃ (Fig. 4b). The
153 control plasmid, T_{gp18}AAA, contains a spacer matching a protospacer shifted by several base-pairs
154 in the viral genome thereby changing the sequence at the PAM position to AAA, which is not
155 functional in CRISPR-Cas immunity. The two constructs were transformed into the LAL14/1
156 $\Delta arrays$ strain, and transformants were tested for viral infectivity. Whereas both SIRV2M and
157 SIRV2Mgp02 were infectious in cells containing the control plasmid T_{gp18}AAA (Fig. 4b), only
158 SIRV2Mgp02 was able to productively infect the cells carrying the functional plasmid-borne
159 subtype I-D CRISPR mini-array (T_{gp18}GTG) (Fig. 4b). A similar experiment using mini-arrays
160 targeting another SIRV2 gene (*gp44*) generated the same result (Supplementary Fig. 3). In summary,
161 our results strongly support the conclusion that SIRV3 gp02 is an anti-CRISPR protein targeting
162 subtype I-D, and this protein was accordingly designated AcrID1 (Anti-CRISPR against subtype I-
163 D).

164 **AcrID1 interacts directly with the Cas10d subunit of the subtype I-D effector complex *in vitro***

165 The subtype I-D interference gene cassette of *S. islandicus* LAL14/1 contains 4 genes, *cas3'*, *csc1*
166 (*cas5d*), *csc2* (*cas7d*), and *cas10d20*. Subtype I-D is an unusual CRISPR-Cas variant that appears to
167 be a hybrid between type I and type III encoding the signature proteins of both types, i.e. a distinct
168 variant of Cas3 (type I hallmark) and a variant of Cas10 (type III hallmark), with the latter
169 containing an inactivated Palm domain but an active HD nuclease domain³⁴. Thus, in subtype I-D,
170 the Cas10d protein is predicted to function as the large subunit of the effector complex and the
171 target-cleaving nuclease but not as part of a signal transduction pathway as shown for type III
172 systems^{35,36}

173 In order to detect a potential direct interaction between AcrID1 and the I-D system, recombinant
174 proteins of the LAL14/1 I-D effector complex were expressed in *E. coli* with a C-terminal His-tag.
175 Using this method, Cas10d and Csc2 could be purified in isolation whereas Cas3' and Csc1 did not
176 express. Heat-treated (78°C) *E. coli* lysates containing Cas10d or Csc2 were incubated separately
177 with a heat-treated (85°C) lysate containing untagged SIRV3 AcrID1 protein and a Ni-NTA pull-
178 down was carried out (Supplementary Fig. 4A). AcrID1 was clearly detected by Western blotting in
179 the eluate fraction with Cas10d, but not in the eluate with Csc2 (Supplementary Fig. 4B) suggesting
180 that AcrID1 interacts specifically with Cas10d. Next, we purified AcrID1 and Cas10d separately to
181 homogeneity, mixed the proteins in an approximate equimolar ratio and analyzed the individual
182 proteins and the complex separately by gel filtration chromatography (Fig. 4c). AcrID1 elutes with
183 an apparent molecular weight of ~30 kDa, suggesting it is a dimer, whereas isolated Cas10d appears
184 as a monomer of approximately ~90 kDa (Fig. 4c, green). Most significantly, AcrID1 and Cas10d
185 co-eluted from the column as a large, stable complex (Fig. 4c and 4d) with an apparent molecular
186 weight of ~220 kDa (Fig. 4c). Together, these results strongly suggest that AcrID1 binds to Cas10d
187 in a 2:2 molecular ratio, possibly through a dimer of AcrID1 tethering two molecules of Cas10d.
188 Thus, we conclude that AcrID1 specifically interacts with the subtype I-D Cas10d protein, the large

189 subunit of the effector complex, and that this interaction blocks the CRISPR-Cas response allowing
190 virus propagation to continue. Given that Cas10d is unique for subtype I-D, these findings also
191 imply that the inhibitory activity of AcrID1 is highly specific for this CRISPR-Cas variant.

192 **AcrID1 homologs in archaeal viruses**

193 Protein sequence database searches (see Methods for details) identified about 50 proteins
194 homologous to AcrID1 encoded by viruses that infect hyperthermophilic crenarchaeota of the order
195 Sulfolobales, including rudiviruses, lipothrixviruses, fuselloviruses and monocaudaviruses
196 (Supplementary Fig. 5). The sequences of these proteins typically show at least 25% identity with
197 highly significant expectation values ($e < 10^{-5}$) after one to three PSI-BLAST iterations, depending
198 on the virus family, supporting the conclusion on their homology. Examination of the multiple
199 alignment of the AcrID1 homolog sequences shows conservation of multiple hydrophobic as well
200 as charged amino acid residues (Supplementary Fig. 5 and below).

201 The phylogenetic tree of the AcrID1 protein family splits into two branches, with the majority
202 clustering with SIRV3 AcrID1 (the SIRV2 gp03 group) and less than one third clustering with
203 SIRV2 gp02 (Fig. 5, Supplementary Fig. 5, and Supplementary Fig. 6). There is a notable
204 expansion of the family in some of the archaeal viruses, in particular, SIRV2, which encodes 12
205 paralogous family members (Fig. 5 and Supplementary Fig. 6). Despite the small size of the
206 proteins that limits the resolution of phylogenetic analysis, the tree contains several well-supported
207 branches with varying numbers of family members from different viruses (Fig. 5). This pattern
208 implies a complex evolutionary history of the Acr family that seems to include multiple gene
209 duplications, losses, and transfer of Acr genes between viruses.

210 Similarly to the previously characterized bacteriophage Acrs, the *acrID1*-related genes of SIRV1,
211 SIRV2 and SIRV3 are all located adjacent to genes encoding a helix-turn-helix (HTH) DNA-

212 binding domain (Supplementary Fig. 6). In particular, *acrID1* is adjacent to SIRV2 gp01 gene
213 which encodes a HTH domain-containing protein³⁷. However, many viruses of Sulfolobales,
214 including multiple isolates of rudiviruses³⁸, lack detectable members of the AcrID1 family.
215 Nevertheless, most of these viruses possess genes for gp01 homologs located upstream of arrays of
216 genes encoding uncharacterized small proteins that could be Acrs unrelated to AcrID1. Collectively,
217 these findings suggest that viruses of Sulfolobales encode multiple Acr proteins, many of which
218 belong to the AcrID1 family, whereas others appear to have unrelated origins. In all cases, the
219 HTH-containing proteins most likely regulate the expression of the Acr genes. Previous
220 transcriptomic analyses have shown early, high-level transcription of the SIRV2 genomic termini
221 including *gp01* and the *acrID1*³⁹. This finding suggests that, because CRISPR-Cas response occurs
222 early during viral infection, anti-CRISPR activities are needed immediately post infection.

223 **SIRV3 AcrID1 displays a compact, dimeric $\alpha\beta$ -fold with a tendency to form fibers**

224 His-tagged SIRV3 AcrID1 (gp02) was crystallized and the structure determined to 2 Å using Se-
225 Met substitution (see Supplementary Table 2 and Methods for details). The structure contains 6
226 molecules of AcrID1 in the asymmetric unit (ASU) and has R = 24.1% (R_{free} = 26.9%). AcrID1 is a
227 compact $\alpha\beta$ -sandwich protein comprised of a five-stranded, anti-parallel β sheet and two α -helices,
228 both located on the same side of the sheet (Fig. 6a). The first β -strand of the sheet is broken in two
229 ($\beta 1$ and $\beta 1'$) and the topology can therefore appropriately be described as $\alpha 1$ - $\beta 1$ - $\beta 1'$ - $\alpha 2$ - $\beta 2$ - $\beta 3$ - $\beta 4$ - $\beta 5$.
230 All six molecules in the ASU engage in identical dimers, thus compatible with the observed
231 behavior of the protein in solution (Fig. 4c). The dimer interaction is head-to-tail, with the helices
232 on the same side of the sheet and the two $\beta 1$ strands adjacent to each other (Fig. 6b). The dimer has
233 a total surface area of 10,920 Å² of which about 1900 Å² are buried at the interface. Interactions
234 between the monomers involve main chain hydrogen bonds between the $\beta 1$ strands, and the dimer
235 thus effectively consists of an extended, 10-stranded anti-parallel β -sheet structure, which is likely

the biologically relevant form of the protein. The dimer interface is further stabilized by hydrophobic interactions in the region between the β -sheet and the helices and polar interactions between the helices themselves (Fig. 6b). In particular, the hydrophobic interactions, Ile19-Phe26 and Ile22-Leu24, which are present twice at the interface due to the two-fold symmetry, as well as Met9-Met9, involve residues that are highly conserved among the homologous archaeal virus proteins (Fig. 6c and S5). Furthermore, the protein contains a range of conserved surface residues, including Glu21, Lys34, Tyr55, Glu81, Arg92, and Trp91, which might be involved in protein-protein interactions (Fig. 6c). Packing of dimers in the crystal extends the β -sheet further, in this case with a 180° rotation perpendicular to the strands and thus with the helices on alternate sides (Supplementary Fig. 7). Although this interaction appears weaker than the dimer interaction, the continuous packing of β -sheets results in the crystal effectively consisting of long, parallel fibers of protein (Supplementary Fig. 7).

The protein databank contains two other structures of archaeal viral proteins with unknown function that are structurally highly similar to AcrID1, *S. islandicus* Filamentous Virus ORF14 (SIFV14, PDB ID 2H36^{40,41}) and *Acidianus* Filamentous Virus 1 ORF-99 (AFV1-99, PDB IDs 3DJW and 3DF6⁴²) (Fig. 6d). These proteins share 30% and 22% identity, respectively, with SIRV3 AcrID1, and both structures display highly similar $\alpha\beta$ -folds and form dimers as well as extended fiber-like structures in the crystal. These consistent properties of AcrID1 homologs suggest that the tendency to dimerize and polymerize could be a biologically relevant feature of this protein family. Calculation of the surface electrostatic potential shows that SIRV3 AcrID1 as well as the two structurally similar proteins are highly negatively charged, suggesting that they might act as DNA mimics as previously proposed for some of the bacteriophage Acrs⁴³ (Fig S8). This negative charge density appears to be particularly pronounced in SIRV3 AcrID1. For AFV1-99, extreme acid and temperature stability, so that the native structures is maintained even at pH 0 and 95°C⁴². Thus,

260 SIRV3 AcrID1 contains a structural fold that is conserved in a broad range of archaeal virus
261 proteins and is likely to be highly pH and temperature stable.

262 The AcrID1 family is represented in archaeal viruses from four diverse families. The wide
263 distribution and the apparent abundance of Acr proteins in the viruses of Sulfolobales correlates
264 with the high tolerance of *Sulfolobus* CRISPR-Cas systems to protospacer mismatches²³, which
265 could make these viruses strongly dependent on Acrs rather than point mutations to circumvent
266 CRISPR-Cas-mediated host defense. The presence of as many as 12 *acrID1* paralogs in SIRV2 as
267 well as a smaller number of AcrID1 family members in other viruses of Sulfolobales
268 (Supplementary Fig. 6), is intriguing. Whether these Acrs function independently or cooperatively,
269 and whether they target different CRISPR-Cas subtypes within a single host or the same subtypes in
270 different hosts, will be the focus of future studies. Moreover, the I-A and III-B systems in LAL14/1
271 do not provide immunity against SIRV2 despite the presence of multiple spacers matching the virus
272 genome (with 2 to 6 mismatches) in the respective CRISPR arrays^{20,39}. Thus, some of the AcrID1
273 paralogs or other putative Acrs encoded by SIRV2 are likely to inhibit these CRISPR-Cas subtypes.

274 **METHODS**

275 **Strains and growth conditions**

276 The hyperthermophilic archaeal host *S. solfataricus* 5E6₂₆ carries CRISPR loci A-D deletion, *S.*
277 *islandicus* LAL549 is a Δ *pyrF* clone derived from *S. islandicus* LAL14/1 based on which Δ arrays⁴⁴
278 and Δ *cas6* were constructed. *Sulfolobus* cells were grown at 78°C in 50 mL GCVY medium (basic
279 salts medium⁴⁵ was supplemented with 0.2% glucose, 0.2% casamino acids, 0.005% yeast extract
280 and a vitamin mixture). Uracil (20 µg/ml) was added to the medium for the cultivation of the Δ *pyrF*
281 strains. To induce SIRV3 gp02 expression from the arabinose promoter (Fig. 3D), we grew cultures
282 in ACVY medium (0.2% arabinose was substituted for glucose). *E. coli* DH5 α and Rosetta were
283 used for cloning and recombinant protein expression, respectively.

284 **Plasmid construction**

285 Gene fragments were amplified by PCR or fusion PCR using primers listed in Supplementary Table
286 1, digested with two restriction enzymes (Supplementary Table 1) and ligated into pEXA2 (for
287 SIRV3 gp02 and *cas6* expression) linearized with the same set of enzymes, or ligated to blunt end
288 pJET1.2 directly. For the construction of pJET1.2/flank2223-SIRV3frag plasmids (Fig. 2B), we
289 cloned homologous flanking arms into pJET1.2 to make pJET1.2/flank2223, then inserted different
290 SIRV3 fragments into pJET1.2/flank2223 which were both digested with *NheI* and *KpnI*. The
291 ligated DNAs were then transformed into *E. coli* DH5 α and plasmid clones were purified using
292 Plasmid DNA Mini Kit II (D6945-02; Omega).

293 To construct the mini-array containing plasmids (Fig. 5), the spacer DNA was prepared by
294 annealing two complementary oligonucleotides and the resultant dsDNA carrying 3 nt overhangs
295 was inserted into the *SapI* site of pGE244.

296 **Construction of gene knockout mutants**

297 Type I-A and type I-D *cas6* were deleted from *S. islandicus* LAL549 based on the marker insertion
298 and unmarked target gene deletion method (MID)⁴⁶ using plasmids pMID-*cas6*⁴⁷ and pMID-*cas6*(I-
299 D) (Supplementary Table 1), respectively. Gene deletion in Δ *cas6*(I-A) and Δ *cas6*(I-D) was
300 confirmed (Supplementary Fig. 2) by PCR with primers listed in Supplementary Table 1.

301 **Electroporation procedures**

302 Electroporation was used to transform *Sulfolobus* strains according to the procedure described
303 previously⁴⁸ with the following modifications: (1) all manipulations were carried out at room
304 temperature; (2) after electroporation of 500 ng DNA into 50 μ L *Sulfolobus* competent cells (OD₆₀₀
305 around 10), the cells were immediately transferred into 950 μ L pre-warmed basic salts medium pre-
306 mixed with SIRV2M (multiplicity of infection, 1) and incubated at 75°C for 30 min without

307 shaking; (3) all cells were transferred into 50 mL SCVU medium (0.2% sucrose was substituted for
308 glucose) and grown at 78°C.

309 **Plaque assay**

310 Three mL *S. islandicus* LAL14/1 Δ arrays cells (OD₆₀₀ around 0.2) were premixed with 100 μ L
311 serial dilutions of viruses and incubated at 78°C for 30 min without shaking. The infected cultures
312 were then mixed with 1 ml 0.4% Gelrite and plated onto pre-warmed Gelrite plates containing SCV
313 medium. After the top layer was set, the plates were put into a plastic box, tightly closed and
314 incubated at 78°C for two days. The plaque forming units (PFU) were calculated from the plates
315 with a proper number of plaques.

316 **Cloning and protein expression**

317 *acrID1*, *cas10d* and *csc2* were cloned into pET30a (Novogene) between the *NdeI* and *XhoI*
318 restriction sites, resulting in constructs encoding proteins with or without a C-terminal His tag.
319 Overlap extension PCR (fusion PCR) was used to introduce a silent mutation in a natural *NdeI*
320 restriction site in the coding sequence of *Cas10d*. The primers applied for construction of the
321 vectors are listed in Supplementary Table 1. *Csc2* was expressed in Rosetta™ cells (Novogene)
322 cultivated in Luria-Bertani (LB) medium and induced with 0.5 mM isopropyl β -D-1-
323 thiogalactopyranoside (IPTG) at 15°C. *AcrID1* and *Cas10d* were expressed in *E. coli* BL21 DE3
324 Rosetta2 as C-terminal His-tagged fusion proteins. Cells were grown in LB medium supplemented
325 with 50 μ g/ml kanamycin and 34 μ g/ml chloramphenicol at 37°C in a shaking incubator until OD₆₀₀
326 reached 0.8 and induced by addition of 0.5-1 mM IPTG. Expression was allowed to continue
327 overnight at 18-20°C in a shaking incubator. Cells were harvested by centrifugation (12.000 g/15
328 mins), washed in ice water and stored at -20°C.

329 **Purification of AcrID1 and Cas10d**

330 AcrID1 and Cas10d were purified by similar protocols. Cells were resuspended in lysis buffer (50
331 mM Tris-Cl, pH 8, 300 mM NaCl, 5 mM MgCl₂, 10% glycerol, 20 mM imidazole, 3 mM β-
332 mercaptoethanol (BME), 1 mM phenylmethylsulfonyl fluoride (PMSF), and 10 µg/ml DNase I) and
333 lyzed by sonication. After centrifugation to remove cell debris, the supernatant was incubated for 20
334 mins at 78°C to precipitate the majority of *E. coli* proteins, clarified again, and loaded on a 5 mL
335 HisTrap HP column (GE Healthcare). After washing in several volumes of lysis buffer including 30
336 mM imidazole the protein was eluted in lysis buffer including 250 mM imidazole. The eluted
337 protein was then diluted and loaded on a 1 mL SourceQ column (GE Healthcare) equilibrated in
338 buffer A (50 mM Tris-Cl, pH 8, 5 mM BME) and eluted using a linear gradient into 50 mM Tris-Cl,
339 pH 8, 1 M NaCl, and 5 mM BME (buffer B). Peak fractions were concentrated using a Vivaspinn
340 centrifugal filter (5 kDa MWCO, Sartorius) and loaded on a Superdex 75 10/300 GL column (GE
341 Healthcare) equilibrated in 25 mM Hepes, pH 7, 150 mM NaCl (GF buffer). Finally, the protein
342 sample was concentrated to 10-20 mg/ml using Vivaspinn filters (5 kDa MWCO) before
343 crystallization.

344 Se-Met substituted AcrID1 was produced in *E. coli* 834 DE3 (Met auxotroph) by growing to OD₆₀₀
345 = 0.6-0.8 in 500 mL LB medium, then pelleting the cells (11.000 g, 15 min) and resuspending them
346 in 30 mL SelenoMet Medium Base (MD12-501, Molecular Dimensions) and transferring 10 mL of
347 this solution into 2 L SelenoMet Medium Base + SelenoMet Nutrient Mix (MD12-502, Molecular
348 Dimensions) as described by the manufacturer, supplemented with 50 µg/µl kanamycin and allowed
349 to grow for 3 h at 37°C in a shaking incubator. Then, 4 mL of a 10 mg/ml stock solution of seleno-
350 methionine was added and the cells allowed to grow until OD₆₀₀ reached 0.8 at which point
351 expression was induced with 1 mM IPTG. The remaining expression and purification protocol was
352 identical to the native protein.

353 **Analysis of Cas10d-AcrID1 interaction *in vitro***

354 For analysis of protein interaction *in vitro*, approximately 20 μ l of a 3.5 mg/ml sample of pure His-
355 tagged AcrID1 was mixed with pure His-tagged Cas10d in 1:1 or 2:1 molar ratios and incubated for
356 90 seconds at 78°C, spun 5 minutes at 13 krpm in a benchtop centrifuge to remove aggregates and
357 separated on a 24 mL Superdex200 increase 10/300 GL column (GE Healthcare) equilibrated in 25
358 mM Hepes-KOH, pH 7.5, 150 mM NaCl, and 5 mM BME. Samples of the isolated proteins were
359 analyzed separately as control.

360 **Crystallization and structure determination of AcrID1**

361 Single crystals were identified using the JCSG+ screen (Molecular Dimensions) in sitting drops
362 containing 0.2 μ l protein + 0.2 μ l reservoir solution (18-20% (w/v) PEG 8000, 0.2 M MgCl₂, 0.1 M
363 Tris-Cl, pH 8.5) in trays that had been incubated 3 days at 37°C and then transferred to 19°C for
364 several days. Crystals of the Se-Met substituted protein appeared in similar conditions but were
365 improved by streak-seeding from smaller crystals into clear drops. Crystals were cryo-protected by
366 quick transfer through a buffer similar to the reservoir but including 40% PEG 8000, and frozen in
367 liquid N₂ before data collection. Complete x-ray diffraction data were collected at beamline P13 at
368 the EMBL PETRA synchrotron, Hamburg from single native and Se-Met substituted crystals, in the
369 latter case at the peak absorption wavelength for Se, 0.97947 Å (Supplementary Table 2).

370 Diffraction images were integrated using the xia2/XDS pipeline and analyzed using CCP4
371 (POINTLESS)^{49,50} and Phenix.xtriage⁵¹. The crystals belong to the monoclinic space group P12₁1
372 and analysis of the native Patterson function reveals that they suffer from pseudo-centric
373 translational non-crystallographic symmetry (peak near $u,v,w=0.5, 0.5, 0.5$), which results in all
374 reflections with odd numbered sum of indices ($h+k+l = 2n+1$) to have a significantly lower value
375 than the other half of the reflections. Despite this issue, phasing was carried out using the SHELXD
376 hkl2map pipelines⁵², which also produced a good initial map and model. This initial model was

improved in BUCCANEER⁵³ before being manually adjusted and improved in Coot⁵⁴. Initial refinement was carried out in Phenix.refine using a standard protocol including positional and individual B factor refinement and NCS restraints. The partly refined model was transferred to the native data set using the same 5% free set of reflections as in the Se-Met data, and iterative model building and refinement continued. The final model consists of three dimers (6 molecules) per ASU, and shows good statistics but high refinement residuals ($R/R_{\text{free}}=24.1\%/26.9\%$) due to the relationships between reflections resulting from the translational NCS. Structure coordinates have been deposited in the Protein Data Bank with ID 6EXP.

Bioinformatics analysis

All small proteins (<150 amino acids in length) encoded in the vicinity of SIRV2 gp01 and its close homolog gp54 (encoded on the opposite end of the genome) were used as queries for PSI-BLAST⁵⁵ search against the virus subset of non-redundant NCBI protein database (E-value cutoff 0.1, composition based statistics and low complexity filtering turned off; the search was run until convergence). In cases when the sets of sequences retrieved in these searches overlapped, their union was taken for further analysis. Additional searches for potential distant homologs were performed using the HHpred software⁵⁶ seeded with profiles obtained from multiple alignments of the respective proteins which were constructed using MUSCLE⁵⁷. The multiple alignment of the homologs of SIRV2 gp03 (SIRV3 gp02) was used for phylogenetic tree reconstruction with the FastTree program⁵⁸, with the WAG evolutionary model and the discrete gamma model with 20 rate categories. The same program was used to compute support values. The alignment in Supplementary Fig. 5 was colored using CLUSTAL X⁵⁹.

Data availability Accession codes for protein sequence data presented herein are provided in Fig. 6. The AcrID1 accession code is 6EXP.

400 **References**

- 401 1 Koonin, E. V., Makarova, K. S. & Wolf, Y. I. Evolutionary Genomics of Defense Systems
402 in Archaea and Bacteria. *Annu Rev Microbiol*, doi:10.1146/annurev-micro-090816-093830
403 (2017).
- 404 2 Mohanraju, P. *et al.* Diverse evolutionary roots and mechanistic variations of the CRISPR-
405 Cas systems. *Science* **353**, aad5147, doi:10.1126/science.aad5147 (2016).
- 406 3 Barrangou, R. *et al.* CRISPR provides acquired resistance against viruses in prokaryotes.
407 *Science* **315**, 1709-1712, doi:10.1126/science.1138140 (2007).
- 408 4 Koonin, E. V., Makarova, K. S. & Zhang, F. Diversity, classification and evolution of
409 CRISPR-Cas systems. *Curr Opin Microbiol* **37**, 67-78, doi:10.1016/j.mib.2017.05.008
410 (2017).
- 411 5 Brouns, S. J. *et al.* Small CRISPR RNAs guide antiviral defense in prokaryotes. *Science* **321**,
412 960-964, doi:10.1126/science.1159689 (2008).
- 413 6 Charpentier, E., Richter, H., van der Oost, J. & White, M. F. Biogenesis pathways of RNA
414 guides in archaeal and bacterial CRISPR-Cas adaptive immunity. *FEMS microbiology*
415 *reviews* **39**, 428-441, doi:10.1093/femsre/fuv023 (2015).
- 416 7 Sternberg, S. H., Richter, H., Charpentier, E. & Qimron, U. Adaptation in CRISPR-Cas
417 Systems. *Molecular cell* **61**, 797-808, doi:10.1016/j.molcel.2016.01.030 (2016).
- 418 8 Jackson, S. A. *et al.* CRISPR-Cas: Adapting to change. *Science* **356**,
419 doi:10.1126/science.aal5056 (2017).
- 420 9 Plagens, A., Richter, H., Charpentier, E. & Randau, L. DNA and RNA interference
421 mechanisms by CRISPR-Cas surveillance complexes. *FEMS microbiology reviews* **39**, 442-
422 463, doi:10.1093/femsre/fuv019 (2015).
- 423 10 Maxwell, K. L. *et al.* The solution structure of an anti-CRISPR protein. *Nature*
424 *communications* **7**, 13134, doi:10.1038/ncomms13134 (2016).
- 425 11 Rauch, B. J. *et al.* Inhibition of CRISPR-Cas9 with Bacteriophage Proteins. *Cell* **168**, 150-
426 158.e110, doi:10.1016/j.cell.2016.12.009 (2017).
- 427 12 Borges, A. L., Davidson, A. R. & Bondy-Denomy, J. The Discovery, Mechanisms, and
428 Evolutionary Impact of Anti-CRISPRs. *Annu Rev Virol*, doi:10.1146/annurev-virology-
429 101416-041616 (2017).
- 430 13 Pawluk, A., Davidson, A. R. & Maxwell, K. L. Anti-CRISPR: discovery, mechanism and
431 function. *Nat Rev Microbiol* **16**, 12-17, doi:10.1038/nrmicro.2017.120 (2018).

432 14 Bondy-Denomy, J., Pawluk, A., Maxwell, K. L. & Davidson, A. R. Bacteriophage genes
433 that inactivate the CRISPR/Cas bacterial immune system. *Nature* **493**, 429-432,
434 doi:10.1038/nature11723 (2013).

435 15 Pawluk, A., Bondy-Denomy, J., Cheung, V. H., Maxwell, K. L. & Davidson, A. R. A new
436 group of phage anti-CRISPR genes inhibits the type I-E CRISPR-Cas system of
437 *Pseudomonas aeruginosa*. *MBio* **5**, e00896, doi:10.1128/mBio.00896-14 (2014).

438 16 Pawluk, A. *et al.* Naturally Occurring Off-Switches for CRISPR-Cas9. *Cell* **167**, 1829-1838
439 e1829, doi:10.1016/j.cell.2016.11.017 (2016).

440 17 Pawluk, A. *et al.* Inactivation of CRISPR-Cas systems by anti-CRISPR proteins in diverse
441 bacterial species. *Nat Microbiol* **1**, 16085, doi:10.1038/nmicrobiol.2016.85 (2016).

442 18 Hynes, A. P. *et al.* An anti-CRISPR from a virulent streptococcal phage inhibits
443 *Streptococcus pyogenes* Cas9. *Nat Microbiol* **2**, 1374-1380, doi:10.1038/s41564-017-0004-7
444 (2017).

445 19 Prangishvili, D. *et al.* The enigmatic archaeal virosphere. *Nat Rev Microbiol* **15**, 724-739,
446 doi:10.1038/nrmicro.2017.125 (2017).

447 20 Jaubert, C. *et al.* Genomics and genetics of *Sulfolobus islandicus* LAL14/1, a model
448 hyperthermophilic archaeon. *Open biology* **3**, 130010, doi:10.1098/rsob.130010 (2013).

449 21 Wiedenheft, B. *et al.* RNA-guided complex from a bacterial immune system enhances target
450 recognition through seed sequence interactions. *Proc Natl Acad Sci U S A* **108**, 10092-10097,
451 doi:10.1073/pnas.1102716108 (2011).

452 22 Semenova, E. *et al.* Interference by clustered regularly interspaced short palindromic repeat
453 (CRISPR) RNA is governed by a seed sequence. *Proc Natl Acad Sci U S A* **108**, 10098-
454 10103, doi:10.1073/pnas.1104144108 (2011).

455 23 Manica, A., Zebec, Z., Steinkellner, J. & Schleper, C. Unexpectedly broad target recognition
456 of the CRISPR-mediated virus defence system in the archaeon *Sulfolobus solfataricus*.
457 *Nucleic Acids Res* **41**, 10509-10517, doi:10.1093/nar/gkt767 (2013).

458 24 Mousaei, M., Deng, L., She, Q. & Garrett, R. A. Major and minor crRNA annealing sites
459 facilitate low stringency DNA protospacer binding prior to Type I-A CRISPR-Cas
460 interference in *Sulfolobus*. *RNA biology*, 0, doi:10.1080/15476286.2016.1229735 (2016).

461 25 Bize, A. *et al.* A unique virus release mechanism in the Archaea. *Proc Natl Acad Sci U S A*
462 **106**, 11306-11311, doi:10.1073/pnas.0901238106 (2009).

463 26 Okutan, E. *et al.* Novel insights into gene regulation of the rudivirus SIRV2 infecting
464 Sulfolobus cells. *RNA biology* **10**, 875-885, doi:10.4161/rna.24537 (2013).

465 27 Deng, L. *et al.* Unveiling cell surface and type IV secretion proteins responsible for archaeal
466 rudivirus entry. *J Virol* **88**, 10264-10268, doi:10.1128/JVI.01495-14 (2014).

467 28 He, F., Chen, L. & Peng, X. First experimental evidence for the presence of a CRISPR toxin
468 in sulfolobus. *J Mol Biol* **426**, 3683-3688, doi:10.1016/j.jmb.2014.09.016 (2014).

469 29 Guo, Y., Kragelund, B. B., White, M. F. & Peng, X. Functional characterization of a
470 conserved archaeal viral operon revealing single-stranded DNA binding, annealing and
471 nuclease activities. *J Mol Biol* **427**, 2179-2191, doi:10.1016/j.jmb.2015.03.013 (2015).

472 30 Martinez-Alvarez, L., Bell, S. D. & Peng, X. Multiple consecutive initiation of replication
473 producing novel brush-like intermediates at the termini of linear viral dsDNA genomes with
474 hairpin ends. *Nucleic Acids Res* **44**, 8799-8809, doi:10.1093/nar/gkw636 (2016).

475 31 Martinez-Alvarez, L., Deng, L. & Peng, X. Formation of a Viral Replication Focus in
476 Sulfolobus Cells Infected by the Rudivirus Sulfolobus islandicus Rod-Shaped Virus 2. *J*
477 *Virol* **91**, doi:10.1128/JVI.00486-17 (2017).

478 32 Erdmann, S., Le Moine Bauer, S. & Garrett, R. A. Inter-viral conflicts that exploit host
479 CRISPR immune systems of Sulfolobus. *Molecular microbiology* **91**, 900-917,
480 doi:10.1111/mmi.12503 (2014).

481 33 Shah, S. A., Erdmann, S., Mojica, F. J. & Garrett, R. A. Protospacer recognition motifs:
482 mixed identities and functional diversity. *RNA biology* **10**, 891-899, doi:10.4161/rna.23764
483 (2013).

484 34 Makarova, K. S. *et al.* Evolution and classification of the CRISPR-Cas systems. *Nat Rev*
485 *Microbiol* **9**, 467-477, doi:10.1038/nrmicro2577 (2011).

486 35 Niewoehner, O. *et al.* Type III CRISPR-Cas systems produce cyclic oligoadenylate second
487 messengers. *Nature* **548**, 543-548, doi:10.1038/nature23467 (2017).

488 36 Kazlauskienė, M., Kostiuk, G., Venclovas, C., Tamulaitis, G. & Siksnys, V. A cyclic
489 oligonucleotide signaling pathway in type III CRISPR-Cas systems. *Science* **357**, 605-609,
490 doi:10.1126/science.aao0100 (2017).

491 37 Peeters, E. *et al.* DNA-Interacting Characteristics of the Archaeal Rudiviral Protein
492 SIRV2_Gp1. *Viruses* **9**, doi:10.3390/v9070190 (2017).

493 38 Bautista, M. A., Black, J. A., Youngblut, N. D. & Whitaker, R. J. Differentiation and
494 Structure in *Sulfolobus islandicus* Rod-Shaped Virus Populations. *Viruses* **9**,
495 doi:10.3390/v9050120 (2017).

496 39 Quax, T. E. *et al.* Massive activation of archaeal defense genes during viral infection. *J Virol*
497 **87**, 8419-8428, doi:10.1128/JVI.01020-13 (2013).

498 40 Goulet, A. *et al.* Crystallization and preliminary X-ray diffraction analysis of protein 14
499 from *Sulfolobus islandicus* filamentous virus (SIFV). *Acta Crystallogr Sect F Struct Biol*
500 *Cryst Commun* **62**, 884-886, doi:10.1107/S1744309106029150 (2006).

501 41 Goulet, A. *et al.* The crystal structure of ORF14 from *Sulfolobus islandicus* filamentous
502 virus. *Proteins* **76**, 1020-1022, doi:10.1002/prot.22448 (2009).

503 42 Goulet, A. *et al.* The thermo- and acido-stable ORF-99 from the archaeal virus AFV1.
504 *Protein Sci* **18**, 1316-1320, doi:10.1002/pro.122 (2009).

505 43 Shin, J. *et al.* Disabling Cas9 by an anti-CRISPR DNA mimic. *Sci Adv* **3**, e1701620,
506 doi:10.1126/sciadv.1701620 (2017).

507 44 He, F., Vestergaard, G., Peng, W., She, Q. & Peng, X. CRISPR-Cas type I-A Cascade
508 complex couples viral infection surveillance to host transcriptional regulation in the
509 dependence of Csa3b. *Nucleic Acids Res* **45**, 1902-1913, doi:10.1093/nar/gkw1265 (2017).

510 45 Zillig, W. *et al.* Screening for *Sulfolobales*, their Plasmids and their Viruses in Icelandic
511 Solfataras. *Systematic and Applied Microbiology* **16**, 609-628, doi:10.1016/s0723-
512 2020(11)80333-4 (1993).

513 46 Zhang, C. *et al.* Revealing the essentiality of multiple archaeal pcna genes using a mutant
514 propagation assay based on an improved knockout method. *Microbiology* **156**, 3386-3397,
515 doi:10.1099/mic.0.042523-0 (2010).

516 47 Peng, W. *et al.* Genetic determinants of PAM-dependent DNA targeting and pre-crRNA
517 processing in *Sulfolobus islandicus*. *RNA biology* **10**, 738-748, doi:10.4161/rna.23798
518 (2013).

519 48 Deng, L., Zhu, H., Chen, Z., Liang, Y. X. & She, Q. Unmarked gene deletion and host-
520 vector system for the hyperthermophilic crenarchaeon *Sulfolobus islandicus*. *Extremophiles*
521 **13**, 735-746, doi:10.1007/s00792-009-0254-2 (2009).

522 49 Evans, P. Scaling and assessment of data quality. *Acta Crystallogr D Biol Crystallogr* **62**,
523 72-82, doi:10.1107/S0907444905036693 (2006).

- 524 50 Winn, M. D. *et al.* Overview of the CCP4 suite and current developments. *Acta Crystallogr*
525 *D Biol Crystallogr* **67**, 235-242, doi:10.1107/S0907444910045749 (2011).
- 526 51 Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular
527 structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221,
528 doi:10.1107/S0907444909052925 (2010).
- 529 52 Pape, T. & Schneider, T. R. HKL2MAP: a graphical user interface for macromolecular
530 phasing with SHELX programs. *Journal of Applied Crystallography* **37**, 843-844,
531 doi:10.1107/S0021889804018047 (2004).
- 532 53 Cowtan, K. Fitting molecular fragments into electron density. *Acta Crystallogr D Biol*
533 *Crystallogr* **64**, 83-89, doi:10.1107/S0907444907033938 (2008).
- 534 54 Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta*
535 *Crystallogr D Biol Crystallogr* **60**, 2126-2132, doi:10.1107/S0907444904019158 (2004).
- 536 55 Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein database
537 search programs. *Nucleic Acids Res* **25**, 3389-3402 (1997).
- 538 56 Soding, J., Biegert, A. & Lupas, A. N. The HHpred interactive server for protein homology
539 detection and structure prediction. *Nucleic Acids Res* **33**, W244-248, doi:10.1093/nar/gki408
540 (2005).
- 541 57 Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
542 throughput. *Nucleic Acids Res* **32**, 1792-1797, doi:10.1093/nar/gkh340 (2004).
- 543 58 Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood
544 trees for large alignments. *PloS one* **5**, e9490, doi:10.1371/journal.pone.0009490 (2010).
- 545 59 Larkin, M. A. *et al.* Clustal W and Clustal X version 2.0. *Bioinformatics* **23**, 2947-2948,
546 doi:10.1093/bioinformatics/btm404 (2007).

547 **Acknowledgments**

548 The authors thank Mart Krupovic, Nick Grishin, David Prangishvili, Qunxin She and Roger A.
549 Garrett for useful discussions, Regitze Bertelsen for help with protein purification and
550 crystallization, and Prof. Gregers Rom Andersen and beamline staff at the P13 beamline at PETRA,
551 Hamburg, for help with data collection. This work was supported by EU FP7 project HotZyme
552 [265933] and Danish Council for Independent Research/Technology and Production (grant number
553 DFF-7017-00060) to XP and the Danish National Research Foundation's Centre for Bacterial Stress

554 Response and Persistence (BASP, grant no. DNR120) to DEB. KSM and EVK are supported by
555 intramural funds of the US Department of Health and Human Resources (to the National Library of
556 Medicine).

557

558 **Author contributions**

559 FH and XP conceived the experiments. FH, YBC, LBV, ALK, MD performed the experiments.
560 DEB analysed the crystal structure of AcrID1. KSM and EVK performed phylogenetic analysis of
561 the AcrID homologs. FH, YBC, ALK, EVK, DEB and XP wrote the paper which was read and
562 approved by all authors.

563

564 **Competing financial interests statement**

565 None

566

567 **Figure legends**

568 **Fig. 1. Isolation of a SIRV2 mutant, SIRV2M, that is non-infectious in *S. islandicus* LAL14/1.**

569 **a.** Schematic diagram of the left genomic terminus of SIRV2 and the recombination-mediated
570 deletion of a 4kb fragment spanning gp02 – gp09 (grey bar). R, 13 bp repeat. gp01 and gp10 are
571 indicated by block arrows. **b** and **c.** PCR analyses of the region using the supernatant of infected
572 cultures of the strains 5E6 (CRISPR -) and LAL14/1 (CRISPR +) (**b**) or purified SIRV2M (**c**) as
573 template. The annealing positions of the PCR primers are indicated by grey arrows above the
574 genome maps. DNA size ladders (L) with the sizes of selected bands are shown. PCR products
575 derived from the wild-type (wt) SIRV2 or the mutant SIRV2M (m) are indicated by dark arrows. **d.**
576 Growth curves of *S. islandicus* LAL14/1 after viral infection. Results in **b** and **c** are representatives
577 of three experiments. The growth curve in **d** is a representative of a single condition that was
578 confirmed by two independent experiments with different sampling time.

579 **Fig. 2. SIRV3 gp02 is essential and sufficient for restoring SIRV2M infectivity in LAL14/1. a.**

580 Schematic diagrams of left genomic termini of SIRV2 and SIRV3. Genes are depicted by arrows
581 and high sequence homologies (> 50% aa identity) are color-coded. **b.** Schematic illustration of the
582 recombination (double crossover) between SIRV2M and SIRV3 fragments (SIRV3frag) cloned
583 between the recombination arms (purple and light blue boxes) in pJET1.2. **c** and **d.** PCR analysis of
584 the recombination events using primers depicted in **b** as horizontal grey arrows. Plasmid-borne
585 SIRV3 genes that were electroporated into LAL14/1 before adding SIRV2M are indicated on top of
586 the gels. Supernatants of the cultures two days post electroporation (**c**) and two days post dilution of
587 the first culture supernatants presented in panel c into fresh cultures (**d**) were used as DNA template
588 in the PCR reactions. Control 1, water was electroporated into the cells but no SIRV2M was added;
589 control 2, water was electroporated into the cells after which SIRV2M was added. The PCR product
590 derived from SIRV2M is indicated by a black arrow in **c**, and the PCR bands derived from

591 recombinant viruses are labeled with asterisks (**c** and **d**). Results in **c** and **d** are representatives of
592 two experiments.

593 **Fig. 3. SIRV3 gp02 facilitates SIRV2M propagation in LAL14/1.** **a.** Growth curves showing
594 infection of SIRV2M and SIRV2Mgp02 in LAL14/1 cells (WT). No virus was added in the control
595 culture. **b.** Spot assay showing the infectivity of the three viruses. An aliquot of 10 μ l was spotted
596 onto a cell lawn of LAL14/1 after serial dilutions (dilution factors are indicated below the picture)
597 of the viral preparations. The titer of the three vial solutions was measured with a CRISPR null
598 mutant and was adjusted to about 3.8×10^9 PFU/ml (plaque forming units/ml). **c.** Plaque forming
599 units in culture supernatants were measured for both SIRV2M and SIRV2Mgp02 after different
600 hours post infection (hpi) in LAL14/1. Error bars, s.d. of two technical replicates. **d.** SIRV3 gp02
601 was cloned into the pEXA2 vector, transformed into LAL14/1 cells, then infected with SIRV2M.
602 Either glucose or arabinose was added to the media as carbon source with the latter inducing gp02
603 expression. Results in **b** are representatives of three experiments. The growth curves in **a** and **d** are
604 representatives of a single condition that were confirmed by two independent experiments with
605 different sampling time.

606

607 **Fig. 4. SIRV3 gp02 (AcrID1) protein interacts with Cas10d protein (c and d) thereby**
608 **inhibiting CRISPR-Cas subtype I-D immunity (a and b).** **a.** Growth of LAL14/1 $\Delta cas6(I-A)$
609 (left), $\Delta cas6(I-D)$ with (right) and without (middle) a plasmid-borne subtype I-D *cas6*
610 complementation; **b.** Growth of LAL14/1 Δ arrays cells carrying a plasmid-borne spacer matching
611 gp18 of SIRV2 (and hence also of SIRV2M and SIRV2Mgp02) with (T_{gp18}GTG) or without
612 (T_{gp18}AAA) the right PAM sequence. SIRV2M or SIRV2Mgp02 was added to the cultures (**a** and **b**)
613 at a MOI (multiplicity of infection) of 0.2. Control: 10 mM Tris acetate pH 6 was added. **c.**

614 Separation of Cas10d (green), AcrID1 (blue), and the Cas10d:AcrID1 complex (red) by gel
615 filtration using Superdex 200 Increase 10/300 GL. Separation of the proteins is shown by
616 absorption at 280 nm (solid lines) and 260 nm (dashed lines). Elution volumes for standard proteins
617 are indicated by arrows and masses in kDa at the top. **d.** SDS-PAGE with fractions of AcrID1,
618 Cas10d, and the Cas10d:AcrID1 complex as indicated. Standard molecular mass (kDa) markers are
619 indicated to the right. The growth curves in **a** and **b** are representatives of a single condition that
620 were confirmed by two independent experiments with different sampling time. Results in **c** and **d**
621 are representatives of three experiments.

622 **Fig. 5. Phylogeny of AcrID1 homologs.** The approximate maximum likelihood phylogenetic tree
623 was constructed as described under Methods. The support values estimated by FASTTREE
624 (percentage points) are at each internal branch. For each sequence, the GenBank identifier, the virus
625 name and the gene name (where available) are indicated. Pink, SIRV1 genes; blue, SIRV2 genes;
626 red, AcrID1s studied here; green, viruses that carry AcrID1-like genes but do not belong to the
627 family *Rudiviridae*.

628 **Fig. 6. Crystal structure of AcrID1.** **a.** Overview of the AcrID1 $\alpha\beta$ -domain fold with N- and C
629 termini as well as secondary structure elements labelled. **b.** The AcrID1 dimer with interactions at
630 the dimer interface shown with labelled side chain sticks. The side chain of Glu18 interacts with the
631 N-terminal NH₂-group of Met1. **c.** Conserved residues shown with colored sticks on the AcrID1
632 dimer in two perpendicular views. Residues are colored using the CLUSTAL X scheme:
633 Hydrophobic (A, I, L, M, F, W, V, C, light blue); aromatic (F, Y, W, teal); basic (K, R, red); acidic
634 (D, E, purple); polar (S, T, N, Q, green). **d.** Overlay of AcrID1 (blue) and two structurally similar
635 proteins, *S. islandicus* Filamentous Virus ORF14 (SIFV14, PDB ID 2H36^{40,41}, green) and *Acidianus*
636 Filamentous Virus 1 ORF-99 (AFV1-99, PDB IDs 3DJW and 3DF6⁴², red).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Anti-CRISPR proteins encoded by archaeal lytic viruses inhibit subtype I-D immunity

Fei He, Yuvaraj Bhoobalan-Chitty, Lan B Van, Anders L Kjeldsen, Matteo Dedola,
Kira S Makarova, Eugene V Koonin, Ditlev E Brodersen, Xu Peng

Correspondence to: peng@bio.ku.dk

This file includes:

Supplementary Tables 1 – 2

Supplementary Figures 1 – 9

Supplementary Table 1. Primer list

Construct	Primer ^a	DNA sequence (5'-3') ^b
Left terminus check primers 1	01S	CGGTCAAACAAGGTATTTTCGTAG
	10A	CTTGAATCTGGAACTCCAAC TAG
Left terminus check primers 2	VirFor	TGAAAGTCGAGAGATACAAATGT
	VirRev	ATCTACTAGAATCTTCATATGATCAC
	22For	AATTGATAGTTTTCGGGAATGG
pJET1.2/Flank2223	2223 <i>KpnI</i> <i>NheI</i> Rev	AATTCCTACGGGTACCGCTAGCCTTACACTTTTTATTTTCATCAGACAT
	2223 <i>NheI</i> <i>KpnI</i> For	AAAGTGTAAGGCTAGCGGTACCCGTAGGAATTCAGAAATGCGT
	23Rev	AGACTTGTTATAGATCCTTTAAAGTG
SIRV3frag gp02	gp02-V4 <i>NheI</i> For	CGGCTAGCGTGAAAAAATAAAGAGATAGAGAAGT
	gp02-V4 <i>KpnI</i> Rev	GGGGTACCTAGTTCATAGTTTTAGTTAGTAAGAAG
SIRV3frag gp02-03	gp02-V4 <i>NheI</i> For	CGGCTAGCGTGAAAAAATAAAGAGATAGAGAAGT
	gp03-V4 <i>KpnI</i> Rev	GGGGTACCAAATAGAATTATCATTCCAAACTTG
SIRV3frag gp02-03-04	gp02-V4 <i>NheI</i> For	CGGCTAGCGTGAAAAAATAAAGAGATAGAGAAGT
	gp04-V4 <i>KpnI</i> Rev	GGGGTACCTCTATCTTTTTACATCTCTATCTCTC
SIRV3frag gp02-03-04-05	gp02-V4 <i>NheI</i> For	CGGCTAGCGTGAAAAAATAAAGAGATAGAGAAGT
	gp05-V4 <i>KpnI</i> Rev	GGGGTACCTCTTGAATTAGTATCACAAACTTG
SIRV3frag gp03	gp03-V4 <i>NheI</i> For	CGGCTAGCTCTTCTTACTAACTAAACTATGAAGT
	gp03-V4 <i>KpnI</i> Rev	GGGGTACCAAATAGAATTATCATTCCAAACTTG
SIRV3frag gp03-04-05	gp03-V4 <i>NheI</i> For	CGGCTAGCTCTTCTTACTAACTAAACTATGAAGT
	gp05-V4 <i>KpnI</i> Rev	GGGGTACCTCTTGAATTAGTATCACAAACTTG
SIRV3frag gp04-05	gp04-V4 <i>NheI</i> For	CGGCTAGCGATTCAATACAATTAATTAGATATCCAAG
	gp05-V4 <i>KpnI</i> Rev	GGGGTACCTCTTGAATTAGTATCACAAACTTG
pEXA2-gp02	gp02-V4 <i>NdeI</i> For	CCCATATGAATTACAAAGAATTAGAAAAAATG
	gp02-V4 <i>NotI</i> Rev	TTGCGGCCGCTAGTTCATAGTTTTAGTTAGTAAGAAG
<i>C</i> _{gp44} GTT	gp44 GTTfwd	aacCTGAAATTTTATCAGTATCTGCTACTTCAACTCCTA
	gp44 GTTrev	tacTAGGAGTTGAAGTAGCAGATACTGATAAAATTCAG
<i>C</i> _{gp44} AAA	gp44 AAAfwd	aacTTTTATCAGTATCTGCTACTTCAACTCCTAGTAATG
	gp44 AAArev	tacCATTACTAGGAGTTGAAGTAGCAGATACTGATAAAA
<i>T</i> _{gp18} GTG	gp18 GTGfwd	aacGAAATTTGGAAAGTAGGAGAAAAGAACAATCTATAGATA
	gp18 GTGrev	tacTATCTATAGATTGTTCTTTTCTCTACTTTCCAAATTC
<i>T</i> _{gp18} AAA	gp18 AAAfwd	aacTTTGAAAGTAGGAGAAAAGAACAATCTATAGATATTCA
	gp18 AAArev	tacTGAATATCTATAGATTGTTCTTTTCTCTACTTTCCAAA
pMID-cas6(I-D)	cas6I-D Lfwd- <i>NcoI</i>	AATTCCATGGGAAAAGGAATGAGGATGCAAT
	cas6I-D Lrev- <i>XhoI</i>	ATCCTCGAGGGAGACACTGAGTTTTAATCTG
	cas6I-D Gfwd- <i>SalI</i>	AATTGTCGACGAGTACGAAAGAATTGAAAATGA
	cas6I-D Grev- <i>MluI</i>	CCACGCGTCAAAGAACAGAAATGGTTCTAAG
	cas6I-D Rfwd- <i>XhoI</i>	ATCCTCGAGCCTTAGAACCATTTCTGTTCTTT
	cas6I-D Rrev- <i>SphI</i>	AATTGCATGCTTATCCTTAGCTTTTAATTTACTTC
Δ cas6(I-A) and Δ cas6(I-D) check primers	cas6I-A con fwd	AAGGAGTGTGAAGAAGTAAGT
	cas6I-A con rev	GAAC TAAGAAAAGAGTCAACAAC
	cas6I-D con fwd	CGTTCAAGGAGTTAATAAAGTATT
	cas6I-D con rev	ATGCTTGCTTGCTTAATTCT
pcas6(I-D)	cas6I-D <i>NdeI</i> fwd	CGCCATATGCCTCTAATCAGATTAAACT
	cas6I-D <i>NotI</i> rev	ATTTGCGGCCGCTAAGGATTCTGTTTCGTCG
pET30a constructs	FPCas10d <i>NdeI</i>	TAGCGCATATGAGAAATTTGAAAAGAATAGTGA
	RPCas10d <i>XhoI</i>	TGACGCTCGAGTGATTATCACCTTGAAGGACT
	FPCas10ddel <i>NdeI</i>	TTTTCAATAGCCTATGCATTAAAAAGA
	RPCas10ddel <i>NdeI</i>	TCTTTAATGCATAGGCTATTGAAAA

FpCsc2 <i>NdeI</i>	<u>ATTACATATGAATGGGATTGAAGCA</u>
RpCsc2 <i>XhoI</i>	AATCCCGAGCTTTTAAATGAATTCCTCTATACT
FPgp02 <i>NdeI</i>	<u>CGCCATATGAATTACAAAGAATTAGAAAAAATG</u>
RPgp02 <i>XhoI</i>	CCGCTCGAGTAGTTCATAGTTTATAGTTAGTAAGAAG
FPgp02His <i>NdeI</i>	<u>CGCCATATGAATTACAAAGAATTAGAAAAAATG</u>
RPgp02His <i>XhoI</i>	CCGCTCGAGCTTTCTAATTTCTCCTCCAAAAATG

^a. Restriction enzymes used to digest the respective PCR product and the vector (pEXA2 or pJET1.2) before ligation were indicated in the primer names. ^b. Restriction sites are indicated in italics, sequences shared between primers in fusion PCR are underlined.

Supplementary Table 2. Crystallographic data collection and refinement statistics

	Se-Met	Native
Data collection		
Space group	P12 ₁ 1	P12 ₁ 1
Cell dimensions		
a, b, c (Å)	54.61, 79.98, 106.09	54.75, 80.44, 106.32
α, β, γ (°)	90.00, 101.08, 90.00	90.00, 101.23, 90.00
Wavelength (Å)	0.97947 (peak)	0.98020
Resolution (Å) ^a	32 - 2.03 (2.07-2.03)	27 - 1.93 (1.96 - 1.93)
No. of unique reflections	52.996 (1673)	67.149 (3320)
Completeness (%)	91.4 (57.9)	98.7 (97.8)
Anom. completeness (%)	90.8 (57.8)	-
Multiplicity	6.1 (4.0)	3.4 (3.2)
Anom. multiplicity	3.1 (2.0)	-
I/σ _I	15.5 (1.1)	13.2 (1.2)
Anomalous signal	1.486 (0.632)	-
R _{merge} (%)	6.6 (110.6)	3.6 (97.8)
R _{meas} (%)	7.1 (127.4)	4.3 (117.0)
CC _{1/2} (%)	99.8 (55.8)	99.9 (59.8)
Wilson B factor (Å ²)	57.131	48.087
Refinement		
Resolution (Å)	-	26.9 - 1.93
No. of reflections used	-	67,129
R _{work} /R _{free}		23.5 / 26.9
No. of atoms		
protein	-	5,235
solvent	-	57
Average B factor (Å ²)		
protein	-	73.3
solvent	-	63.7
rmsds from ideality		
bonds (Å)	-	0.016
angles (°)	-	1.4
Ramachandran plot		
favored	-	92.8
allowed	-	6.9
not allowed	-	0.4
Clash score	-	8.9

^aValues in parentheses correspond to the highest resolution shell.

Supplementary Figure 1. Recombination of the gp02-gp09 fragment into SIRV2M genome facilitating viral propagation in LAL14/1.

Electroporation was performed on LAL14/1 cells using either the gp02-gp09 containing PCR fragment (lane labeled SIRV2M + gp02-09) or water (lanes labeled control and SIRV2M). SIRV2M was immediately added to the cells (lanes labeled SIRV2M and SIRV2M + gp02-09) which were incubated for two days (A). The culture supernatant was diluted 1000 times into fresh LAL14/1 cells followed by two days of incubation (B). PCR was performed on the supernatant using primers VirFor and VirRev flanking the deleted 4 kb region (Supplementary Table 1). The result is a representative of a single experiment.

Supplementary Figure 2. PCR verification of gene deletion in Δ cas6(I-A) and Δ cas6(I-D). PCR was conducted with “ Δ cas6(I-A) and Δ cas6(I-D) check primers” (Table S1) using genomic DNAs extracted from *S. islandicus* LAL14/1 Δ *pyrF* strain LAL549 (WT), and respective deletion mutants (Δ). -, negative control. L, DNA size ladder. The result is a representative of two experiments.

Supplementary Figure 3. Further confirmation of the anti-CRISPR subtype I-D activity of SIRV3 gp02 (AcrID1). Growth of LAL14/1 Δ arrays cells carrying a plasmid-borne spacer matching the coding strand of gp44 of SIRV2 (and hence also of SIRV2M and SIRV2Mgp02) with (C_{gp44}GTT) or without (C_{gp44}AAA) the right PAM sequence. SIRV2M and SIRV2Mgp02 were added at a MOI of 1 and 10mM Tris.Ac pH 6 was added in the controls. The growth curves are representatives of a single condition that were confirmed by two experiments with different sampling time.

Supplementary Figure 4. Cas10d, but not Csc2, interacts with AcrID1. (A) Diagram showing the work flow. Heat treated cell lysates were made from *E. coli* cells expressing the bait (Cas10d-His or Csc2-His) and the prey (SIRV3 AcrID1) proteins. Equal amount of prey protein lysate was mixed with either of the bait protein lysates, purified with Ni-NTA agarose beads followed by SDS-PAGE. (B) Results from Coomassie blue staining (top) or Western blot using antibody raised against SIRV3 AcrID1 (bottom). Two eluents (1 and 2) from each purification were shown. Bands corresponding to Cas10d-His, Csc2-His or AcrID1 were indicated. Results in **b** is a representative of two experiments.

SIRV2 gp03 group

SIRV3 gp02 AcrID1 (sec. structure)

SIRV2 gp02 group

Supplementary Figure 5. Multiple alignment of SIRV2 gp03 homologs. Twelve SIRV2 gp03 homologs are highlighted by light blue on the left. The observed secondary structure of SIRV3 AcrID1 (gp02) (this study) is shown above the sequences. The alignment was colored using the CLUSTAL X scheme: Hydrophobic (A, I, L, M, F, W, V, C, light blue); aromatic (F, Y, W, teal); basic (K, R, red); acidic (D, E, purple); polar (S, T, N, Q, green). Sequences were aligned using the MUSCLE program.

Supplementary Figure6. AcrID1 and gp01 expansion in SIRV1, SIRV2 and SIRV3.

Colored arrows represent core genes conserved in *Sulfolobus* rudiviral genomes

(SIRV1 to SIRV11) (1). The viral gp numbers are shown above the arrows in pink boxes.

Reference

1. Bautista, M. A., Black, J. A., Youngblut, N. D. & Whitaker, R. J. Differentiation and Structure in *Sulfolobus islandicus* Rod-Shaped Virus Populations. *Viruses* 9, doi:10.3390/v9050120 (2017).

Supplementary Figure 7. Fiber-like structures observed in the AcrID1 crystal structure.

Supplementary Figure 8. Surface electrostatic potential for Acr proteins. Left, front (helix side) and back (sheet side) views of the SIRV3 gp02 AcrID structure (PDB ID 6EXP, this study) shown both as ribbons and in surface representation colored by electrostatic potential (-5.0 to $+5.0$ $K_B T/e_c$); Middle, views of the *S. islandicus* Filamentous Virus ORF14 (SIFV14, PDB ID 2H3624); Right, views of *Acidianus* Filamentous Virus 1 ORF-99 (AFV1-99, PDB ID 3DF6).

Supplementary Figure 9. Raw gels

Fig.1b

Fig.1b

Fig.1c

We are not able to find the full raw gel for Fig. 1c, thus provide an earlier gel in which the same PCR product was run (indicated by arrow)

SI Fig.1A

SI Fig.1A

Fig.2c

Fig.2c

SI Fig.1B

SI Fig.1B

Fig.2d

Fig.2d

Photographs used in Fig. 3b

Fig.4d

Fig.4d

SI Fig.1 gels are found together with Fig. 2 gels

SI Fig.2

SI Fig.4B

SI Fig.4B

SI Fig.4B

SI Fig.4B