AARHUS UNIVERSITY

Coversheet

This is the publisher's PDF (Version of Record) of the article.

This is the final published version of the article.

How to cite this publication:

Sim, T. G., Swindles, G. T., Morris, P. J., Gałka, M., Mullan, D., & Galloway, J. M. (2019). Pathways for ecological change in Canadian High Arctic wetlands under rapid twentieth century warming. *Geophysical Research Letters*, 46, 4726–4737. <https://doi.org/10.1029/2019GL082611>

Publication metadata

Title:	Pathways for ecological change in Canadian High Arctic wetlands under rapid twentieth century warming
Author(s):	Sim, T. G., Swindles, G. T., Morris, P. J., Gałka, M., Mullan, D., & Galloway, J. M.
Journal:	<i>Geophysical Research Letters</i>
DOI/Link:	https://doi.org/10.1029/2019GL082611
Document version:	Publisher's PDF (Version of Record)

General Rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

If the document is published under a Creative Commons license, this applies instead of the general rights.

Geophysical Research Letters

RESEARCH LETTER

10.1029/2019GL082611

Key Points:

- The ecological, hydrological, and C accumulation responses of Arctic wetlands to climate warming may be strongly influenced by wetland type
- Contrasting site-specific responses to an increase in growing degree days include increased moss diversity and a shift to shrub dominance
- Intensive grazing from Arctic geese may be an important driver for recent vegetation change in High Arctic coastal wetlands

Supporting Information:

- Supporting Information S1
- Data Set S1

Correspondence to:

T. G. Sim,
gy12tgs@leeds.ac.uk

Citation:

Sim, T. G., Swindles, G. T., Morris, P. J., Galka, M., Mullan, D., & Galloway, J. M. (2019). Pathways for ecological change in Canadian High Arctic wetlands under rapid twentieth century warming. *Geophysical Research Letters*, 46, 4726–4737. <https://doi.org/10.1029/2019GL082611>

Received 5 MAR 2019

Accepted 5 APR 2019

Accepted article online 11 APR 2019

Published online 15 MAY 2019

Pathways for Ecological Change in Canadian High Arctic Wetlands Under Rapid Twentieth Century Warming

T. G. Sim¹ , G. T. Swindles^{1,2} , P. J. Morris¹ , M. Galka³, D. Mullan⁴, and J. M. Galloway^{5,6}

¹School of Geography, University of Leeds, Leeds, UK, ²Ottawa-Carleton Geoscience Centre and Department of Earth Sciences, Carleton University, Ottawa, Ontario, Canada, ³Department of Geobotany and Plant Ecology, Faculty of Biology and Environmental Protection, University of Łódź, Łódź, Poland, ⁴School of Natural and Built Environment, Queen's University Belfast, Belfast, UK, ⁵Aarhus Institute of Advanced Studies (AIAS), Aarhus University, Aarhus, Denmark, ⁶Natural Resources Canada/Ressources naturelles Canada, Geological Survey of Canada/Commission géologique du Canada, Calgary, Alberta, Canada

Abstract We use paleoecological techniques to investigate how Canadian High Arctic wetlands responded to a mid-twentieth century increase in growing degree days. We observe an increase in wetness, moss diversity, and carbon accumulation in a polygon mire trough, likely related to ice wedge thaw. Contrastingly, the raised center of the polygon mire showed no clear response. Wet and dry indicator testate amoebae increased concomitantly in a valley fen, possibly relating to greater inundation from snowmelt followed by increasing evapotranspiration. This occurred alongside the appearance of generalist hummock mosses. A coastal fen underwent a shift from sedge to shrub dominance. The valley and coastal fens transitioned from minerogenic to organic-rich wetlands prior to the growing degree days increase. A subsequent shift to moss dominance in the coastal fen may relate to intensive grazing from Arctic geese. Our findings highlight the complex response of Arctic wetlands to warming and have implications for understanding their future carbon sink potential.

Plain Language Summary The response of Arctic wetland ecosystems and carbon stores to climate change is uncertain. We investigate the response of wetland ecosystems in the Canadian High Arctic to twentieth century climate warming. We use proxies for changes in vegetation (plant macrofossils) and wetness (testate amoebae) preserved in the wetland soil in combination with radiocarbon dating to reconstruct the past ecology of these wetlands. This approach allows us to explore beyond the timeframe of monitoring studies. Our results suggest that wetland type is an important determinant of the response of ecological, hydrological, and soil carbon accumulation to climate warming. Our findings highlight the clear but complex response of Arctic wetlands to twentieth century warming. This has important implications for understanding the future carbon sink potential of these ecosystems.

1. Introduction

Climate warming over the last century has been greatest in the Arctic and is projected to continue in the 21st century at a rate above the global average (Christensen et al., 2013). Arctic warming is causing widespread permafrost thaw (Jorgenson et al., 2006; Payette et al., 2004) and alteration of terrestrial ecosystem hydrology (Smith et al., 2007), vegetation (Myers-Smith et al., 2011), and wildfire regimes (Gibson et al., 2018; Myers-Smith et al., 2008). Changes to high-latitude ecosystems are likely to influence their carbon budgets, including global warming through enhanced carbon emissions (Dorrepaal et al., 2009). Permafrost soils in general could become carbon sources with warming through greater aerobic (CO₂) and anaerobic (CH₄) decomposition rates (Natali et al., 2015; Schuur et al., 2015). However, longer, warmer growing seasons and changes in Arctic precipitation regimes (Bintanja & Selten, 2014; Kattsov et al., 2007; Kopec et al., 2016) may stimulate carbon capture through enhanced plant productivity in peatlands and the transition of minerotrophic wetlands into organic peatlands (Charman et al., 2013, 2015; Gallego-Sala et al., 2018; Morris et al., 2018). Recent evidence shows an inconsistent response of Arctic and sub-Arctic peatlands to warming in terms of carbon accumulation (Zhang, Gallego-Sala, et al., 2018).

In the Arctic, wetlands occupy ~7% of the nonglaciated land area (Walker et al., 2005) and—although high-latitude carbon stocks are poorly constrained—wetlands store a disproportionate amount of carbon for their extent (Tarnocai et al., 2009). Arctic wetlands exemplify the complexity of landscapes underlain by

Figure 1. Study sites, nearby weather stations and the lake cores used for pollen-based temperature reconstruction in Gajewski (2015). Permafrost and the tree line data from Brown et al. (2002).

permafrost and typically occur in three locales: on ground affected by ice wedge formations (polygon mires), on previously glaciated terrain with favorable topographic depressions (known as patchy wetlands), and in coastal zones of isostatic uplift (coastal wetlands; Glenn & Woo, 1997; Woo & Young, 2006). Across Arctic wetlands, warming has been linked to greater plant biomass (Hill & Henry, 2011), vegetation composition changes and desiccation (Woo & Young, 2006, 2014; Zhang, Piilo, et al., 2018). Ice wedge polygon mires specifically are complex and dynamic systems (de Klerk et al., 2011; Fritz et al., 2016), and degradation in response to recent warming has led to changes in vegetation and drainage (Fraser et al., 2018; Jorgenson et al., 2015; Liljedahl et al., 2016; Perreault et al., 2017). Similarly, vegetation in some sub-Arctic and Arctic coastal wetlands has been altered by increasing bird grazing pressures in recent decades (Jefferies & Rockwell, 2002; Peterson et al., 2013).

Future hydrological changes, vegetation shifts and degradation of permafrost have been identified as key areas of uncertainty in the prediction of permafrost carbon dynamics (Abbott et al., 2016). There is limited information on the response of Arctic wetlands to climate change and their potential to transition into productive peatlands on centennial timescales. For the first time in the Canadian High Arctic, we use a high-resolution paleoecological approach (testate amoebae and plant macrofossils) to reconstruct past moisture conditions and vegetation histories in peat-forming wetlands. We aim to investigate the ecological response to twentieth century climate warming in the three main types of High Arctic wetland: (i) a polygon mire, (ii) a valley fen (a type of patchy wetland), and (iii) a coastal fen.

2. Study Sites, Materials, and Method

The study region is in the continuous permafrost zone of the western Canadian Arctic tundra, between the latitudes 68.5 and 74.5 °N (Figure 1). We study two wetlands on northern Banks Island, Northwest Territories: a polygon mire (74.459 °N, 121.04 °E) and a valley fen (74.05 °N, 118.429 °E). The distinctive topography of the polygon mire was captured by extracting a monolith from a raised center mound and from a surrounding trough. Our third site is a coastal fen (68.65 °N, 105.455 °E) located on an island ~50-km south of Cambridge Bay, Nunavut, where two monoliths <10 m apart were collected as replicate paleorecords.

Figure 2. Recent warming in western Canadian Arctic. (a) Annual temperature data from nearby weather stations. (b) GDD₀ modelled for each site from climate reanalysis data (Compo et al., 2011). Stepped lines indicate changepoints. (c) Reconstructed mean July temperature for western Arctic during the Holocene (Gajewski, 2015). Shading indicates one standard deviation.

Monoliths were sampled to the base of the active layer and subsampled in the laboratory at 1-cm depth intervals. Bulk density and organic matter content were calculated in accordance with Chambers et al. (2011). Testate amoebae were prepared for analysis using the method outlined by Booth et al. (2010). In the absence of an appropriate transfer function, testate amoebae were grouped into categories of hydrological preference based on a detailed literature review (Table S2 in the supporting information). Plant macrofossils were prepared in accordance with Gałka, Tobolski, et al. (2017). Aboveground plant macrofossils and bulk peat (where reliable plant macrofossils were not present) were radiocarbon dated (Table S1). Radiocarbon dates were calibrated using Clam 2.2 (Blaauw, 2010) in R 3.4.3 (R Core Team, 2018). The IntCal13 (Reimer et al., 2013) calibration curve for pre-bomb dates and Hua et al. (2013) for post-bomb dates were used. When post-bomb dates demonstrated multiple probability distributions during calibration, we applied the principle of superposition to determine the most likely probability distribution. Chronological uncertainty can be introduced by cryoturbation in permafrost soils, particularly toward to base of the active layer (Bockheim & Tarnocai, 1998; Ping et al., 2008). These factors are considered in our interpretation and are comprehensively discussed in the supplementary materials (Table S1 and Text S3). The most appropriate age-depth model for each chronology was selected including linear interpolation, cubic spline, and polynomial regression. Growing degree days above 0 °C (GDD₀) were calculated for each site from monthly climate reanalysis data spanning the period AD 1851–2011 (Compo et al., 2011). Changepoint analysis (Killick & Eckley, 2014) was conducted in R 3.4.3 (R Core Team, 2018) on GDD₀ time series data for each site. See full methods and study site information in the supporting information (Figures S1 to S6, Tables S1 to S2, and Text S1 to S3).

3. Results

3.1. Twentieth Century Warming and Lengthening Growing Seasons in the Western Arctic

In the western Arctic following a warm period from ~10 to 7.5 Ka BP, there was general cooling throughout the Holocene (Gajewski, 2015; Figure 2c). Weather station records beginning ~AD 1950 document twentieth

century climatic warming near to our sites (Figure 1) and across the Arctic (McBean et al., 2005). The Cambridge Bay and Sachs Harbour stations show pronounced warming rates of 0.3 and 0.5 °C per decade, respectively, contrasted to 0.1 °C per decade at Mould Bay, the highest latitude station (Figure 2a). Reanalysis data (Compo et al., 2011) demonstrated a pronounced increase in GDD₀ across all wetland sites with changepoints detected between 1931 and 1940 (Figure 2b). Following respective changepoints, mean GDD₀ at the polygon mire increased from 49 ± 50 to 152 ± 46 °C days year⁻¹ (changepoint: 1940) and at the valley fen, from 105 ± 76 to 260 ± 6 °C days year⁻¹ (changepoint: 1936; error terms indicate standard deviation). After the 1931 changepoint, mean GDD₀ at the coastal fen increased from 400 ± 132 to 605 ± 68.3 °C days year⁻¹.

The rise in GDD₀ in the 1930s (Figure 2b) is consistent with a well-documented climatic fluctuation (~1920–1940) around the North Atlantic. This event began in the 1920s and resulted in positive Arctic-wide average temperature anomalies for most of the 1930s (Johannessen et al., 2004; Polyakov et al., 2003). Causes of the climatic fluctuation remain open to debate, with internal atmospheric variability, anthropogenic greenhouse gas forcing, solar variability, volcanic forcing, and regional dynamic feedbacks proposed in the literature (Wood & Overland, 2010). Bengtsson et al. (2004) suggested natural variability was the likely cause, as reduced sea ice cover was critical in initiating warming of ~1.7 °C during the peak period between 1930 and 1940 at 60–90 °N. However, Wood and Overland (2010) interpreted the event as an intrinsically forced albeit essentially random climatic phenomenon, superimposed on rising temperatures associated with anthropogenic forcing. GDD₀ have remained high following this event (Figure 2b) and increased elsewhere in the Canadian High Arctic in the past few decades (Woo & Young, 2014).

3.2. Paleoecological Results

3.2.1. Polygon Mire

The polygon mire demonstrates contrasting responses to recent warming between the raised center and the trough. The raised center monolith (11 cm; Figure 3a) is mainly composed of partially decomposed *Scorpidium cossonii* (80%–90.5%), bulk density is relatively high (0.36 ± 0.1 g cm⁻³), while organic matter is relatively low ($29.8\% \pm 9.4\%$), and carbon accumulation increases toward the top of the profile. The trough monolith (17 cm; Figure 3b) is more dynamic by comparison, recording multiple vegetation shifts. The lower phase (17–7 cm) consists of a light brown, peaty soil undergoing a gradual transition (~AD 750–1800) from *S. cossonii* (40%–85%) and *Warnstorfia sarmentosa* (0%–35%) moss to a more sedge (Cyperaceae) and herb-dominated wetland. During this phase, carbon accumulation is low (6.2 ± 2.1 g m⁻² year⁻¹), organic matter is low ($27.8\% \pm 8.7\%$), and there is relatively high bulk density (0.51 ± 0.16 g cm⁻³). In the upper phase (0–7 cm), organic matter increases (max: 77.4%) and bulk density decreases (min: 0.04 g cm⁻³). A sedge and herb-dominated ecosystem is present from ~AD 1800 before an increase in moss diversity at ~AD 2000, including the presence of *S. cossonii* (20%–70%), *Calliergon* spp. (10%–30%), and *Campylium* cf. *stellatum* (10%–20%). Coincident with this ~AD 2000 is a shift in hydrological conditions, with wet-indicator testate amoeba taxa increasing dramatically ($68.8\% \pm 17.9\%$) and carbon accumulation increasing to between 45.7 and 102.8 g m⁻² year⁻¹.

3.2.2. Valley Fen

The valley fen monolith (13 cm; Figure 3c) is composed of three main phases. Notwithstanding chronological uncertainty, our age-depth model suggests that the base of the record (13–9 cm) accumulated before ~AD 1780. The base phase is characterized by a pale-colored mineral layer of low organic matter content (7.2%–11.6%), high bulk density (1.08 – 1.52 g cm⁻³), and sedge-dominated vegetation (>75%). At ~7 cm (~AD 1800), there is a sharp increase in organic matter (11.6%–33.4%) and a decrease in bulk density (1.08 – 0.38 g cm⁻³). Throughout the middle section of the profile (9–4 cm), there is a gradual increase in organic matter to 56.5% and decrease in bulk density to 0.23 g cm⁻³ (~AD 1950). This change represents the development of a more productive intermediate or rich fen system dominated by brown mosses (60%–75%), *Calliergon* spp. (<20%), and *Hamatocaulis vernicosus* (<10%), with increasing carbon accumulation (10.9 – 20.1 g m⁻² year⁻¹). In the top 4 cm (~AD 1950 onward), the wetland demonstrates properties more characteristic of a peatland. Organic matter continues to increase to 78.8% and bulk density decreases further to 0.06 g cm⁻³. *Tomentypnum nitens* (30%–50%) and *Cinclidium stygium* (0%–20%) now dominate the valley fen record alongside other mosses. These mosses (including *T. nitens* and *C. stygium*) may have been present under the Brown moss spp. classification; however, decomposition made species identification impossible.

Figure 3. Summary stratigraphic diagram of paleoecological variables from polygon mire raised center (a), polygon mire trough (b), valley fen (c), and coastal fen (d and e; yellow arrows indicate sampling locations). GDD₀ CP is changepoint in GDD₀. For detailed stratigraphic diagrams see Figures S7–S21. GDD₀ = growing degree days.

3.2.3. Coastal Fen

The peat monoliths analyzed from coastal fen A (CFA) and coastal fen B (CFB) demonstrate highly similar responses, each recording two distinct phases (Figures 3d and 3e). Notwithstanding chronological uncertainty, our age-depth model suggests the lower phase (CFA 17–12 cm and CFB 16–10 cm) accumulated before ~AD 1950. This lower phase is characterized by a pale-colored minerogenic layer dominated by sedges and herbs, with low organic matter (mostly 2%–26%) and a high bulk density (max: 2.44 g cm⁻³), which decreases up the succession (min: 0.38 g cm⁻³). The upper phase (CFA 12–0 cm and CFB 10–0 cm) is characterized by a darker-colored peaty layer of poorly decomposed plant material and a sharp rise in organic matter that remains consistently high (average of upper phases in both monoliths = 87% ± 4.8%) alongside a relatively low bulk density (average of upper phases in both monoliths = 0.12 ± 0.02 g cm⁻³). The vegetation in the upper phase initially changes from sedge and herb dominance, to a transitional phase dominated

by the moss *Calliergon richardsonii* (CFA 9–8 cm and CFB 9–8 cm). Shrubs then dominate (CFA 8–2 cm and CFB 8–1 cm), before a return to a dominance of *C. richardsonii* (>50%) alongside the appearance of *Brachythecium mildeanum* (in CFA; <40%). At the coastal fen site, the ecosystem shift to shrub dominance is coincident with fast carbon accumulation rates (~AD 1980) of $>100 \text{ g m}^{-2} \text{ year}^{-1}$.

4. Discussion

Under 21st century warming and in the presence of adequate moisture, Arctic wetlands such as our study sites may become more productive and transition into peatlands—this could to some extent mitigate carbon losses from degrading peatlands farther south (Charman et al., 2015; Gallego-Sala et al., 2018). The Arctic wetlands we studied did demonstrate ecological responses to a mid-twentieth century increase in GDD₀. However, the pathway of these responses varied between wetland types, and therefore, their future carbon balance is likely to be complex and may also vary between wetland types. In particular, our results highlight the importance of ground ice dynamics, changing seasonal hydrological regimes, and herbivory.

4.1. Ground Ice Dynamics and Permafrost Feedbacks

Ground ice dynamics and feedbacks in permafrost regions are a key uncertainty in understanding the carbon cycling response of these systems to warming (Grosse et al., 2016; Schuur et al., 2015). We believe our polygon mire site exemplifies the influence ground ice dynamics can have upon wetland structure and ecology with warming. A changepoint increase in GDD₀ was detected at the polygon mire in 1940 (Figure 2b). A further increase in GDD₀ from 1956 to 2016 at Sach Harbour has been documented and linked to increasing active layer depth (Fraser et al., 2018). At the polygon mire, the raised center monolith showed no response to recent warming, perhaps because permafrost thaw has not yet induced the collapse of the raised center structure, as may be the case in some Siberian peatlands (c.f. Teltewskoi et al., 2016). In contrast, the trough demonstrates high sedge abundance from ~AD 1800, consistent with stable or initially degrading ice wedges, while the shift to dominance of hydrophilic moss (*S. cossonii*) from ~AD 2000 is characteristic of advanced ice wedge degradation (Figure 3.6b; Jorgenson et al., 2015). Recent ice wedge degradation is further supported by an increase in wet indicator testate amoeba from ~AD 2000 (Figure 3.5b). This decadal or subdecadal time-scale of trough wetting is supported by observations from across the Arctic since the mid-twentieth century (Fraser et al., 2018; Jorgenson et al., 2015; Liljedahl et al., 2016). The previous periods of sedge and herb dominance and moss dominance in the record (Figure 3.6b) may relate to previous ice wedge degradation and stabilization cycles. However, accurate interpretation of these previous trough vegetation shifts is limited by the absence of a testate amoeba hydrological record (barren below 7 cm). Local geomorphological and autogenic vegetation processes are key in the formation and variability of polygon mires, but climate change may modify these patterns (Ellis & Rochefort, 2004). We suggest the magnitude of recent warming (Figure 2) alongside background autogenic processes caused a threshold of ice wedge thaw to be reached ~AD 2000 in the polygon mire.

Under advanced ice wedge degradation and the establishment of hydrological connectivity between troughs, landscape-scale drainage can be initiated (Liljedahl et al., 2016). Drainage could threaten wetland persistence and expose partially decomposed carbon to increased aerobic decomposition—elevating CO₂ emissions (Schädel et al., 2016). However, the establishment of moss vegetation ~AD 2000, under wetter conditions (Figure 3b), may represent a negative ecological feedback to ice wedge thaw in troughs. The accumulated layer of moss since ~AD 2000 may act as an insulating layer alongside slumped material from trough banks to retard further thaw (Jorgenson et al., 2006, 2015). The slumping process was observed onsite (Figure S2) and may explain an older bulk peat date (1–2 cm; AD 1956 ± 1) stratigraphically above a reliable plant macrofossil date (4–5 cm; AD 1998 ± 2; Table S1). The negative ecological and geomorphological (slumping) feedbacks could contribute to the deceleration of ice wedge degradation. Our results highlight the importance of ground ice dynamics and autogenic factors in polygon mires with warming (c.f. Fritz et al., 2016; Vardy et al., 2005).

4.2. Changing Hydrological Regime and Precipitation With Warming

The short growing season of around 2 months in the High Arctic creates a highly seasonal hydrological regime, which our results suggest is being influenced by warming. When temperatures rise above 0 °C there is an influx of water to catchments from the thaw of ground ice and snowbanks that have accumulated over

the winter (Glenn & Woo, 1997)—this is particularly relevant at our valley fen site. An increase in GDD_0 at the valley fen from AD 1936 (Figure 2b) is followed at ~AD 1950 by a marked increase in wet indicator testate amoeba taxa and slight increase in dry indicator taxa (Figure 3.5c). The apparent increase in wetness is likely as a result of increased catchment thaw under warming conditions—a process that has been observed in other Arctic and alpine regions (Fontana et al., 2010; Woo & Young, 2014). The increase in dry indicator testate amoeba taxa may represent increasingly dry conditions toward the end of the growing season as a result of increased evapotranspiration over a longer and warmer summer (Oechel et al., 1998; Woo & Young, 2014; Zhang, Piilo, et al., 2018). This increased variability in hydrological conditions from ~AD 1950 is further supported by the presence of the hummock mosses *T. nitens* (30%–50%) and *Aulacomnium palustre* (<5%). These mosses have broad climatic and ecological ranges (Minke et al., 2009; Nicholson & Gignac, 1995) making them suitable for the increasingly variable moisture conditions. *T. nitens* specifically has been observed in both wet High Arctic (Steere & Scotter, 1979) and dry Boreal fens (Gignac et al., 1991), suggesting a certain resilience to warming temperatures alongside more variable hydrology.

Under the likely scenario of further Arctic warming (Christensen et al., 2013), alterations to the hydrology of the valley fen may threaten future persistence. Greater summer snowmelt may increase inundation in the short term. However, there may come a point where snowbanks become sufficiently depleted by increased evapotranspiration and no longer have the capacity to sustain wetlands (Woo & Young, 2006). Possible increases in 21st century Arctic late autumn and winter precipitation (Bintanja & Selten, 2014; Kattsov et al., 2007) may counteract increased evaporative losses to some extent. However, large uncertainties remain in predicting future precipitation patterns. Since ~AD 1950, there has been no clear shift in precipitation at the closest weather stations to our sites (Figure S1). Furthermore, increased future permafrost thaw and active layer thickening could increase permeability and thus drainage (Avis et al., 2011). Under the scenario of drying, CO_2 emissions from aerobic decomposition are likely to increase and the wetland may become a carbon source (Oechel et al., 1998). Carbon accumulation rates since ~AD 1950 remain consistent around $20 \text{ g m}^{-2} \text{ year}^{-1}$ before a recent increase to $61.3 \text{ g m}^{-2} \text{ year}^{-1}$ (Figure 3.4c). This increase hints at improved productivity with increased GDD_0 , indicating that the wetland ecosystem has not reached a moisture-limited point where carbon loss to decomposition outweighs input from litter. However, this carbon accumulation rate should be treated with caution because incomplete decomposition of recently accumulated organic matter was not accounted for. Our results demonstrate a clear ecosystem shift in response to recent warming, linked to a changing hydrological regime. The balance of possible increases in 21st century precipitation against summer evapotranspiration losses—alongside potential changes to drainage with permafrost thaw—is likely to determine the future sustainability and carbon accumulation capacity of this valley fen and potentially similar Arctic wetlands.

4.3. Potential Impact of Herbivory on Arctic Coastal Wetlands

The timing and magnitude of the ecosystem shift from sedges to shrubs is well established in both coastal fen records by ~AD 1950 and is likely linked to warming. The subsequent shift to moss dominance in both records since ~AD 2000 is likely linked to changes in Arctic geese grazing pressures. The increased dominance of shrubs in this coastal wetland ~AD 1950 follows an increase in GDD_0 at AD 1931 (Figure 2b). Northward expansion of shrub communities and increased productivity in response to recent warming has been widely documented across the tundra (Elmendorf et al., 2012; Myers-Smith et al., 2011; Tape et al., 2006) and directly linked to increased shrub and decreased sedge abundance in an Alaskan coastal wetland (Carlson et al., 2018). Isostatic uplift can lead to a drop in relative sea level and lowering of the water table, increasing dryness to facilitate sedge to shrub succession (Klinger & Short, 1996). However, any hydrological shift associated with an increased abundance of shrubs is unclear in our testate amoeba data. Furthermore, the projected range of sea-level change for nearby Cambridge Bay by 2100 relative to 2010 is between a 15 cm fall and 30 cm rise, depending upon future emissions scenarios (James et al., 2011). Therefore, isostatic uplift is likely to be more relevant in regions where uplift outpaces global eustasy, such as Hudson Bay and the Northwest Passage (James et al., 2015). We suggest the increase in GDD_0 is the most plausible mechanism driving this shift to shrub dominance at our site. The role of climate is further supported by evidence of a recent increase in shrub taxa in sub-Arctic permafrost peatlands in Sweden (Gałka, Szal, et al., 2017) and Alaska (Gałka et al., 2018)—areas of limited isostatic uplift (Geruo et al., 2013; Peltier, 2004).

The contemporary wetland ecosystem demonstrates minimal shrub cover and contains large sections of bare substrate and patches of dead moss (Figure S4). Both coastal fen monoliths demonstrate a recent shift to mosses ~AD 2000 (Figures 3d and 3e). Wildfire has reduced shrub cover in similar environments (Higuera et al., 2008; Mack et al., 2011), but the absence of macrocharcoal at our site rules this out. An alternative possibility for the reappearance of moss is increased grazing by Arctic geese, for example, lesser snow geese (*Chen caerulescens*) and Ross's geese (*Anser rossii*). Snow geese have demonstrated a preference for herbivory of vascular plants and therefore across the coastal wetlands of northern Canada, areas that have been subjected to destructive grazing often experience an increase in moss vegetation abundance (Abraham & Jefferies, 1997; Alisauskas et al., 2006). In areas of more intense grazing, replacement of moss by bare peat has been documented (Conkin & Alisauskas, 2017). Similarly, the recolonization of mosses toward the edge of the wetland (Figure S4) is typical of disturbed wetlands (Speed et al., 2010).

Arctic geese populations are being driven up by improved food availability from agricultural production (via artificial fertilizers and subsidy policies) in the wintering grounds of the southern United States (Fox et al., 2005; Jefferies et al., 2004). Our coastal fen site is located ~20-km northwest of Queen Maud Gulf Migratory Bird Sanctuary, a major summer nesting site for Arctic geese (Cooch et al., 2001) and bird presence can be confirmed unambiguously at our coastal fen site by an abundance of tracks and excrement (Figure S4). The Queen Maud Gulf Migratory Bird Sanctuary has seen Arctic geese nesting populations rise from ~44,300 in 1965 to ~2,251,900 in 2006, with over 50% of this increase occurring since 1998 (Kerbes et al., 2014). The likely recent intensification of grazing in established summer nesting sites may have caused snow geese to seek out more pristine habitats further north as they warm, such as our site, and begin to degrade them. Therefore, we suggest either edge recolonization following Arctic geese disturbance or selective Arctic geese herbivory provides a plausible explanation for the dramatic shift from a shrub to moss-dominated ecosystem ~AD 2000. It remains unclear what implications this shift from shrub to moss dominance will have upon long-term carbon accumulation rates; however, our findings highlight the potential importance of Arctic geese grazing (substantially influenced by nonclimate factors) in the context of future Arctic carbon cycling.

5. Conclusions

Warming of the western Canadian Arctic has caused a mid-twentieth century increase in growing degree days above 0 °C (GDD₀). In the absence of long-term monitoring data our high-resolution multiproxy paleoecological approach—the first of its kind in the Canadian High Arctic—allowed us to identify alternate pathways for ecological responses to recent warming in three wetlands. The main findings of our study are as follows:

1. An increase in moss diversity, decrease in sedges and herbs, and an increase in carbon accumulation rate occurred in the trough of a high centered polygon mire following the shift in GDD₀. Testate amoeba data suggest increased wetness at this time which may be related to ice wedge thaw driven by recent climate warming. In comparison, the raised center of the polygon mire showed no clear response to recent warming.
2. In a valley fen, the appearance of generalist mosses coincides with the shift in GDD₀. This occurs alongside an increase in both dry and wet indicator testate amoebae that may suggest increased seasonality in hydrological conditions (wetter conditions during snowmelt and drier conditions in the late summer owing to increased evapotranspiration). This site shifted from a minerogenic to an organic-rich wetland sometime prior to the shift in GDD₀.
3. A coastal fen site experienced an ecosystem shift coincident with an increase in GDD₀ from a sedge-dominated fen to a shrub-dominated fen with a high carbon accumulation rate. This site shifted from a minerogenic wetland to an organic-rich peatland sometime before the shift in GDD₀. The results from this site have important implications for understanding the expansion of shrub communities in the Arctic.
4. A subsequent reduction in shrub cover in the coastal fen since ~AD 2000 and recolonization by mosses may be related to an increase in destructive grazing by Arctic geese. This may affect long-term carbon accumulation rates.
5. Our results show clear, albeit complex ecological responses to warming in these Arctic wetlands. These complexities have important implications for understanding of the future carbon sink potential of these ecosystems under climate warming and are important areas for future research.

Acknowledgments

T. G. S. is funded by the Leeds-York Natural Environment Research Council Doctoral Training Partnership (NE/L002574/1). We acknowledge the generous support of Natural Resources Canada/Geological Survey of Canada in funding our radiocarbon dating and thank the André E. Lalonde AMS Laboratory, University of Ottawa, for the rapid analysis. This paper represents NRCan Contribution number/Numéro de contribution de RNCAN: 20180298. For the Banks Island sample collection, we thank Rod Smith of Geological Survey Canada, René Gysler of Great Slave Helicopters and Andrew Durbano. Sample collection occurred during a Geological Survey of Canada Geo-Mapping for Energy and Minerals Program (GEM Western Arctic) field program. For the Elu Inlet sample collection, we thank Thomas Hadlari, Attima Hadlari, Elisabeth Jansen-Hadlari, Francis Emingak, Braden Gregory, and Nawaf Nasser. Funding for field logistics was provided by a Polar Knowledge Canada/Savoir polaire Canada grant number 1516-149 to Jennifer Galloway and Timothy Patterson and from the Geological Survey of Canada Environmental Geoscience Program. For the helpful discussion and comments regarding the identification and classification of the ciliate Vaginicolidae (included alongside testate amoebae as a paleohydrological indicator), we thank Matthew Amesbury, Anatoly Bobrov, Louis Beyens, Dan Charman, Clément Duckert, Diego Fontaneto, Anna Kosakyan, Mariusz Lamentowicz, Michelle McKeown, Edward Mitchell, Thomas Roland, Ferry Siemensma, and Dave Wilkinson. We thank Lars Hedenäs for help in the identification of brown moss species in the plant macrofossil analysis. We thank the contributors to the Climate Explorer site for the re-analysis and station data used in this paper and are grateful to Environment Canada for the extended temperature records for the Sachs Harbour station. We acknowledge receiving isostatic uplift data from <http://grace.jpl.nasa.gov> website. We also thank Konrad Gajewski for his advice on interpretation of past temperature reconstructions and for directing us to pollen temperature reconstruction datafiles. All paleoecological data is available in the supporting information.

References

- Abbott, B. W., Jones, J. B., Schuur, E. A. G., Chapin III, F. S., Bowden, W. B., Bret-Harte, M. S., et al. (2016). Biomass offsets little or none of permafrost carbon release from soils, streams, and wildfire: An expert assessment. *Environmental Research Letters*, 11(3), 034014. <https://doi.org/10.1088/1748-9326/11/3/034014>
- Abraham, K. F., & Jefferies, R. L. (1997). High goose populations: Causes, impacts and implications. In B. D. J. Batt (Ed.), *Arctic ecosystems in peril: Report of the Arctic Goose Habitat Working Group. Arctic Goose Joint Venture Special Publication* (pp. 7–72). Ottawa, Ontario: Canadian Wildlife Service.
- Alisauskas, R. T., Charlwood, J. W., & Kellett, D. K. (2006). Vegetation correlates of the history and density of nesting by Ross's geese and lesser snow geese at Karrak Lake, Nunavut, Arctic. Arctic Institute of North America. <https://doi.org/10.2307/40512794>
- Amesbury, M. J., Roland, T. P., Royles, J., Hodgson, D. A., Convey, P., Griffiths, H., & Charman, D. J. (2017). Widespread biological response to rapid warming on the Antarctic Peninsula. *Current Biology*, 27(11), 1616–1622. <https://doi.org/10.1016/j.cub.2017.04.034>
- Avis, C. A., Weaver, A. J., & Meissner, K. J. (2011). Reduction in areal extent of high-latitude wetlands in response to permafrost thaw. *Nature Geoscience*, 4(7), 444–448. <https://doi.org/10.1038/ngeo1160>
- Bengtsson, L., Semenov, V. A., & Johannessen, O. M. (2004). The early twentieth-century warming in the Arctic—A possible mechanism. *Journal of Climate*, 17(20), 4045–4057. [https://doi.org/10.1175/1520-0442\(2004\)017<4045:TETWIT>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<4045:TETWIT>2.0.CO;2)
- Bintanja, R., & Selten, F. M. (2014). Future increases in Arctic precipitation linked to local evaporation and sea-ice retreat. *Nature*, 509(7501), 479–482. <https://doi.org/10.1038/nature13259>
- Blaauw, M. (2010). Methods and code for 'classical' age-modelling of radiocarbon sequences. *Quaternary Geochronology*, 5(5), 512–518. <https://doi.org/10.1016/J.QUAGEO.2010.01.002>
- Bockheim, J., & Tarnocai, C. (1998). Recognition of cryoturbation for classifying permafrost-affected soils. *Geoderma*, 81(3–4), 281–293. [https://doi.org/10.1016/S0016-7061\(97\)00115-8](https://doi.org/10.1016/S0016-7061(97)00115-8)
- Booth, R. K., Lamentowicz, M., & Charman, D. J. (2010). Preparation and analysis of testate amoebae in peatland paleoenvironmental studies. *Mires and Peat*, 7(2), 1–7.
- Brown, J., Ferrians, O., Heginbottom, J. A., & Melnikov, E. (2002). Circum-Arctic map of permafrost and ground-ice conditions, Version 2. Boulder, Colorado USA: National Snow and Ice Data Center.
- Carlson, L. G., Beard, K. H., & Adler, P. B. (2018). Direct effects of warming increase woody plant abundance in a subarctic wetland. *Ecology and Evolution*, 8(5), 2868–2879. <https://doi.org/10.1002/ece3.3902>
- Chambers, F. M., Beilman, D. W., & Yu, Z. (2011). Methods for determining peat humification and for quantifying peat bulk density, organic matter and carbon content for paleostudies of climate and peatland carbon dynamics. *Mires and Peat*, 7(07), 1–10.
- Charman, D. J., Amesbury, M. J., Hinchliffe, W., Hughes, P. D. M., Mallon, G., Blake, W. H., et al. (2015). Drivers of Holocene peatland carbon accumulation across a climate gradient in northeastern North America. *Quaternary Science Reviews*, 121, 110–119. <https://doi.org/10.1016/J.QUASCIREV.2015.05.012>
- Charman, D. J., Beilman, D. W., Blaauw, M., Booth, R. K., Brewer, S., Chambers, F. M., et al. (2013). Climate-related changes in peatland carbon accumulation during the last millennium. *Biogeosciences*, 10(2), 929–944. <https://doi.org/10.5194/bg-10-929-2013>
- Christensen, J. H., Kumar, K. K., Aldrian, E., An, S.-I., Cavalcanti, I. F. A., de Castro, M., et al. (2013). Climate Phenomena and their relevance for future regional climate change. In T. F. Stocker et al. (Eds.), *The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 1217–1310). Cambridge, UK, and New York, NY, USA: Cambridge University Press.
- Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Matsui, N., Allan, R. J., Yin, X., et al. (2011). The twentieth century reanalysis project. *Quarterly Journal of the Royal Meteorological Society*, 137(654), 1–28. <https://doi.org/10.1002/qj.776>
- Conkin, J., & Alisauskas, R. T. (2017). Conversion of tundra to exposed peat habitat by snow geese (*Chen caerulescens caerulescens*) and Ross's geese (*C. rossii*) in the central Canadian Arctic. *Polar Biology*, 40(3), 563–576. <https://doi.org/10.1007/s00300-016-1979-x>
- Cooch, E., Rockwell, R. F., & Brault, S. (2001). Retrospective analysis of demographic responses to environmental change: A lesser snow goose example. *Ecological Monographs*, 71(3), 377–400. [https://doi.org/10.1890/0012-9615\(2001\)071\[0377:RAODRT\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2001)071[0377:RAODRT]2.0.CO;2)
- de Klerk, P., Donner, N., Karpov, N. S., Minke, M., & Joosten, H. (2011). Short-term dynamics of a low-centred ice-wedge polygon near Chokurdakh (NE Yakutia, NE Siberia) and climate change during the last ca 1250 years. *Quaternary Science Reviews*, 30(21–22), 3013–3031. <https://doi.org/10.1016/J.QUASCIREV.2011.06.016>
- Dorrepaa, E., Toet, S., van Logtestijn, R. S. P., Swart, E., van de Weg, M. J., Callaghan, T. V., & Aerts, R. (2009). Carbon respiration from subsurface peat accelerated by climate warming in the subarctic. *Nature*, 460(7255), 616–619. <https://doi.org/10.1038/nature08216>
- Ellis, C. J., & Rochefort, L. (2004). Century-scale development of polygon-patterned tundra wetland, Bylot Island (73° N, 80° W). *Ecology*, 85(4), 963–978. <https://doi.org/10.1890/02-0614>
- Elmendorf, S. C., Henry, G. H. R., Hollister, R. D., Björk, R. G., Boulanger-Lapointe, N., Cooper, E. J., et al. (2012). Plot-scale evidence of tundra vegetation change and links to recent summer warming. *Nature Climate Change*, 2(6), 453–457. <https://doi.org/10.1038/nclimate1465>
- Fontana, F. M. A., Trishchenko, A. P., Luo, Y., Khlopenkov, K. V., Nussbaumer, S. U., & Wunderle, S. (2010). Perennial snow and ice variations (2000–2008) in the Arctic circumpolar land area from satellite observations. *Journal of Geophysical Research*, 115, F04020. <https://doi.org/10.1029/2010JF001664>
- Fox, A. D., Madsen, J., Boyd, H., Kuijken, E., Norriss, D. W., Tombre, I. M., & Stroud, D. A. (2005). Effects of agricultural change on abundance, fitness components and distribution of two arctic-nesting goose populations. *Global Change Biology*, 11(6), 881–893. <https://doi.org/10.1111/j.1365-2486.2005.00941.x>
- Fraser, R., Kokelj, S., Lantz, T., McFarlane-Winchester, M., Olthof, I., & Lacelle, D. (2018). Climate sensitivity of high arctic permafrost terrain demonstrated by widespread ice-wedge thermokarst on Banks Island. *Remote Sensing*, 10(6), 954. <https://doi.org/10.3390/rs10060954>
- Fritz, M., Wolter, J., Rudaya, N., Palagushkina, O., Nazarova, L., Obu, J., et al. (2016). Holocene ice-wedge polygon development in northern Yukon permafrost peatlands (Canada). *Quaternary Science Reviews*, 147, 279–297. <https://doi.org/10.1016/J.QUASCIREV.2016.02.008>
- Gajewski, K. (2015). Quantitative reconstruction of Holocene temperatures across the Canadian Arctic and Greenland. *Global and Planetary Change*, 128, 14–23. <https://doi.org/10.1016/J.GLOPLACHA.2015.02.003>
- Galka, M., Swindles, G. T., Szal, M., Fulweber, R., & Feurdean, A. (2018). Response of plant communities to climate change during the late Holocene: Paleocological insights from peatlands in the Alaskan Arctic. *Ecological Indicators*, 85, 525–536. <https://doi.org/10.1016/J.ECOLIND.2017.10.062>

- Galka, M., Szal, M., Watson, E. J., Gallego-Sala, A., Amesbury, M. J., Charman, D. J., et al. (2017). Vegetation succession, carbon accumulation and hydrological change in subarctic peatlands, Abisko, Northern Sweden. *Permafrost and Periglacial Processes*, 28(4), 589–604. <https://doi.org/10.1002/ppp.1945>
- Galka, M., Tobolski, K., Lamentowicz, Ł., Ersek, V., Jassey, V. E. J., van der Knaap, W. O., & Lamentowicz, M. (2017). Unveiling exceptional Baltic bog ecohydrology, autogenic succession and climate change during the last 2000 years in CE Europe using replicate cores, multi-proxy data and functional traits of testate amoebae. *Quaternary Science Reviews*, 156, 90–106. <https://doi.org/10.1016/j.quascirev.2016.11.034>
- Gallego-Sala, A. V., Charman, D. J., Brewer, S., Page, S. E., Prentice, I. C., Friedlingstein, P., et al. (2018). Latitudinal limits to the predicted increase of the peatland carbon sink with warming. *Nature Climate Change*, 8(10), 907–913. <https://doi.org/10.1038/s41558-018-0271-1>
- Geruo, A., Wahr, J., & Zhong, S. (2013). Computations of the viscoelastic response of a 3-D compressible Earth to surface loading: An application to glacial isostatic adjustment in Antarctica and Canada. *Geophysical Journal International*, 192(2), 557–572. <https://doi.org/10.1093/gji/ggs030>
- Gibson, C. M., Chasmer, L. E., Thompson, D. K., Quinton, W. L., Flannigan, M. D., & Olefeldt, D. (2018). Wildfire as a major driver of recent permafrost thaw in boreal peatlands. *Nature Communications*, 9(1), 3041. <https://doi.org/10.1038/s41467-018-05457-1>
- Gignac, L. D., Vitt, D. H., Zoltai, S. C., & Bayley, S. E. (1991). Bryophyte response surfaces along climatic, chemical, and physical gradients in peatlands of western Canada. *Nova Hedwigia*, 53, 27–71.
- Glenn, M. S., & Woo, M. (1997). Spring and summer hydrology of a valley-bottom wetland, Ellesmere Island, Northwest Territories, Canada. *Wetlands*, 17(2), 321–329. <https://doi.org/10.1007/BF03161420>
- Grosse, G., Goetz, S., McGuire, A. D., Romanovsky, V. E., & Schuur, E. A. G. (2016). Changing permafrost in a warming world and feedbacks to the Earth system. *Environmental Research Letters*, 11(4), 040201. <https://doi.org/10.1088/1748-9326/11/4/040201>
- Higuera, P. E., Brubaker, L. B., Anderson, P. M., Brown, T. A., Kennedy, A. T., & Hu, F. S. (2008). Frequent fires in ancient shrub tundra: Implications of paleorecords for Arctic environmental change. *PLoS ONE*, 3(3), e0001744. <https://doi.org/10.1371/journal.pone.0001744>
- Hill, G. B., & Henry, G. H. R. (2011). Responses of High Arctic wet sedge tundra to climate warming since 1980. *Global Change Biology*, 17(1), 276–287. <https://doi.org/10.1111/j.1365-2486.2010.02244.x>
- Hua, Q., Barbetti, M., & Rakowski, A. Z. (2013). Atmospheric radiocarbon for the period 1950–2010. *Radiocarbon*, 55(04), 2059–2072. https://doi.org/10.2458/azu_js_rc.v55i2.16177
- James, T. S., Henton, J. A., Leonard, L. J., Darlington, A., Forbes, D. L., & Craymer, M. (2015). Tabulated values of relative sea-level projections in Canada and the adjacent mainland United States; Geological Survey of Canada, Open File 7942, 81. <https://doi.org/10.4095/297048>
- James, T. S., Simon, K. M., Forbes, D. L., Dyke, A. S., & Mate, D. J. (2011). Sea-level projections for five pilot communities of the Nunavut Climate change partnership. Geological Survey of Canada, Open File 6715, 23. <https://doi.org/10.4095/288019>
- Jefferies, R. L., & Rockwell, R. F. (2002). Foraging geese, vegetation loss and soil degradation in an Arctic salt marsh. *Applied Vegetation Science*, 5(1), 7–16. <https://doi.org/10.1111/j.1654-109X.2002.tb00531.x>
- Jefferies, R. L., Rockwell, R. F., & Abraham, K. F. (2004). The embarrassment of riches: Agricultural food subsidies, high goose numbers, and loss of Arctic wetlands—A continuing saga. *Environmental Reviews*, 11(4), 193–232. <https://doi.org/10.1139/a04-002>
- Johannessen, O. M., Bengtsson, L., Miles, M. W., Kuzmina, S. I., Semenov, V. A., Alekseev, G. V., et al. (2004). Arctic climate change: observed and modelled temperature and sea-ice variability. *Tellus A: Dynamic Meteorology and Oceanography*, 56(4), 328–341. <https://doi.org/10.3402/tellusa.v56i4.14418>
- Jorgenson, M. T., Kanevskiy, M., Shur, Y., Moskalenko, N., Brown, D. R. N., Wickland, K., et al. (2015). Role of ground ice dynamics and ecological feedbacks in recent ice wedge degradation and stabilization. *Journal of Geophysical Research: Earth Surface*, 120, 2280–2297. <https://doi.org/10.1002/2015JF003602>
- Jorgenson, M. T., Shur, Y. L., & Pullman, E. R. (2006). Abrupt increase in permafrost degradation in Arctic Alaska. *Geophysical Research Letters*, 33, L02503. <https://doi.org/10.1029/2005GL024960>
- Kattsov, V. M., Walsh, J. E., Chapman, W. L., Govorkova, V. A., Pavlova, T. V., & Zhang, X. (2007). Simulation and projection of Arctic freshwater budget components by the IPCC AR4 Global Climate Models. *Journal of Hydrometeorology*, 8(3), 571–589. <https://doi.org/10.1175/JHM575.1>
- Kerbes, R. H., Meeres, K. M., & Alisauskas, R. T. (2014). *Surveys of Nesting Lesser Snow Geese and Ross's Geese in Arctic Canada, 2002–2009, Arctic Goose Joint Venture Special Publication*. Washington, DC, and Ottawa, Ontario: U.S. Fish and Wildlife Service and Canadian Wildlife Service.
- Killick, R., & Eckley, I. A. (2014). changepoint: An R Package for changepoint analysis. *Journal of Statistical Software*, 58(3), 1–19. <https://doi.org/10.18637/jss.v058.i03>
- Klinger, L. F., & Short, S. K. (1996). Succession in the Hudson Bay Lowland, Northern Ontario, Canada. *Arctic and Alpine Research*, 28(2), 172–183. <https://doi.org/10.1080/00040851.1996.12003163>
- Kopec, B. G., Feng, X., Michel, F. A., & Posmentier, E. S. (2016). Influence of sea ice on Arctic precipitation. *Proceedings of the National Academy of Sciences of the United States of America*, 113(1), 46–51. <https://doi.org/10.1073/pnas.1504633113>
- Liljedahl, A. K., Boike, J., Daanen, R. P., Fedorov, A. N., Frost, G. V., Grosse, G., et al. (2016). Pan-Arctic ice-wedge degradation in warming permafrost and its influence on tundra hydrology. *Nature Geoscience*, 9(4), 312–318. <https://doi.org/10.1038/ngeo2674>
- Mack, M. C., Bret-Harte, M. S., Hollingsworth, T. N., Jandt, R. R., Schuur, E. A. G., Shaver, G. R., & Verbyla, D. L. (2011). Carbon loss from an unprecedented Arctic tundra wildfire. *Nature*, 475(7357), 489–492. <https://doi.org/10.1038/nature10283>
- McBean, G., Alekseev, G., Chen, D., Førland, E., Fyfe, J., Groisman, P. Y., et al. (2005). Arctic Climate: Past and Present. In C. Symon, L. Arris & B. Heal (Eds.), *Arctic Climate Impact Assessment* (pp. 22–60). Cambridge, United Kingdom: Cambridge University Press.
- Minke, M., Donner, N., Karpov, N., de Klerk, P., & Joosten, H. (2009). Patterns in vegetation composition, surface height and thaw depth in polygon mires in the Yakutian Arctic (NE Siberia): A microtopographical characterisation of the active layer. *Permafrost and Periglacial Processes*, 20(4), 357–368. <https://doi.org/10.1002/ppp.663>
- Morris, P. J., Swindles, G. T., Valdes, P. J., Ivanovic, R. F., Gregoire, L. J., Smith, M. W., et al. (2018). Global peatland initiation driven by regionally asynchronous warming. *Proceedings of the National Academy of Sciences of the United States of America*, 115(19), 4851–4856. <https://doi.org/10.1073/pnas.1717838115>
- Myers-Smith, I. H., Forbes, B. C., Wilmking, M., Hallinger, M., Lantz, T., Blok, D., et al. (2011). Shrub expansion in tundra ecosystems: Dynamics, impacts and research priorities. *Environmental Research Letters*, 6(4), 045509. <https://doi.org/10.1088/1748-9326/6/4/045509>
- Myers-Smith, I. H., Harden, J. W., Wilmking, M., Fuller, C. C., McGuire, A. D., & Chapin, F. S. I. (2008). Wetland succession in a permafrost collapse: interactions between fire and thermokarst. *Biogeosciences*, 5(12), 1273–1286.

- Natali, S. M., Schuur, E. A. G., Mauritz, M., Schade, J. D., Celis, G., Crummer, K. G., et al. (2015). Permafrost thaw and soil moisture driving CO₂ and CH₄ release from upland tundra. *Journal of Geophysical Research: Biogeosciences*, 120, 525–537. <https://doi.org/10.1002/2014JG002872>
- Nicholson, B. J., & Gignac, L. D. (1995). Ecotope dimensions of peatland bryophyte indicator species along gradients in the Mackenzie River Basin, Canada. *The Bryologist*, 98(4), 437. <https://doi.org/10.2307/3243583>
- Oechel, W. C., Vourlitis, G. L., Hastings, S. J., Ault, R. P., & Bryant, P. (1998). The effects of water table manipulation and elevated temperature on the net CO₂ flux of wet sedge tundra ecosystems. *Global Change Biology*, 4(1), 77–90. <https://doi.org/10.1046/j.1365-2486.1998.00110.x>
- Payette, S., Delwaide, A., Caccianiga, M., & Beauchemin, M. (2004). Accelerated thawing of subarctic peatland permafrost over the last 50 years. *Geophysical Research Letters*, 31, L18208. <https://doi.org/10.1029/2004GL020358>
- Peltier, W. R. (2004). Global glacial isostasy and the surface of the ice-age Earth: The ICE-5G (VM2) model and GRACE. *Annual Review of Earth and Planetary Sciences*, 32(1), 111–149. <https://doi.org/10.1146/annurev.earth.32.082503.144359>
- Perreault, N., Lévesque, E., Fortier, D., Gratton, D., & Lamarque, L. J. (2017). Remote sensing evaluation of High Arctic wetland depletion following permafrost disturbance by thermo-erosion gully processes. *Arctic Science*, 3(2), 237–253. <https://doi.org/10.1139/as-2016-0047>
- Peterson, S. L., Rockwell, R. F., Witte, C. R., & Koons, D. N. (2013). The Legacy of destructive snow goose foraging on supratidal marsh habitat in the Hudson Bay Lowlands. *Arctic, Antarctic, and Alpine Research*, 45(4), 575–583. <https://doi.org/10.1657/1938-4246.45.4.575>
- Ping, C. L., Michaelson, G. J., Kimble, J. M., Romanovsky, V. E., Shur, Y. L., Swanson, D. K., & Walker, D. A. (2008). Cryogenesis and soil formation along a bioclimate gradient in Arctic North America. *Journal of Geophysical Research*, 113, G03S12. <https://doi.org/10.1029/2008JG000744>
- Polyakov, I. V., Bekryaev, R. V., Alekseev, G. V., Bhatt, U. S., Colony, R. L., Johnson, M. A., et al. (2003). Variability and trends of air temperature and pressure in the maritime Arctic, 1875–2000. *Journal of Climate*, 16(12), 2067–2077. [https://doi.org/10.1175/1520-0442\(2003\)016<2067:VATOAT>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<2067:VATOAT>2.0.CO;2)
- R Core Team (2018). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from. <https://www.r-project.org>
- Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Ramsey, C. B., et al. (2013). IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon*, 55(4), 1869–1887. https://doi.org/10.2458/azu_js_rc.55.16947
- Schädel, C., Bader, M. K. F., Schuur, E. A. G., Biasi, C., Bracho, R., Čapek, P., et al. (2016). Potential carbon emissions dominated by carbon dioxide from thawed permafrost soils. *Nature Climate Change*, 6(10), 950–953. <https://doi.org/10.1038/nclimate3054>
- Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., et al. (2015). Climate change and the permafrost carbon feedback. *Nature*, 520(7546), 171–179. <https://doi.org/10.1038/nature14338>
- Smith, L. C., Sheng, Y., & MacDonald, G. M. (2007). A first pan-Arctic assessment of the influence of glaciation, permafrost, topography and peatlands on Northern Hemisphere lake distribution. *Permafrost and Periglacial Processes*, 18(2), 201–208. <https://doi.org/10.1002/ppp.581>
- Speed, J. D. M., Cooper, E. J., Jónsdóttir, I. S., van der Wal, R., & Woodin, S. J. (2010). Plant community properties predict vegetation resilience to herbivore disturbance in the Arctic. *Journal of Ecology*, 98(5), 1002–1013. <https://doi.org/10.1111/j.1365-2745.2010.01685.x>
- Steere, W. C., & Scotter, G. W. (1979). Bryophytes of Banks Island, Northwest Territories, Canada. *Canadian Journal of Botany*, 57(10), 1136–1149. <https://doi.org/10.1139/b79-137>
- Tape, K., Sturm, M., & Racine, C. (2006). The evidence for shrub expansion in Northern Alaska and the Pan-Arctic. *Global Change Biology*, 12(4), 686–702. <https://doi.org/10.1111/j.1365-2486.2006.01128.x>
- Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., & Zimov, S. (2009). Soil organic carbon pools in the northern circumpolar permafrost region. *Global Biogeochemical Cycles*, 23, GB2023. <https://doi.org/10.1029/2008GB003327>
- Teltewskoi, A., Beermann, F., Beil, I., Bobrov, A., de Klerk, P., Lorenz, S., et al. (2016). 4000 years of changing wetness in a permafrost polygon peatland (Kytalyk, NE Siberia): A comparative high-resolution multi-proxy study. *Permafrost and Periglacial Processes*, 27(1), 76–95. <https://doi.org/10.1002/ppp.1869>
- Vardy, S. R., Warner, B. G., & Asada, T. (2005). Holocene environmental change in two polygonal peatlands, south-central Nunavut, Canada. *Boreas*, 34(3), 324–334. <https://doi.org/10.1111/j.1502-3885.2005.tb01104.x>
- Walker, D. A., Reynolds, M. K., Daniëls, F. J. A., Einarsson, E., Elvebakk, A., Gould, W. A., et al., & The other members of the CAVM Team (2005). The circumpolar Arctic vegetation map. *Journal of Vegetation Science*, 16(3), 267–282. <https://doi.org/10.1111/j.1654-1103.2005.tb02365.x>
- Woo, M., & Young, K. L. (2006). High Arctic wetlands: Their occurrence, hydrological characteristics and sustainability. *Journal of Hydrology*, 320(3–4), 432–450. <https://doi.org/10.1016/J.JHYDROL.2005.07.025>
- Woo, M., & Young, K. L. (2014). Disappearing semi-permanent snow in the High Arctic and its consequences. *Journal of Glaciology*, 60(219), 192–200. <https://doi.org/10.3189/2014JoG13J150>
- Wood, K. R., & Overland, J. E. (2010). Early 20th century Arctic warming in retrospect. *International Journal of Climatology*, 30(9), 1269–1279. <https://doi.org/10.1002/joc.1973>
- Zhang, H., Gallego-Sala, A. V., Amesbury, M. J., Charman, D. J., Piilo, S. R., Väilänta, M. M., & Zhang, H. (2018). Inconsistent response of Arctic permafrost peatland carbon accumulation to warm climate phases. *Global Biogeochemical Cycles*, 32, 1605–1620. <https://doi.org/10.1029/2018GB005980>
- Zhang, H., Piilo, S. R., Amesbury, M. J., Charman, D. J., Gallego-Sala, A. V., & Väilänta, M. M. (2018). The role of climate change in regulating Arctic permafrost peatland hydrological and vegetation change over the last millennium. *Quaternary Science Reviews*, 182, 121–130. <https://doi.org/10.1016/J.QUASCIREV.2018.01.00>

References From the Supporting Information

- Amesbury, M. J., Mallon, G., Charman, D. J., Hughes, P. D. M., Booth, R. K., Daley, T. J., & Garneau, M. (2013). Statistical testing of a new testate amoeba-based transfer function for water-table depth reconstruction on ombrotrophic peatlands in north-eastern Canada and Maine, United States. *Journal of Quaternary Science*, 28(1), 27–39. <https://doi.org/10.1002/jqs.2584>
- Beyens, L., & Chardez, D. (1995). An annotated list of testate amoebae observed in the Arctic between the longitudes 27° E and 168° W. *Archiv für Protistenkunde*, 146(2), 219–233. [https://doi.org/10.1016/S0003-9365\(11\)80114-4](https://doi.org/10.1016/S0003-9365(11)80114-4)

- Beyens, L., Chardez, D., de Baere, D., de Bock, P., & Jaques, E. (1990). Ecology of terrestrial testate amoebae assemblages from coastal Lowlands on Devon Island (NWT, Canadian Arctic). *Polar Biology*, 10(6), 431–440. <https://doi.org/10.1007/BF00233691>
- Bick, H. (1972). *Ciliated Protozoa: An Illustrated Guide to the Species used as Biological Indicators in Freshwater Biology*. Geneva, Switzerland: World Health Organization.
- Bobrov, A. A., Charman, D. J., & Warner, B. G. (1999). Ecology of testate amoebae (Protozoa: Rhizopoda) on peatlands in Western Russia with special attention to niche separation in closely related taxa. *Protist*, 150(2), 125–136. [https://doi.org/10.1016/S1434-4610\(99\)70016-7](https://doi.org/10.1016/S1434-4610(99)70016-7)
- Booth, R. K. (2008). Testate amoebae as proxies for mean annual water-table depth in *Sphagnum*-dominated peatlands of North America. *Journal of Quaternary Science*, 23(1), 43–57. <https://doi.org/10.1002/jqs.1114>
- Charman, D. J., Blundell, A., & Abrupt Climate Change Over the European Land Mass project members (2007). A new European testate amoebae transfer function for palaeohydrological reconstruction on ombrotrophic peatlands. *Journal of Quaternary Science*, 22(3), 209–221. <https://doi.org/10.1002/jqs.1026>
- Charman, D. J., Hendon, D., & Woodland, W. A. (2000). The identification of testate amoebae (Protozoa: Rhizopoda) in peats: QRA Technical Guide No. 9. Quaternary Research Association. London: Quaternary Research Association. <https://doi.org/10.1016/j.quascirev.2004.02.012>
- Corliss, J. O. (1979). *The Ciliated Protozoa: Characterization, Classification and Guide to the Literature* (2nd ed.). Oxford, United Kingdom: Pergamon Press.
- de Vleeschouwer, F., Chambers, F. M., & Swindles, G. T. (2010). Coring and sub-sampling of peatlands for palaeoenvironmental research. *Mires and Peat*, 7, 1–10.
- Flora of North America Editorial Committee (2007). *Flora of North America North of Mexico, Volume 27: Bryophytes, Part 1: Mosses*. Oxford, United Kingdom: Oxford University Press.
- Grimm, E. C. (1987). CONISS: a FORTRAN 77 program for stratigraphically constrained cluster analysis by the method of incremental sum of squares. *Computers & Geosciences*, 13(1), 13–35. [https://doi.org/10.1016/0098-3004\(87\)90022-7](https://doi.org/10.1016/0098-3004(87)90022-7)
- Hadenäs, L. (2003). The European species of the Calliergon–Scorpidium–Drepanocladus complex, including some related or similar species. *Meylania*, 28, 1–116.
- Hammer, Ø., Harper, D. A., & Ryan, P. D. (2001). PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaentologia Electronica*, 4(1), 9.
- Juggins, S. (2007). C2 Version 1.5 User guide. Software for ecological and paleoecological data analysis and visualisation. Newcastle upon Tyne, UK: Newcastle University.
- Juggins, S. (2018). Rioja: Analysis of quaternary science data. R package version 0.9-15.1. Retrieved from. <https://cran.r-project.org/package=rioja>
- Kralik, U. (1961). Ein Beitrag zur Biologie von loricaten peritrichen Ziziaten, insbesondere von *Platycola truncata* Fromentel 1874. *Archiv für Protistenkunde*, 105, 201–258.
- Lamarre, A., Magnan, G., Garneau, M., & Boucher, É. (2013). A testate amoeba-based transfer function for paleohydrological reconstruction from boreal and subarctic peatlands in northeastern Canada. *Quaternary International*, 306, 88–96. <https://doi.org/10.1016/J.QUAINT.2013.05.054>
- Lamentowicz, L., Lamentowicz, M., & Gąbka, M. (2008). Testate amoebae ecology and a local transfer function from a peatland in western Poland. *Wetlands*, 28(1), 164–175. <https://doi.org/10.1672/07-92.1>
- Mauquoy, D., & van Geel, B. (2007). Mire and peat macros. In S. A. Elias (Ed.), *Encyclopedia of Quaternary Science* (pp. 2315–2336). Amsterdam, Netherlands: Elsevier.
- Microworld. (2018). Microworld: World of amoeboid organisms. Retrieved from <https://www.arcella.nl/>
- Mitchell, E. A. D., Buttler, A. J., Warner, B. G., & Gobat, J.-M. (1999). Ecology of testate amoebae (Protozoa: Rhizopoda) in *Sphagnum* peatlands in the Jura mountains, Switzerland and France. *Écoscience*, 6(4), 565–576. <https://doi.org/10.1080/11956860.1999.11682555>
- Nacken, M., & Reise, K. (2000). Effects of herbivorous birds on intertidal seagrass beds in the northern Wadden Sea. *Helgolander Marine Research*, 54(2–3), 87–94. <https://doi.org/10.1007/s101520050006>
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., et al. (2017). vegan: Community ecology package. Retrieved from <https://cran.r-project.org/package=vegan>
- Qin, Y., Mitchell, E. A. D., Lamentowicz, M., Payne, R. J., Lara, E., Gu, Y., et al. (2013). Ecology of testate amoebae in peatlands of central China and development of a transfer function for paleohydrological reconstruction. *Journal of Paleolimnology*, 50(3), 319–330. <https://doi.org/10.1007/s10933-013-9726-6>
- Reimer, P. J., Brown, T. A., & Reimer, R. W. (2004). Discussion: Reporting and calibration of post-bomb ^{14}C data. *Radiocarbon*, 46(3), 1299–1304. <https://doi.org/10.1017/S0033822200033154>
- Smith, A. J. E. (2004). *The Moss Flora of Britain and Ireland*, (2nd ed.). Cambridge, United Kingdom: Cambridge University Press.
- Smith, H. G. (1992). Distribution and ecology of the testate rhizopod fauna of the continental Antarctic zone. *Polar Biology*, 12(6–7), 629–634.
- Swindles, G. T., Amesbury, M. J., Turner, T. E., Carrivick, J. L., Woulds, C., Raby, C., et al. (2015). Evaluating the use of testate amoebae for palaeohydrological reconstruction in permafrost peatlands. *Palaogeography, Palaeoclimatology, Palaeoecology*, 424, 111–122. <https://doi.org/10.1016/J.PALAEO.2015.02.004>
- Taylor, L. S., Swindles, G. T., Morris, P. J., & Galka, M. (2019). Ecology of peatland testate amoebae in the Alaskan continuous permafrost zone. *Ecological Indicators*, 96, 153–162. <https://doi.org/10.1016/J.ECOLIND.2018.08.049>
- Vardy, S. R., Warner, B. G., Turunen, J., & Aravena, R. (2000). Carbon accumulation in permafrost peatlands in the Northwest Territories and Nunavut, Canada. *The Holocene*, 10(2), 273–280. <https://doi.org/10.1191/095968300671749538>
- Warner, B. G., & Charman, D. J. (1994). Holocene changes on a peatland in northwestern Ontario interpreted from testate amoebae (Protozoa) analysis. *Boreas*, 23(3), 270–279. <https://doi.org/10.1111/j.1502-3885.1994.tb00949.x>
- Warner, B. G., & Chengalath, R. (1991). *Habrotricha angusticollis* (Bdelloidea, Rotifera): A new paleoecological indicator in Holocene peat deposits in Canada. *SIL Proceedings*, 24(5), 2738–2740. <https://doi.org/10.1080/03680770.1989.11899146>