REVIEW ARTICLE

What is an appropriate caries diagnosis?

VIBEKE BÆLUM^{1,2}¹*School of Dentistry and* ²*Department of Epidemiology, School of Public Health, Faculty of Health Sciences, Aarhus University, Aarhus, Denmark*

Abstract

This aim of this paper is to spur a discussion of the direction of caries-lesion detection activities in clinical dental practice. It is argued that since the dental clinician's caries-related decision making is a script-matching enterprise in which clinical decisions are made on the basis of 'this-lesion-needs-this-kind-of-treatment' reasoning, the methods and strategies employed for caries lesion detection should accommodate this fact. This may be done by employing a clinical visual–tactile method for caries lesion detection that evaluates the two aspects that are crucial for appropriate caries management: lesion activity and surface integrity. The use of diagnostic methods that do not assess these features directly but involve assumptions about activity status and surface integrity should be avoided. This includes the use of bite-wing radiography for the detection of approximal caries lesions, as it may be shown that plain reliance on radiographs leads to considerable overtreatment. If clinical dentistry is to retain its status as a profession committed to doing good, changes in diagnostic practices along these lines are warranted.

Key Words: *Adverse effects, caries, clinical practice, diagnostic methods, screening*

Introduction

In the course of just a few years, several conferences have been held [1–3] and books have been written [4] that have addressed the issue of how best to go about diagnosing caries in various settings, which may include daily dental practice, clinical research, epidemiology or population oral health surveillance programmes. However, among the settings listed, the daily dental practice setting remains the most important since the reasons why we continue to carry out clinical and epidemiological caries research and continue to run population oral health surveillance programmes all originate in the fundamental fact that people, and in some populations all people, continue to experience new dental caries lesions over time. Dental practitioners may play a central role in helping patients to control their caries situation, as they may be able to observe caries lesions long before they are sufficiently advanced to produce the symptoms that would otherwise prompt the patient to seek treatment. Thereby, the key 'raison d'être' for the dental profession, which also includes those of us who are involved in clinical and epidemiologic research on dental

caries, is our ability to help people control their caries situation. In most countries, the organization of the dental healthcare services is based on the idea that the general dental practitioner constitutes the key to the control and management of dental caries, and it is for this reason that any discussion of better caries diagnostic methods should take as their starting point what goes on in the meeting between the patient and their general dental practitioner.

Clarification of terms

As used here, the term 'patient' refers to any member of the public who is consulting a general dental practitioner. Patients may do so for one of two principal reasons. The patient has either gone to have their regular dental check-up, or they have gone because of a tangible problem for which they seek assistance from the dental practitioner. In the check-up situation the patient has no perception of any dental treatment being necessary but goes in the hope of being told that all is well, and possibly also to receive preventive care, which may include variable combinations of

Correspondence: Vibeke Baelum, Department of Epidemiology, School of Public Health, Aarhus University, Bartholins Alle 2, DK-8000 Aarhus C, Denmark.
E-mail: baelum@soci.au.dk

(Received 3 December 2009; accepted 4 December 2009)

ISSN 0001-6357 print/ISSN 1502-3850 online © 2010 Informa UK Ltd. (Informa Healthcare, Taylor & Francis AS)

DOI: 10.3109/00016350903530786

fluoride applications, scaling and polishing, fissure sealants and preventive advice [5,6]. The core activity carried out by the practitioner during the check-up visit is a dental examination of the oral mucosa, the periodontium and the teeth. Thereby, the check-up visit effectively becomes an oral screening procedure [6,7]. In the check-up situation, differential diagnostic considerations play little role except for possible mucosal lesions. When it comes to the periodontium and the teeth, the dental practitioners know exactly what they are looking for, and in the case of dental caries the objects sought for are caries lesions in need of treatment.

There is a growing recognition among researchers in the field of dental caries diagnosis that the same terms may have very different meanings in different quarters [8–10]. A higher degree of precision in the terminology used has therefore been advocated [11]. Even so, the terms used in the field of caries diagnosis continue to be an object of confusion, and despite several attempts to delineate their precise distinction [1,8,9,11,12] it is not evident how all of the terms ‘caries measurement’, ‘caries scoring’, ‘caries detection’, ‘caries diagnosis’, ‘caries assessment’ and ‘caries monitoring’ are helpful in bringing clarity to the field of caries diagnosis. In this review, the term ‘caries detection’ denotes the process of identifying caries lesions in need of treatment during a clinical examination in a general dental practice setting. The key reasons for not using the term ‘caries diagnosis’ are that proper caries diagnoses are rarely made in clinical dentistry [13,14] and differential diagnostic considerations are greatly downplayed and replaced by a pattern-recognition process [13,15].

Caries detection—caries script matching

Dental school taught us that any diligently exercised caries examination begins with a careful visual–tactile inspection, in which each single tooth surface in the dentition is evaluated in a standardized and sequential manner for deviations from its normal appearance with respect to colour, lustre, surface texture and surface integrity [16]. Thereby, a single caries examination involves the visual–tactile inspection of up to 148 tooth surfaces in the patient. As a normal working day in general dental practice typically involves the examination of dozens of patients, it is hardly surprising that this diagnostic process soon becomes rather more automated, and that the discrete differential diagnostic steps outlined above are no longer undertaken. Instead, the caries diagnostic process soon develops into a pattern-recognition process, which means that each tooth surface is scanned for its similarity to one of a finite set of clinical patterns or caries scripts retained in the clinician’s mind.

A caries script is a highly summarized version of the clinician’s cumulative experience with similar clinical presentations [13]. If, during inspection, a departure from ‘normal’ is detected, a rapid and unconscious review is undertaken of the dental clinician’s inventory of caries scripts to establish if one of them matches the presentation. Importantly, once a matching caries script has been identified, the treatment decision is often given by virtue of the script. Thereby, diagnostic considerations are altogether skipped and replaced by scripts in the form of ‘this-type-of-lesion-needs-this-kind-of-treatment’ decisions [10,13,14], and dentists often ‘diagnose’ lesions for treatment using the intended treatment to identify and describe the observation in charts and patient records [13,17].

If the observations made during clinical inspection do not match one of the many caries scripts retained in the clinician’s mind, the clinician is typically prompted to collect additional information using ancillary diagnostic devices. Unless the use of bite-wing radiographic information is already incorporated in the script-matching process, which may indeed occur [18,19], the ancillary information would most commonly be provided by the use of bite-wing radiographs, although other sense-enhancing methods might also be considered. The clinician scans the radiographic image and uses this additional information to improve the matching of the clinical and radiographic presentations to one of the caries scripts.

The key question is whether our caries-related treatment decisions inevitably result in the best decisions. Unfortunately, the answer to this rhetorical question is ‘no’. This conclusion follows from the numerous observations that dentists vary considerably in their caries detection and treatment decisions [14,20–43] for the same teeth and the same patients. It is just a matter of simple logic to conclude that if two dentists disagree on the presence of a lesion on a surface, at least one of them is wrong. Similarly, if two dentists agree on the presence of a lesion but decide to intervene using different treatments, it logically follows that one of them has provided a less-desirable treatment [14,44,45]. It follows that the seemingly simple and straightforward process of detection of deviations from normal and the sorting of abnormal observations into distinct ‘this-lesion-needs-this-kind-of-treatment’ groups hides several problems that jeopardize the value of the activity.

As shown in Figure 1, the many sources of influence on the caries-related treatment decisions made in clinical dental practice broadly fall into two categories: the patient factors embrace the nature and strength of signals emanating from the patient presentation, whereas the dentist factors include the elements that form and shape the dentist’s inventory of caries scripts. It is within the realms of dentistry to control the dentist factors, and for reasons that will be detailed later in this article it is high time that the

Figure 1. Conceptual model explaining how dentists make caries-related treatment decisions. Adapted from Bader & Shugars [14].

profession embarks on this enterprise. A lot could and should be done at the levels of dental education and the dental faculty [10,14,44]. If the problem is to be solved before dentists become branded as dishonest [46] salesmen [47] who are more devoted to personal income and leisurely living than to their professional obligations [48], we must also address the dentists already in clinical dental practice. By virtue of their position these dental practitioners are strong influences for changes in the caries-related treatment practices of the newly graduated dentist, and may therefore easily undermine/overrule what has been taught at dental school. Unfortunately, established practitioners are most unlikely to change their caries-detection behaviours, and it has therefore been suggested that a single professional standard be developed that minimizes the need for subjective interpretation [14]. While this proposal conflicts with the dental professional's implicit freedom to exercise judgment in their work (i.e. the 'art of dentistry'), it is equally clear that some regulation would benefit the interests of the public.

Finding caries lesions that match our treatment options

Since the clinical caries examination is a lesion-detection process involving 'this-lesion-needs-this-kind-of-treatment' decisions, improvement of our caries-related decisions must begin with an overview

of the caries-lesion treatment options that are available to the dental clinician. Step two involves the matching of these treatment options to the different types of caries lesions that may be encountered in the clinic.

Broadly speaking, the caries-lesion treatment armamentarium comprises two distinct forms of treatment: operative and non-operative treatment. While it is recognized that each of these options involves a myriad of alternatives and choices regarding agents and materials, techniques and methods and frequency and dosage, it is also clear that the treatment-decision principles that emanate from these two forms are very simple. Hence, it is generally agreed that a cavitated caries lesion necessitates the insertion of a restoration, the purpose of which is to make the tooth surface amenable to cleansing and to protect the pulpo-dentinal complex [49]. Since cavitated caries lesions may indeed be arrested if they are located in surfaces that are accessible for daily oral hygiene procedures [50,51], the dental clinician may well choose the non-operative treatment route for the motivated patient. Restoring non-cavitated lesions is considered inappropriate since such lesions may be arrested using non-operative procedures [52,53]. Non-cavitated lesions comprise two forms: the arrested and the active caries lesion. There is clearly no reason to institute professional treatment of the arrested caries lesion, whereas it is exceedingly important to be able to control the active non-cavitated caries lesion by non-operative intervention because of an otherwise poor prognosis [54].

The caries-lesion detection process is therefore a matter of detecting caries lesions that match one of these two basic treatment options, and this involves the distinction of cavitated lesions, non-cavitated active lesions and arrested (cavitated and non-cavitated) lesions from the normal appearance of the tooth surface. Only the visual-tactile caries-lesion detection method can achieve this directly. The use of any other caries-lesion detection methods involves assumptions, because these methods are based on physical principles that do not adequately reflect the decisive issues of surface integrity and lesion-activity status. For example, the information provided by bite-wing radiography relates to lesion depth, but there is no unequivocal link between radiographic lesion depth and lesion characteristics in terms of surface integrity, and the bite-wing radiograph provides no information whatsoever related to caries-lesion activity status [55]. The widely held idea that a radiographic lesion extending into dentine is an indication of lesion cavitation has been largely undermined by research in the field [40,56–65], but remains deeply ingrained in the minds of most dental clinicians. It is equally clear that bite-wing radiographs fail to disclose early caries lesions [62,66–70], i.e. precisely those lesions that are amenable to non-operative caries-lesion control. In spite of these reservations, caries-lesion detection in clinical dental practice continues to be based on a combination of (at least) two diagnostic tests: the visual-tactile examination and the bite-wing radiographic examination.

Value of the caries diagnostic test

The value of the caries diagnostic test has most commonly been evaluated by assessing its criterion validity, i.e. by comparing the test result to a histological 'gold standard' [71,72] thought to represent the caries diagnostic 'truth'. However, the de- and remineralization processes involved in caries-lesion formation are dynamic and as long as the resulting

net mineral gains or losses stay within limits that do not result in clinically discernible lesion formation or jeopardize the integrity of the tooth, we need not be concerned [10,73]. Moreover, the dynamic nature of the de- and remineralization processes has the additional effect of rendering the 'true' state rather elusive with respect to mineral loss, and the search for a caries diagnostic 'truth' therefore makes little sense [10,73].

Despite the frequent statement that bite-wing radiographs are valuable for caries diagnosis [74] a simple example will reveal how wrong this perception may be. A systematic review [75] concluded that the sensitivity and specificity values for bite-wing radiography in the detection of approximal cavities were 0.66 and 0.95, respectively. Depending on the true, but unknown, prevalence of approximal cavitated caries lesions in the population, the bite-wing radiographic test yields the results shown in Table I. Using the 1% true cavity prevalence as the example, and assuming that the dental clinician decides to restore all the bite-wing radiography-positive surfaces, it may be calculated that a total of 561 (66+495) surfaces will be restored. Among these restored surfaces, 495 (88%) have been unnecessarily restored, i.e. they represent overtreatment. Considering the true cavities, on the other hand, 34 (34%) will remain undiagnosed and thereby undertreated. The total number of decisional errors made amounts to 529 (34+495), and the main result of using bite-wing radiographs for approximal cavity detection is a considerably invasive overtreatment owing chiefly to the less-than-perfect specificity value of 0.95.

The visual-tactile examination has also been evaluated in terms of its diagnostic sensitivity and specificity for the detection of approximal cavities, yielding a sensitivity of 0.52 and a specificity of 0.98 [71]. As shown in Table II, the number of diagnostic errors is much smaller than for bite-wing radiographs, owing to the slightly higher specificity. Using once again the 1% true cavity prevalence to highlight the results, the Table shows that 250 (52+198) bite-wing radiography-positive surfaces will be restored, although 198 (79%) of these surfaces have been overtreated. A total

Table I. Diagnostic results obtained when applying bitewing radiography (BW) for the detection of approximal cavities in a population of 10,000 approximal surfaces. Given for different values of the true, but unknown, prevalence of approximal cavitation. Sensitivity = 66%, specificity = 95% [71].

Cavity prevalence (<i>n</i> = 10,000) (%)	No. of BW+ cavities	No. of BW- cavities	Total no. of cavities	No. of BW+ intact surfaces	No. of BW- intact surfaces	Total no. of intact surfaces
0.1	6.6	3.4	10	585	9405	9990
0.5	33	17	50	497	9453	9950
1	66	34	100	495	9405	9900
2.5	165	85	250	487	9263	9750
5	330	170	500	475	9025	9500

Table II. Diagnostic results obtained when applying visual–tactile (VT) methods for the detection of approximal cavities in a population of 10,000 approximal surfaces. Given for different values of the true, but unknown, prevalence of approximal cavitation. Sensitivity = 52%, specificity = 98% [71].

Cavity prevalence ($n = 10,000$) (%)	No. of VT+ cavities	No. of VT– cavities	Total no. of cavities	No. of VT+ intact surfaces	No. of VT– intact surfaces	Total no. of intact surfaces
0.1	5.2	4.8	10	200	9790	9990
0.5	26	24	50	199	9751	9950
1	52	48	100	198	9702	9900
2.5	130	120	250	195	9555	9750
5	260	240	500	190	9310	9500

of 48 cavities (48%) will remain undiagnosed and thus undertreated, but the total number of decisional errors amounts to 246 (48+198) and is therefore less than half of the errors committed using the bite-wing radiographs.

The results shown in Tables I and II also demonstrate how misleading is the concept of diagnostic yield. The all-pervading argument in favour of the use of bite-wing radiography for the detection of caries lesions is that thereby more lesions can be detected [10], and it is implicit that the additional lesions detected are real lesions that would have gone undetected had bite-wing radiographs not been used [74,76]. However, as shown in Tables I and II, the additional cavitated lesions detected by bite-wing radiography are largely lesions that are not truly present, i.e. false-positive diagnoses. In fact, considering once again the 1% true cavity prevalence example, the tally shows that bite-wing radiographs lead to correct detection of an additional 14 cavities compared with the visual–tactile examination, but this occurs at the expense of the additional ‘detection’ of 297 non-existing cavities. Thereby, for each true cavity that has been ‘saved’ from the perils of going undetected, 21 intact surfaces must suffer the rather deleterious restorative consequences of false cavity detection.

Bite-wing radiographs are usually used as an ancillary diagnostic procedure to the visual–tactile clinical examination, but to the best of my knowledge no studies have looked at the effect of combining test procedures on the treatment decisions made. However, a recent study explored the effects of two different diagnostic strategies in a context conforming with the script-matching caries-detection process [77]. Following a visual examination of each of 96 occlusal surfaces that presented without frank cavitation, each of three examiners was asked to choose between three treatment options: no treatment; non-operative treatment; or operative treatment for each surface. Four weeks later, this procedure was repeated, but this time the examiners had access to the test results from four ancillary tests: bitewing

radiographs, electric conductance measurement, quantitative light fluorescence and laser fluorescence. The availability of additional diagnostic information led to a substantial increase in invasive treatment recommendations, considerably more overtreatment and a reduced frequency of correct treatment recommendations [77].

While it may seem counterintuitive to many people that the use of more diagnostic tests does not necessarily lead to better decisions, the phenomenon is intimately linked with the suboptimal reliability and reproducibility of each diagnostic test [40,78]. When tests are added to each other, each of which has a low, but certainly discernible, error probability, the inevitable result is that more diagnostic and hence decisional errors will be made. This simple truth was pointed out by Murphy [79], who noted that whenever diagnostic tests are imperfect (i.e. having less than 100% sensitivity, specificity and reliability), a healthy person is just someone who has not been sufficiently tested. In the dental clinical scenario, multiple testing occurs either as described above, i.e. when several diagnostic tests are applied during the same clinical session, or when a single diagnostic test is applied over a number of consecutive sessions, i.e. semi-annually or annually, as is common practice for patients participating in routine recall/check-up schemes. It is thus a fact that the regular recall/routine dental attendance rates are rather high in many populations [80–89]. In Sweden, 90–95% of adults visit a dentist regularly, and since the 1980s >80% of appointments have been given at the dentist’s initiative [80].

Regular dental check-ups—inevitably a good thing to have?

Having regular check-up visits or participating in routine recall programmes should be understood and recognized as participation in regular screening programmes. The principles underlying a screening programme for dental caries are illustrated in

SV: Symptom-driven dental attendance pattern
 RV: Regular checkup dental attendance pattern

Figure 2. The principles underlining screening: three possible scenarios depending on earlier disease detection and treatment effect.

Figure 2. It is only for illustrative purposes that the unit considered here is the tooth, and the principles can easily be integrated to cover subjects or populations, just as events to avoid other than tooth extraction may be considered. In fact, in many contemporary populations a much more relevant event to avoid would be the first insertion of a restoration. The idea outlined in Figure 2 is that caries lesions can be detected by a dental professional during a routine check-up visit long before the patient would otherwise have noted the lesion. Thereby, the dental professional may institute a treatment of the lesion that may result in increased longevity of the tooth compared with the no-screening situation. However, as Figure 2 clearly shows, earlier detection of the caries lesion is not associated with a benefit to the patient unless the treatment that is instituted leads to postponed tooth extraction (or any other event to avoid). This means that the observation of a longer ‘survival time’ cannot be taken as evidence of a beneficial effect, but merely reflects the earlier detection of the lesion, a phenomenon called lead-time bias [90].

The claims of benefit made when the dental profession advocates regular dental check-ups can chiefly be described as better oral health conditions (than would have been the case had regular dental check-ups not been carried out). In this context it is noteworthy that the evidence for a beneficial effect of such regular dental check-ups is missing [6,91–93]. The problem is not a lack of studies, but a lack of studies with an appropriate study design. It is thus not difficult to find observational studies that report an association between regular dental attendance and better oral health conditions [81–83,94,95]. However, it is exceedingly difficult to control the biases

inherent in such observational studies [6] and the unequivocal evidence should come from high-quality randomized controlled clinical trials. However, such trials are largely missing [6,91,92].

There can be little doubt that dental professionals thrive on the existence of routine dental recall programmes. Moreover, the rather high routine recall attendance rates observed in many countries could be taken to indicate that check-ups are generally very well accepted and considered beneficial by the public, and a layperson might really wonder how could such regular dental check-ups not be beneficial when they are so strongly advocated by dental professionals? However, what the layperson does not perceive—because no-one tells them—is that like any other screening programme, the routine recall or regular check-up dental screening programme may also cause harm. Moreover, the layperson does not know that a profound belief exists throughout the health professions that early (or earlier) disease detection inarguably provides benefits to the screened populations [7]. This professional belief is so profound that screening programmes, such as those for breast cancer, have been launched in many countries in the absence of clear evidence of their beneficial effects [96]. Professionals tend to adopt a paternalistic “we know what is good for you” attitude and may greatly downplay the harmful effects of screening in the information materials that should allow people to decide whether or not to attend based on informed consent [97,98]. In breast cancer screening, it has thus been estimated that if 2000 women are screened regularly for 10 years, one woman will have her life prolonged as a result (that is, she will avoid dying from breast cancer, but may die from other causes), whereas another 10

healthy women will experience harm in the form of unnecessary breast surgery and radiotherapy [96]. In addition, about 200 women (i.e. 10%) are at risk of considerable and sustained psychological distress because they will receive a false-positive breast cancer diagnosis [99,100]. Paradoxically, in the absence of balanced and adequate information given to the person before they decide whether to attend a screening examination, the screened person will react positively to the screening, irrespective of the outcome. The person who was screened negative is happy to know that they are disease-negative (even though this could be a false-negative observation); the person who was screened positive, but in subsequent more detailed tests found to be negative, is relieved again after a period of worry; and the person who was confirmed disease-positive is happy because they have been operated on and saved from a much worse fate (even though this may have been a false-positive diagnosis). This merely shows that in the absence of a fair and balanced account of the benefits and harms of the screening procedure, the population uptake of a screening programme should not be used as an argument for that programme.

Whilst dental caries is in no way comparable to breast cancer, whether as a public health problem or as a personal problem for those affected, the breast cancer example highlights the issues of benefits and harms that should be addressed in any screening programme, including dental routine recall screening programmes. Moreover, in order that members of the public can make informed choices about whether to participate in screening/routine recall programmes or not it is crucial to be open and fair about the intended as well as unintended consequences of participation in the recall programme. If the dental profession is to retain its status as a profession, it is mandatory that it is perceived by the public as being strongly committed to the hallmarks of professionalism, and as clearly denouncing commercialism as a motive.

Patient autonomy and dental professionalism

The high adherence to routine recall visits in the form of annual or biannual dental screening visits may pose a problem from the public's point of view if these screening visits are unnecessary or if the visits are used for selfish professional advantage, e.g. by prescribing unnecessary treatments, which may impair prognosis [101–103], or by ordering unnecessary diagnostic procedures for which the patient must pay. While the patient may to some extent retain control over their demand for the dental services, i.e. the number of dental visits made, by deciding whether to book or honour an appointment or not, it is also clear that they stand much less chance of maintaining control over their utilization, i.e. the costs per visit, once sat in the

dental chair. Patients lack the appropriate knowledge and expertise needed to allow them to review their dentist and objectively assess the necessity for and the quality of the work provided [104,105], and this opens the possibility for the dentist to exercise supplier inducement [106–109]. Hence, the evidence shows that dentists are indeed able to control both the demand for [106,108] and the utilization of dental services [106–110]. The public has no choice but to trust that the dentist is not driven by selfish motives when advising and counselling them, and such trust is indeed the centrepiece of the social contract between dentistry and society that has granted dentistry the status of a profession [104,111]. This social contract with society implies that the professionals profess a commitment to competence, integrity and morality, altruism and promotion of the public good within their competence domain [104,105,112–114]. In return, the profession is granted monopoly over the use of its knowledge base in the form of autonomy in practice and self-regulation privileges [112,114].

Dentistry can maintain its status as a profession as long as the public trusts that dentists work for the good of the public and refrain from exploiting their monopoly and the vulnerability of the patients [105]. It is equally clear, however, that a conflict of interests exists between the public and the dentists by virtue of the fact that dentists generate their personal income from the services provided to the public. From an income-generation point of view, dentists who work under a fee-for-service reimbursement system have a vested economic interest in providing more services per patient [105], and the possibility to over-service patients is clearly present. Hence, while each single patient may not pay too much attention to the addition of a diagnostic service to the bill, the dentist is clearly knowledgeable about the implications of so doing for their own revenue.

In recent years, increasing concern has been voiced that dentistry is losing its status as a profession and is changing to become a for-profit enterprise [46–48,115]. Dentistry seems to be increasingly utilizing its monopolistic position to exercise commercialism, and from the point of view of continued public trust in the value of attending the dentist, and participating in regular check-up/routine recall screening programmes, such a change is clearly deeply problematic. Moreover, elective treatments such as cosmetic interventions constitute an increasing proportion of the total volume of dental services provided, but they cannot be considered indicated from a dental health perspective. This intermingling of aesthetic problems with oral health problems poses yet another threat to the status of the dental profession [104].

The rise of commercialism in healthcare and the formation of a healthcare marketplace is not reserved for dentistry, but has been noted in other fields. As an example, Hafferty [116] described the pervasiveness

of medical commercialism by stating that “medicine’s traditional ‘one or two bad apples’ has morphed into a megaorchard of physician clinicians and researchers, brimming with commercial proclivities, penchants, and practices” [116]. Such developments clearly pose a threat to the survival of the core professional privileges, which include discretionary decision making, occupational autonomy and the right to control recruitment, training and credentialing of new professionals. It is thus not difficult to imagine what could be termed a bureaucratic response (Table III) to the onwads march of commercialism within the health professions, in the form of more rules and regulations and codes of conduct and ethics being imposed by the state or insurance companies, i.e. outside the profession. Hence, one of the virtues of the bureaucracy is its ability to serve as a corrective to inappropriate and irresponsible discretion (Table III). Noting that the heart of what professionalism must defend is the economic privilege that follows from professionalism’s monopoly over its knowledge base, Freidson [117] predicted that unless current trends are opposed, the professionals are on a course of changing into neutral technical experts serving the needs of state and capital, which will increasingly gain control over performance and costs. Both Hafferty [116] and Freidson [117] noted how the “conspicuous absence of activities conscientiously enforcing professional codes of ethics” has

contributed to the vulnerability of the healthcare professions to attacks, and they conclude that the core of professionalism is the devotion to use disciplined knowledge for the public good.

The status of a profession is delegated to a discipline by the state or by capital [117] and is not an attribute that can be achieved unless the state or capital is convinced that the discipline is of special value to the public at large or to an important interest of the state. This delegation is entirely based on trust. Freidson [117] thus noted that “if professionalism is to be reasserted and regain some of its influence, it must not only elaborate and refine its codes of ethics but also strengthen its methods of adjudicating and correcting their violation”. This is a call for particular reinforcement of the institutional ethics, i.e. those ethics dealing with the economic, political, social and ideological circumstances that create many of the moral problems of work. If, therefore, dentistry is to survive as a profession, the dental associations have a vested interest in organizing a disciplinary dental practice that exercises responsible and accountable discretion in their decisions, which should be based on evidence. Systems should be devised to vigorously enforce the detailed codes of practice ethics, and dental associations should be seen by the public to take serious action whenever the privileged position of dental practitioners is abused for selfish purposes. Durkheim saw the professions as

Table III. Distinguishing characteristics of three ideal-type models that regulate provider–client relationships: commercialism, bureaucracy and professionalism [117].

	Commercialism (Adam Smith)	Bureaucracy (Max Weber)	Professionalism (Eliot Freidson)
Prototype characteristics	Free market	Civil service organizations	Professions, e.g. law, medicine, dentistry
Type of market	Full competition	–	Monopoly
Purpose	Profit maximization	For the public good	For the public good
Efficiency	High	Moderate	Moderate
Public prestige	Low	Moderate	High
Flexibility and responsiveness	High	Moderate	Low
Organization of work	Perfect freedom	Hierarchy, lines of authority Standardization of procedures	Discretionary judgment, based on abstract theory/concepts, even with manual tasks
Labour market characteristics			
Port of entry	Open	Personnel office (regulated)	Practice institution (closed)
Entry requirements	Consumer choices	Formal job description, by position	Training credentials
Typical career-line	Disorderly, irregular	Regular, vertical within firm	Horizontal, across firms
Predominant knowledge	Everyday	Variable by position, firm-specific	Discretionary, transferable
Consumer/client control			
Degree of control	Full consumer control	No consumer control, control exercised by managers	No consumer control, control exercised by professionals
Autonomy	Consumer autonomy in assessment of utility of goods and services	Moderate client autonomy in utility assessment	No consumer autonomy in utility assessment

the ideal organs for preventing the powerful “from tyrannizing over individuals” [118]. However, the dental profession needs to reassure the public that it is not a cat set among pigeons.

Caries detection in a changing disease panorama

Current developments in oral disease patterns offer an opportunity to test the dental profession’s ability to exercise appropriate and responsible discretion. Hence, the epidemiological evidence is overwhelming that the population prevalence, extent and severity of dental caries have declined significantly over past decades, not only in most populations but also in very diverse populations [119–164]. Although the decline has been particularly well documented among children in the high-income populations, the decline is also observed in moderate- and low-income countries, and has clearly also trickled into the adult population groups as well. Concomitant with the caries decline, a considerable improvement has been noted in tooth retention among the older age groups, where tooth loss used to be pronounced.

The data presented by Hugoson et al. [165] from the repeated cross-sectional epidemiological studies carried out in Jönköping, Sweden since 1973 may be used to illustrate how these changes have affected the ‘average’ dental patient in terms of their caries profile. In Figure 3, the mean dft/DFT counts observed among the Jönköping population have been adjusted to represent all persons rather than dentate persons only. Thereby, the Figure illustrates that, over recent decades, the age at which the dft/DFT experience peaks has moved upwards from 30 years in 1973 to 60 years in 2003. Moreover, the peak dft/DFT

experience has declined since the maximum was reached by the 40-year-olds in 1983 (Figure 3). The average dft/DFT experience has shown a steady decline since 1973 among the 10-, 20- and 30-year-olds, whereas the 40- and 50-year-olds show a mixed picture, with caries increases in the early part of the period followed by declines in the later survey years. Among the 60-, 70- and 80-year-olds a gradual increase in the DFT experience is noted (Figure 3), which is paralleled by the increasing tooth retention among these age groups (Figure 4). The cross-sectional epidemiological data presented by Hugoson and co-workers [161,165] thus serve to demonstrate that not only has the caries incidence rate (the number of new lesions per year) decreased substantially, but so has the caries progression rate, and in 2003, 80–90% of all caries lesions observed among the 3–20-year-olds were ‘initial’ lesions, i.e. non-cavitated caries lesions [165].

The improvements are dramatic and widespread, and it would seem prudent to keep a watchful eye on the extent to which these changes are paralleled by appropriate changes in the dental services rendered to the public. It would thus seem obvious that such dramatic changes should have profound implications for our caries-related clinical decisions. Dentists who wish to subscribe to appropriate dentistry and professionalism will fight their inherent temptation to respond to a biological problem using traditional technical solutions [44], and will increasingly turn in the direction of the non-operative caries management procedures. ‘Pre-emptive strikes’ in the form of operative (restorative) treatment given to caries lesions for which cavitation status, let alone activity status, is not clinically verified and is only assumed from ancillary diagnostic test results should no longer be regarded as appropriate caries management. The

Figure 3. Development over three decades in the dft/DFT counts among Swedes aged 10–80 years. Data from Hugoson et al. [165].

Figure 4. Development over three decades in the number of natural teeth present among Swedes aged 10–80 years. Data from Hugoson et al. [165].

Figure 5. Changes in the propensity of dentists in the Norwegian Dental Public Service to restore approximal surfaces among 15-year-olds as a function of the radiographic finding and the year of examination. Data from Gimmestad et al. [166].

substantially lower caries incidence and progression rates that increasingly characterize the populations imply that restorative treatment decisions should gradually be erased from the dental clinician's caries-controlling armamentarium.

Very little is known about general dental practitioners' responses to the considerable decline in dental caries that has been going on for nearly half a century. However, it is known from both Norway and Denmark that caries-related treatment practices have changed dramatically among dentists employed in the public dental services [166,167]. In Norway, the restorative treatment propensity in 1996 was only 16% of what it had been in 1979 [166] (Figure 5), and in Denmark the restorative treatment propensity

Figure 6. Changes in the propensity of dentists in the Danish Dental Public Service to restore approximal surfaces among 17-year-olds as a function of the radiographic finding and the year of examination. Data from Heidmann et al. [167].

in 1984–85 was only 17% of what it had been only 6 years before [167] (Figure 6). In the Danish study these changes were seen as partly resulting from the acknowledgment of slower caries progression rates and a more sceptical attitude towards the results of filling therapy [167], whereas the tentative explanation in the Norwegian study also included a higher dentist workload resulting in non-operative treatments being preferred over operative treatments [166]. It is clear, however, that the responsiveness demonstrated here for public dental services may well have been countered in private dental service settings by the wish to maintain a high income for a reasonable number of working hours. The fact is that dentistry may have arrived at a crossroads. The profession may choose the commercial way, in which case the public must be appropriately informed that dentistry is a for-profit enterprise; or it can choose the professional healthcare-oriented way and respond adequately and ethically to the altered population disease profiles. This involves acceptance that fewer dental services are needed for current users and that more concern be devoted to the underprivileged with whom the major disease burdens will rest in the future [168].

Concluding remarks

Since the early caries lesions that are amenable to non-operative caries treatment are much better detected by means of the visual–tactile examination than by bite-wing radiography [68,169], it would seem timely for the dental profession to place greater reliance on the results of a diligently exercised clinical visual–tactile caries examination and begin phasing out the use of bite-wing radiographs and other ancillary and assumption-ridden diagnostic methods for the detection of 'additional' caries lesions. The visual–tactile caries examination is the centrepiece for the identification of caries lesions that are amenable to treatment based on biologically sound principles, which call for assessment of lesion activity and surface integrity.

The International Caries Detection and Assessment System (ICDAS), which is advertised by some as the new common denominator for modern caries diagnosis in both clinical and research settings [4], is clearly marked by its origin as a tool primarily for researchers and epidemiologists [170]. As such it does not seem to suit the needs of clinicians particularly well. The caries lesion classification employed by ICDAS does not corroborate the script-matching 'this-type-of-lesion-needs-this-kind-of-treatment' type of decision making used by dental clinicians [13], and does not consider lesion-activity assessment. Moreover, to achieve lesion-activity assessment in the ICDAS system, points must be assigned to the caries lesion scores based on assessment of plaque

stagnation characteristics and surface texture, and the total number of points is used to determine the activity status [171,172]. From a strictly procedural point of view this caries-lesion detection system therefore seems to have limited applicability in a dental practice setting, where countless caries-related scripts are matched on a daily basis.

Since appropriate caries treatment selection simply necessitates a distinction between active and arrested, and between cavitated and non-cavitated caries lesions, the criteria employed in the visual-tactile caries examination should focus on two aspects: activity status and cavitation status. In this respect, the Nyvad criteria [78,173] are clearly best suited for use in clinical dental practice settings. They are simple to use and provide a direct link between the best treatment options available and the clinical presentation of the lesion in question. Thereby, these criteria fit well with the script-matching procedure used daily by dental clinicians worldwide to make clinical caries-related treatment decisions.

Much continues to be written about the relative merits of various caries diagnostic approaches in terms of their validity and reliability/reproducibility profiles. However, at the end of the day the best caries diagnostic test is that which results in the best long-term oral health outcomes for the patients. This means that it is not sensible to attempt to separate diagnostic decisions from treatment interventions, because it is the treatments and not the diagnostic procedures that can make our patients fare better than they would have done under different circumstances. Thereby, the 'gold standard' test for caries diagnostic methods is the randomized controlled clinical trial, because this is the only type of investigation that can validly provide the answer to the crucial question: which method of detecting caries lesions for treatment provides the best long-term oral health outcomes for our patients?

Acknowledgements

The author thankfully acknowledges the receipt of the 2009 *Acta Odontologica Scandinavica* award for an excellent contribution to dental research. Professor Bente Nyvad is gratefully thanked for her thoughtful comments on the manuscript.

References

- [1] Pitts NB, Stamm JW. International consensus workshop on caries clinical trials (ICW-CCT)—final consensus statements: agreeing where the evidence leads. *J Dent Res* 2004;83: C125–8.
- [2] Nyvad B, ten Cate JM, Robinson C. Cariology in the 21st century. State of the art and future perspectives. *Caries Res* 2004;38:167–329.
- [3] Horowitz AM. A report on the NIH consensus development conference on diagnosis and management of dental caries throughout life. *J Dent Res* 2004;83:C15–17.
- [4] Pitts N, editor. Detection, assessment, diagnosis and monitoring of caries. Basle, Switzerland: Karger; 2009.
- [5] Frame PS, Sawai R, Bowen WH, Meyerowitz C. Preventive dentistry: Practitioners' recommendations for low-risk patients compared with scientific evidence and practice guidelines. *Am J Prev Med* 2000;18:159–62.
- [6] Beirne PV, Clarkson JE, Worthington HV. Recall intervals for oral health in primary care patients. *Cochrane Database Syst Rev* 2007;(4):CD004346.
- [7] Deep P. Screening for common oral diseases. *J Can Dent Assoc* 2000;66:298–9.
- [8] Pitts NB. Modern concepts of caries measurement. *J Dent Res* 2004;83:C43–7.
- [9] Nyvad B. Diagnosis versus detection of caries. *Caries Res* 2004;38:192–8.
- [10] Baelum V, Heidmann J, Nyvad B. Dental caries paradigms in diagnosis and diagnostic research. *Eur J Oral Sci* 2006;114:263–77.
- [11] Longbottom C, Huysmans M-C, Pitts NB, Fontana M. Glossary of key terms. *Monogr Oral Sci* 2009;21:209–16.
- [12] Pitts NB. Implementation. Improving caries detection, assessment, diagnosis and monitoring. *Monogr Oral Sci* 2009;21:199–208.
- [13] Bader JD, Shugars DA. What do we know about how dentists make caries-related treatment decisions? *Commun Dent Oral Epidemiol* 1997;25:97–103.
- [14] Bader JD, Shugars DA. Variation in clinical decision making related to caries. In: Fejerskov O, Kidd E, editors. *Dental caries. The disease and its clinical management*, 2nd edn. Oxford, UK: Blackwell Munksgaard; 2008. p. 556–74.
- [15] Maupomé G, Sheiham A. Clinical decision-making in restorative dentistry. Content-analysis of diagnostic thinking processes and concurrent concepts used in an educational environment. *Eur J Dent Educ* 2000;4:143–52.
- [16] Nyvad B, Fejerskov O, Baelum V. Visual-tactile caries diagnosis. In: Fejerskov O, Kidd E, editors. *Dental caries. The disease and its clinical management*, 2nd edn. Oxford, UK: Blackwell Munksgaard; 2008. p. 49–68.
- [17] Bader JD, Shugars DA. Need for change in standards of caries diagnosis—epidemiology and health services research perspective. *J Dent Educ* 1993;57:415–21.
- [18] Swan ESC, Lewis DW. Ontario dentists: 1. Radiologic practices and opinions. *J Can Dent Assoc* 1993;59:62–7.
- [19] Frame PS, Sawai R, Bowen WH, Meyerowitz C. Preventive dentistry: practitioners' recommendations for low-risk patients compared with scientific evidence and practice guidelines. *Am J Prev Med* 2000;18:159–62.
- [20] Mileman P, Purdell-Lewis D, van der Weele L. Variation in radiographic caries diagnosis and treatment decisions among university teachers. *Commun Dent Oral Epidemiol* 1982;10:329–34.
- [21] Mileman PA, van den Hout WB. Comparing the accuracy of Dutch dentists and dental students in the radiographic diagnosis of dentinal caries. *Dentomaxillofac Radiol* 2002;31:7–14.
- [22] Rytömaa I, Järvinen V, Järvinen J. Variation in caries recording and restorative treatment plan among university teachers. *Commun Dent Oral Epidemiol* 1979;7:335–9.
- [23] Rosen B, Birkhed D, Nilsson K, Olavi G, Egelberg J. Reproducibility of clinical caries diagnoses on coronal and root surfaces. *Caries Res* 1996;30:1–7.
- [24] Ekstrand KR, Ricketts DNJ, Longbottom C, Pitts NB. Visual and tactile assessment of arrested initial enamel carious lesions: an in vivo pilot study. *Caries Res* 2005;39:173–7.
- [25] Lazarchik DA, Firestone AR, Heaven TJ, Filler SJ, Lussi A. Radiographic evaluation of occlusal caries: effect of training and experience. *Caries Res* 1995;29:355–8.

- [26] Noar SJ, Smith BG. Diagnosis of caries and treatment decisions in approximal surfaces of posterior teeth in vitro. *J Oral Rehabil* 1990;17:209–18.
- [27] Thomas CC, Land MF, Albin-Wilson SM, Stewart GP. Caries detection accuracy by multiple clinicians and techniques. *Gen Dent* 2000;48:334–8.
- [28] Ermis RB, Aydin U. Examiner agreement in the replacement decision of Class I amalgam restorations. *J Contemp Dent Pract* 2004;15:81–92.
- [29] Setcos JC, Khosrave R, Wilson NH, Shen C, Yang M, Mjör IA. Repair or replacement of amalgam restorations: decisions at a USA and a UK dental school. *Oper Dent* 2004;29:392–7.
- [30] Bader JD, Shugars DA. Agreement among dentists' recommendations for restorative treatment. *J Dent Res* 1993;72:891–6.
- [31] Bader JD, Shugars DA. Variation in dentists' clinical decisions. *J Publ Health Dent* 1995;55:181–8.
- [32] Bader JD, Shugars DA. Variation, treatment outcomes, and practice guidelines in dental practice. *J Dent Educ* 1995;59:61–95.
- [33] Espelid I, Tveit AB, Fjelltveit A. Variations among dentists in radiographic detection of occlusal caries. *Caries Res* 1994;28:169–75.
- [34] Espelid I, Tveit AB, Mejare I, Sundberg H, Hallonsten A-L. Restorative treatment decisions on occlusal caries in Scandinavia. *Acta Odontol Scand* 2001;59:21–7.
- [35] Espelid I, Tveit AB. Radiographic diagnosis of mineral loss in approximal enamel. *Caries Res* 1984;18:141–8.
- [36] Espelid I. Radiographic diagnoses and treatment decisions on approximal caries. *Commun Dent Oral Epidemiol* 1986;14:265–70.
- [37] Espelid I, Tveit AB, Riordan PJ. Radiographic caries diagnosis by clinicians in Norway and Western Australia. *Commun Dent Oral Epidemiol* 1994;22:214–19.
- [38] Doméjean-Orliaguet S, Tubert-Jeannin S, Riordan PJ, Espelid I, Tveit AB. French dentists' restorative treatment decisions. *Oral Health Prev Dent* 2004;2:125–31.
- [39] Espelid I, Tveit AB. A comparison of radiographic occlusal and approximal caries diagnoses made by 240 dentists. *Acta Odontol Scand* 2001;59:285–9.
- [40] Hintze H, Wenzel A, Danielsen B, Nyvad B. Reliability of visual examination, fibre-optic transillumination, and bite-wing radiography, and reproducibility of direct visual examination following tooth separation for the identification of cavitated carious lesions in contacting approximal surfaces. *Caries Res* 1998;32:204–9.
- [41] Kay EJ, Nuttall NM, Knill-Jones R. Restorative treatment thresholds and agreement in treatment decision-making. *Commun Dent Oral Epidemiol* 1992;20:265–8.
- [42] Kay EJ, Nuttall NM. Relationship between dentists' treatment attitudes and restorative decisions made on the basis of simulated bite-wing radiographs. *Commun Dent Oral Epidemiol* 1994;22:71–4.
- [43] Kay E, Nuttall N. Clinical decision making—an art or a science? Part I: an introduction. *Br Dent J* 1995;178:76–8.
- [44] Baelum V. Caries management: Technical solutions to biological problems or evidence-based care? *J Oral Rehabil* 2008;35:135–51.
- [45] Bader JD, Ismail AI. A primer on outcomes in dentistry. *J Publ Health Dent* 1999;59:131–5.
- [46] Eckenbarger WH. How honest are dentists? *Readers Digest* 1997; February:50–6.
- [47] Renshaw J. After the first 125 years of the BDJ where might clinical dentistry be heading? *Br Dent J* 2005;199:331–7.
- [48] Berenson A. Boom times for dentists, but not for teeth. Available at http://www.nytimes.com/2007/10/11/business/11decay.html?_r=1. 2007.
- [49] Kidd EAM, Fejerskov O. What constitutes dental caries? Histopathology of carious enamel and dentin related to the action of cariogenic biofilms. *J Dent Res* 2004;83(Special Issue C):C35–8.
- [50] Anderson BG. Clinical study of arresting dental caries. *J Dent Res* 1938;17:443–52.
- [51] Lo ECM, Schwarz E, Wong MCM. Arresting dentine caries in Chinese preschool children. *Int J Paediatr Dent* 1998;8:253–60.
- [52] Nyvad B, Fejerskov O. Active root surface caries converted into inactive caries as a response to oral hygiene. *Scand J Dent Res* 1986;94:281–4.
- [53] Nyvad B, Fejerskov O. Assessing the stage of caries lesion activity on the basis of clinical and microbiological examination. *Commun Dent Oral Epidemiol* 1997;25:69–75.
- [54] Nyvad B, Machiulskiene V, Baelum V. Construct and predictive validity of clinical caries diagnostic criteria assessing lesion activity. *J Dent Res* 2003;82:117–22.
- [55] Wenzel A. Bite-wing and digital bite-wing radiography for detection of caries lesions. *J Dent Res* 2004;83:C72–5.
- [56] de Araujo FB, Rosito DB, Toigo E, dos Santos CK. Diagnosis of approximal caries: radiographic versus clinical examination using tooth separation. *Am J Dent* 1992;5:245–8.
- [57] de Araujo FB, de Araujo DR, dos Santos CK, de Souza MA. Diagnosis of approximal caries in primary teeth: radiographic versus clinical examination using tooth separation. *Am J Dent* 1996;9:54–6.
- [58] Thylstrup A, Bille J, Qvist V. Radiographic and observed tissue changes in approximal carious lesions at the time of operative treatment. *Caries Res* 1986;20:75–84.
- [59] Bille J, Thylstrup A. Radiographic diagnosis and clinical tissue changes in relation to treatment of approximal carious lesions. *Caries Res* 1982;16:1–6.
- [60] Akpata ES, Farid MR, Al-Saif K, Roberts EAU. Cavitation at radiolucent areas on proximal surfaces of posterior teeth. *Caries Res* 1996;30:313–6.
- [61] Mejare I, Malmgren B. Clinical and radiographic appearance of proximal carious lesions at the time of operative treatment in young permanent teeth. *Scand J Dent Res* 1986;94:19–26.
- [62] Pitts NB, Rimmer PA. An in vivo comparison of radiographic and directly assessed clinical caries status of posterior approximal surfaces in primary and permanent teeth. *Caries Res* 1992;26:146–52.
- [63] Ratledge DK, Kidd EAM, Beighton D. A clinical and microbiological study of approximal carious lesions. Part 1: The relationship between cavitation, radiographic lesion depth, the site-specific gingival index and the level of infection of the dentine. *Caries Res* 2001;35:3–7.
- [64] Mariath AAS, Casagrande L, de Araujo FB. Grey levels and radiolucent lesion depth as cavity predictors for approximal dentin caries lesions in primary teeth. *Dentomaxillofac Radiol* 2007;36:377–81.
- [65] Woodward GL, Leake JL. The use of dental radiographs to estimate the probability of cavitation of carious interproximal lesions. Part I: evidence from the literature. *J Can Dent Assoc* 1996;62:731–6.
- [66] Mejare I, Grondahl HG, Carlstedt K, Grever AC, Ottoson E. Accuracy at radiography and probing for the diagnosis of proximal caries. *Scand J Dent Res* 1985;93:178–84.
- [67] Machiulskiene V, Nyvad B, Baelum V. A comparison of clinical and radiographic caries diagnoses in posterior teeth of 12-year-old Lithuanian children. *Caries Res* 1999;33:340–8.
- [68] Machiulskiene V, Nyvad B, Baelum V. Comparison of diagnostic yields of clinical and radiographic caries examinations in children of different age. *Eur J Paediatr Dent* 2004;5:157–62.
- [69] Rimmer PA, Pitts NB. Temporary elective tooth separation as a diagnostic aid in general dental practice. *Br Dent J* 1990;169:87–92.

- [70] Hintze H, Wenzel A, Jones C. In vitro comparison of D- and E-speed film radiography, RVG, and Visualix digital radiography for the detection of enamel approximal and dentinal occlusal caries lesions. *Caries Res* 1994;28:363–7.
- [71] Bader JD, Shugars DA, Bonito AJ. Systematic reviews of selected caries diagnostic and management methods. *J Dent Educ* 2001;65:960–8.
- [72] Bader JD, Shugars DA. A systematic review of the performance of a laser fluorescence device for detecting caries. *J Am Dent Assoc* 2004;135:1413–26.
- [73] Baelum V, Nyvad B, Gröndahl H-G, Fejerskov O. The foundations of good diagnostic practice. In: Fejerskov O, Kidd E, editors. *Dental caries. The disease and its clinical management*, 2nd edn. Oxford, UK: Blackwell Munksgaard; 2008. p. 104–20.
- [74] Kidd EAM, Pitts NB. A reappraisal of the value of the bitewing radiograph in the diagnosis of posterior approximal caries. *Br Dent J* 1990;169:195–200.
- [75] Bader JD, Shugars DA, Bonito AJ. A systematic review of selected caries prevention and management methods. *Commun Dent Oral Epidemiol* 2001;29:399–411.
- [76] Bloemendal E, de Wet HCW, Bouter LM. The value of bitewing radiographs in epidemiological caries research: a systematic review of the literature. *J Dent* 2004;32:255–64.
- [77] Pereira AC, Eggertsson H, Martinez-Mier EA, Mialhe FL, Eckert GJ, Zero DT. Validity of caries detection on occlusal surfaces and treatment decisions based on results from multiple caries-detection methods. *Eur J Oral Sci* 2009;117:51–7.
- [78] Nyvad B, Machiulskiene V, Baelum V. Reliability of a new caries diagnostic system differentiating between active and inactive caries lesions. *Caries Res* 1999;33:252–60.
- [79] Murphy EA. *The logic of medicine*, 2nd edn. Baltimore, MD: Johns Hopkins University Press; 1997.
- [80] Hugoson A, Koch G, Göthberg C, Helkimo AN, Lundin S-Å, Norderyd O, et al. Oral health of individuals aged 3–80 years in Jönköping, Sweden during 30 years (1973–2003). I. Review of findings of dental care habits and knowledge of oral health. *Swed Dent J* 2005;29:125–38.
- [81] Afonso-Souza G, Nadanovsky P, Chor D, Faerstein E, Werneck GL, Lopes CS. Association between routine visits for dental checkup and self-perceived oral health in an adults population in Rio de Janeiro: the Pró-Saúde Study. *Commun Dent Oral Epidemiol* 2007;35:393–400.
- [82] Woolfolk MW, Lang WP, Borgnakke WS, Taylor GW, Ronis DL, Nyquist LV. Determining dental checkup frequency. *J Am Dent Assoc* 1999;130:715–23.
- [83] Sohn W, Ismail AI. Regular dental visits and anxiety in an adult dentate population. *J Am Dent Assoc* 2005;136:58–66.
- [84] Nuttall NM, Bradnock G, White D, Morris J, Nunn J. Dental attendance in 1998 and implications for the future. *Br Dent J* 2001;190:177–82.
- [85] Christensen LB, Petersen PE, Steding-Jessen M. Consumption of dental services among adults in Denmark 1994–2003. *Eur J Oral Sci* 2007;115:174–9.
- [86] Christensen LB, Kjoller M, Petersen PE, Vigild M. Tandstatus og udnyttelsen af tandplejetilbuddet hos voksne i Danmark 1994. Status og udviklingstendenser. *Tandlægebladet* 1996;100:215–22.
- [87] Schwarz E. Changes in demand for dental care among Danish adults, 1975–1990. *Acta Odontol Scand* 1996;54:36–43.
- [88] Suominen-Taipale AL, Widstrom E, Alanen P, Uutela A. Trends in self-reported use of dental services among Finnish adults during two decades. *Commun Dent Health* 2000;17:31–7.
- [89] Sheiham A. Routine check-ups. *Br Dent J* 2000;189:181.
- [90] Last JMA. *dictionary of epidemiology*, 4th edn. Oxford, UK: Oxford University Press; 2001.
- [91] Davenport CF, Elley KM, Fry-Smith A, Taylor-Weetman CL, Taylor RS. The effectiveness of routine dental checks: a systematic review of the evidence base. *Br Dent J* 2003;195:87–98.
- [92] Davenport C, Elley K, Salas C, Taylor-Weetman CL, Fry-Smith A, Bryan S, et al. The clinical effectiveness and cost-effectiveness of routine dental checks: a systematic review and economic evaluation. *Health Technol Assess* 2003;7:No. 7.
- [93] Beirne P, Forgie A, Worthington HV, Clarkson J. Routine scale and polish for periodontal health in adults. *Cochrane Database Syst Rev* 2005;(1):CD004625.
- [94] Gilbert GH, Foerster U, Dolan TA, Duncan RP, Ringelberg ML. Twenty-four month coronal caries incidence: the role of dental care and race. *Caries Res* 2000;34:367–79.
- [95] Bullock C, Boath E, Lewis M, Gardam K, Croft P. A case-control study of differences between regular and casual adult attenders in general dental practice. *Prim Dent Care* 2001;8:35–40.
- [96] Gøtzsche PC, Nielsen M. Screening for breast cancer with mammography. *Cochrane Database Syst Rev* 2009;(4):CD001877.
- [97] Gøtzsche PC, Hartling OJ, Nielsen M, Brodersen J, Jørgensen KJ. Breast screening: the facts—or maybe not. *Br Med J* 2009;338:446–8.
- [98] Jørgensen KJ, Brodersen J, Hartling OJ, Nielsen M, Gøtzsche PC. Informed choice requires information about both benefits and harms. *J Med Ethics* 2009;35:268–9.
- [99] Brodersen J. Measuring psychological consequences of false-positive screening results—breast cancer as an example. Copenhagen, Denmark: Faculty of Health Sciences, University of Copenhagen; 2006.
- [100] Brewer NT, Salz T, Lillie SE. Systematic review: the long-term effects of false-positive mammograms. *Ann Intern Med* 2007;146:502–10.
- [101] Elderton RJ. Iatrogenesis in the treatment of dental caries. *Proc Finn Dent Soc* 1992;88:25–32.
- [102] Elderton RJ. Overtreatment with restorative dentistry: when to intervene? *Int Dent J* 1993;43:17–24.
- [103] Elderton RJ. Preventive (evidence-based) approach to quality general dental care. *Med Princ Pract* 2003;12(Suppl 1):12–21.
- [104] Welie JVM. Is dentistry a profession? Part 3. Future challenges. *J Can Dent Assoc* 2004;70:675–8.
- [105] Welie JVM. Is dentistry a profession? Part 2. The hallmarks of professionalism. *J Can Dent Assoc* 2004;70:599–602.
- [106] Grytten J. The effect of supplier inducement on Norwegian dental services; some empirical findings based on a theoretical model. *Commun Dent Health* 1991;8:221–31.
- [107] Grytten J. The effect of the price of dental services on their demand and utilisation in Norway. *Commun Dent Health* 1990;8:303–10.
- [108] Grytten J. Supplier inducement—its relative effect on demand and utilization. *Commun Dent Oral Epidemiol* 1992;20:6–9.
- [109] Birch S. The identification of supplier-inducement in a fixed price system of health care provision. The case of dentistry in the United Kingdom. *J Health Econ* 1988;7:129–50.
- [110] Grembowski D, Fiset L, Milgrom P, Conrad D, Spadafora A. Does fluoridation reduce the use of dental services among adults? *Med Care* 1997;35:454–71.
- [111] Welie JVM. Is dentistry a profession? Part 1. Professionalism defined. *J Can Dent Assoc* 2004;70:529–32.
- [112] Cruess SR, Johnston S, Cruess RL. “Profession”: a working definition for medical educators. *Teach Learn Med* 2004;16:74–6.
- [113] Cruess RL, Cruess SR, Johnston SE. Professionalism: an ideal to be sustained. *Lancet* 2000;356:156–9.
- [114] Cruess SR, Johnston S, Cruess RL. Professionalism for medicine: opportunities and obligations. *Med J Aust* 2002;177:208–11.

- [115] Reid KI. Respect for patients' autonomy. *J Am Dent Assoc* 2009;140:470–4.
- [116] Hafferty F. The elephant in medical professionalism's kitchen. *Acad Med* 2006;81:906–14.
- [117] Freidson E. Professionalism—the third logic. Cambridge, UK: Polity Press; 2001.
- [118] Durkheim E. Professional ethics and civic morals. New York: Routledge; 1957.
- [119] Sundhedsstyrelsen. Børne- og ungdomstandplejen i Danmark 1988. Primær sundhedstjenestestatistik III 1989;11:7–77.
- [120] Downer MC. Time trends in caries experience of children in England and Wales. *Caries Res* 1992;26:466–72.
- [121] Truin GJ, König KG, Kalsbeek H. Trends in dental caries in The Netherlands. *Adv Dent Res* 1993;7:15–18.
- [122] Marthaler TM, O'Mullane DM, Vrbic V. The prevalence of dental caries in Europe 1990–1995. *Caries Res* 1996;30:237–55.
- [123] Poulsen S. Dental caries in Danish children and adolescents 1988–94. *Commun Dent Oral Epidemiol* 1996;24:282–5.
- [124] Einarsdottir KG, Bratthall D. Restoring oral health. On the rise and fall of dental caries in Iceland. *Eur J Oral Sci* 1996;104:459–69.
- [125] Davies MJ, Spencer AJ, Slade GD. Trends in dental caries experience of school children in Australia—1977 to 1993. *Aust Dent J* 1997;42:389–94.
- [126] Kalsbeek H, Truin GJ, van Rossum GMJM, van Rijkom HM, Poorterman JHG, Verrips GH. Trends in caries prevalence in Dutch adults between 1983 and 1995. *Caries Res* 1998;32:160–5.
- [127] de Liefde B. The decline of caries in New Zealand over the past 40 years. *N Z Dent J* 1998;94:109–13.
- [128] Cleaton-Jones P, Fatti P. Dental caries trends in Africa. *Commun Dent Oral Epidemiol* 1999;27:316–20.
- [129] Poulsen S, Scheutz F. Dental caries in Danish children and adolescents 1988–1997. *Commun Dent Health* 1999;16:166–70.
- [130] Beltrán-Aguilar ED, Estupiñán-Day S, Báez R. Analysis of prevalence and trends of dental caries in the Americas between the 1970s and 1990s. *Int Dent J* 1999;49:322–9.
- [131] Künzel W, Fischer T. Caries prevalence after cessation of water fluoridation in La Salud, Cuba. *Caries Res* 2000;34:20–5.
- [132] Künzel W, Fischer T, Lorenz R, Brühmann S. Decline of caries prevalence after the cessation of water fluoridation in the former East Germany. *Commun Dent Oral Epidemiol* 2000;28:382–9.
- [133] Brown LJ, Wall TP, Lazar V. Trends in total caries experience: permanent and primary teeth. *J Am Dent Assoc* 2000;131:223–31.
- [134] Birkeland JM, Haugejorden O, von der Fehr FR. Some factors associated with the caries decline among Norwegian children and adolescents: age-specific and cohort analyses. *Caries Res* 2000;34:109–16.
- [135] Vrbic V. Reasons for the caries decline in Slovenia. *Commun Dent Oral Epidemiol* 2000;28:126–32.
- [136] Szöke J, Petersen PE. Evidence for dental caries decline among children in an East European country (Hungary). *Commun Dent Oral Epidemiol* 2000;28:155–60.
- [137] Virtanen JI. Changes and trends in attack distributions and progression of dental caries in three age cohorts in Finland. *J Epidemiol Biostat* 2001;6:325–9.
- [138] Vanobbergen J, Martens L, Declerck D. Caries prevalence in Belgian children: a review. *Int J Paediatr Dent* 2001;11:164–70.
- [139] Maltz M, Schoenardie AB, Carvalho JC. Dental caries and gingivitis in schoolchildren from the municipality of Porto Alegre, Brazil in 1975 and 1996. *Clin Oral Invest* 2001;5:199–204.
- [140] Birkeland JM, Haugejorden O. Caries decline before fluoride toothpaste was available: earlier and greater decline in the rural north than in southwestern Norway. *Acta Odontol Scand* 2001;59:7–13.
- [141] Kovac-Kavcic M, Skaleric U. The change of DMFT counts in Slovenia. *Caries Res* 2001;35:247–51.
- [142] Brown LJ, Wall TP, Lazar V. Trends in caries among adults 18 to 45 years old. *J Am Dent Assoc* 2002;133:827–34.
- [143] Birkeland JM, Haugejorden O, von der Fehr FR. Analyses of the caries decline and incidence among Norwegian adolescents 1985–2000. *Acta Odontol Scand* 2002;60:281–9.
- [144] Van Nieuwenhuysen JP, Carvalho JC, D'Hoore W. Caries reduction in Belgian 12-year-old children related to socio-economic status. *Acta Odontol Scand* 2002;60:123–8.
- [145] Poulsen S, Pedersen MM. Dental caries in Danish children: 1988–2001. *Eur J Paediatr Dent* 2002;4:195–8.
- [146] Bönecker M, Cleaton-Jones P. Trends in dental caries in Latin American and Caribbean 5-6- and 11-13-year-old children: a systematic review. *Commun Dent Oral Epidemiol* 2003;31:152–7.
- [147] Petersen PE. The World Oral Health Report 2003. Geneva, Switzerland: World Health Organization; 2003.
- [148] Marthaler TM. Changes in dental caries 1953–2003. *Caries Res* 2004;38:173–81.
- [149] Aleksejuniene J, Holst D, Balciuniene I. Factors influencing the caries decline in Lithuanian adolescents—trends in the period 1993–2001. *Eur J Oral Sci* 2004;112:3–7.
- [150] Carvalho JC, D'Hoore W, Van Nieuwenhuysen JP. Caries decline in the primary dentition of Belgian children over 15 years. *Commun Dent Oral Epidemiol* 2004;32:277–82.
- [151] Pieper K, Schulte AG. The decline in dental caries among 12-year-old children in Germany between 1994 and 2000. *Commun Dent Health* 2004;21:199–206.
- [152] Bourgeois DM, Roland E, Desfontaine J. Caries prevalence 1987–1998 in 12-year-olds in France. *Int Dent J* 2004;54:193–200.
- [153] Stecksen-Blicks C, Sunnegårdh K, Borssen E. Caries experience and background factors in 4-year-old children: time trends 1967–2002. *Caries Res* 2004;38:149–55.
- [154] Beltrán-Aguilar ED, Barker LK, Canto MT, Dye BA, Gooch BF, Griffin SO, et al. Surveillance for dental caries, dental sealants, tooth retention, edentulism, and enamel fluorosis—United States 1988–1994 and 1999–2002. *MMWR Morb Mortal Wkly Rep* 2005;54(SS-3):1–48.
- [155] Adam C, Eid A, Riordan PJ, Wolikow M, Cohen F. Caries experience in the primary dentition among French 6-year-olds between 1991 and 2000. *Commun Dent Oral Epidemiol* 2005;33:333–40.
- [156] Marthaler T, Menghini G, Steiner M. Use of the Significant Caries Index in quantifying the changes in caries in Switzerland from 1964 to 2000. *Commun Dent Oral Epidemiol* 2005;33:159–66.
- [157] Cleaton-Jones P, Fatti P, Bönecker M. Dental caries trends in 5- to 6-year-old and 11- to 13-year-old children in three UNICEF designated regions—Sub Saharan Africa, Middle East and North Africa, Latin America and Caribbean: 1970–2004. *Int Dent J* 2006;56:294–300.
- [158] Demertzi A, Topitsoglou V, Muronidis S. Caries prevalence of 11.5 year-olds between 1989 and 2001 in a province of North-Eastern Greece. *Commun Dent Health* 2006;23:140–6.
- [159] Souza ML, Bastos JL, Peres MA. Trends in dental caries rates in 12- and 13-year-old school children from Florianópolis (Brazil) between 1971 and 2005. *Oral Health Prev Dent* 2006;4:187–92.
- [160] Ferro R, Besostri A, Olivieri A, Stellini E, Mazzoleni S. Preschoolers' dental caries experience and its trend over 20 years in a North-East Italian Health District. *Eur J Paediatr Dent* 2007;8:199–204.

- [161] Hugoson A, Koch G, Helkimo AN, Lundin S-Å. Caries prevalence and distribution in individuals aged 3-20 years in Jönköping, Sweden, over a 30-year period (1973-2003). *Int J Paediatr Dent* 2008;18:18-26.
- [162] Sgan-Cohen HD, Amram-Liani D, Livny A. Changing dental caries levels in the 1980's, 1990's and 2005 among children of a Jerusalem region. *Commun Dent Health* 2009;26:62-4.
- [163] Irigoyen Camacho ME, Sánchez Pérez L, García Pérez A, Zepeda Zepeda MA. Relationship between severe early childhood caries, mother's oral health and mutans streptococci in a low-income group: changes from 1996 to 2007. *J Clin Pediatr Dent* 2009;33:241-6.
- [164] Carvalho JC, Figueiredo MJ, Vieira EO, Mestrinho HD. Caries trends in Brazilian non-privileged preschool children in 1996 and 2006. *Caries Res* 2009;43:2-9.
- [165] Hugoson A, Koch G, Göthberg C, Helkimo AN, Lundin S-Å, Norderyd O, et al. Oral health of individuals aged 3-80 years in Jönköping, Sweden during 30 years (1973-2003). II. Review of clinical and radiographic findings. *Swed Dent J* 2005;29:139-55.
- [166] Gimmestad AL, Holst D, Fylkenes K. Changes in restorative caries treatment in 15-year-olds in Oslo, Norway, 1979-1996. *Commun Dent Oral Epidemiol* 2003;31:246-51.
- [167] Heidmann J, Helm S, Helm T, Poulsen S. Changes in prevalence of approximal caries in 17-year-olds and related restorative treatment strategies over a 6-year period. *Commun Dent Oral Epidemiol* 1988;16:167-70.
- [168] Baelum V, van Palenstein Helderman W, Hugoson A, Yee R, Fejerskov O. 'For richer, for poorer, in sickness and in health...' The role of dentistry in controlling caries and periodontitis globally. In: Fejerskov O, Kidd E, editors. *Dental caries. The disease and its clinical management*, 2nd edn. Oxford, UK: Blackwell Munksgaard; 2008. p. 575-605.
- [169] Machiulskiene V, Nyvad B, Baelum VA. A comparison of clinical and radiographic caries diagnoses in posterior teeth of 12-year-old Lithuanian children. *Caries Res* 1999;33:340-8.
- [170] Topping GVA, Pitts NB. Clinical visual caries detection. *Monogr Oral Sci* 2009;21:15-41.
- [171] Ekstrand KR, Zero DT, Martignon S, Pitts NB. Lesion activity assessment. *Monogr Oral Sci* 2009;21:63-90.
- [172] Braga MM, Mendes FM, Martignon S, Ricketts DNJ, Ekstrand KR. In vitro comparison of Nyvad's system and ICDAS-II with lesion activity assessment for evaluation of severity and activity of occlusal caries lesions in primary teeth. *Caries Res* 2009;43:405-12.
- [173] Nyvad B, Machiulskiene V, Baelum V. The Nyvad criteria for assessment of caries lesion activity. In: Stookey G, editor. *Clinical models workshop: Remin-demin, precavitation, caries. Proceedings of the 7th Indiana Conference*. Indianapolis, IN: Indiana University School of Dentistry; 2006. p. 99-116.