

A Case for Sequencing the Genome of *Musca domestica* (Diptera: Muscidae)

J. G. SCOTT,¹ N. LIU,² M. KRISTENSEN,³ AND A. G. CLARK⁴

J. Med. Entomol. 46(2): 175–182 (2009)

ABSTRACT House flies are carriers of >100 devastating diseases that have severe consequences for human and animal health. Despite the fact that it is a passive vector, a key bottleneck to progress in controlling the human diseases transmitted by house flies is lack of knowledge of the basic molecular biology of this species. Sequencing of the house fly genome will provide important inroads to the discovery of novel target sites for house fly control, understanding of the house fly immune response, rapid elucidation of insecticide resistance genes, and understanding of numerous aspects of the basic biology of this insect pest. The ability of the house fly to prosper in a remarkably septic environment motivates analysis of its innate immune system. Its polymorphic sex determination system, with male-determining factors on either the autosomes or the Y chromosome, is ripe for a genomic analysis. Sequencing of the house fly genome would allow the first opportunity to study the interactions between a pest insect and its parasitoid (*Nasonia vitripennis*) at the whole genome level. In addition, the house fly is well placed phylogenetically to leverage analysis of the multiple Dipteran genomes that have been sequenced (including several mosquito and *Drosophila* species). The community of researchers investigating *Musca domestica* are well prepared and highly motivated to apply genomic analyses to their widely varied research programs.

KEY WORDS genome sequencing, insect immunity, insecticide resistance, sex determination, comparative genomics

House Flies

Biology and Importance. House flies, *Musca domestica* L. (Diptera: Muscidae), are cosmopolitan, ubiquitous, and transmit >100 human and animal diseases (Scott and Lettig 1962, Greenberg 1965, Keiding 1986), including bacterial infections such as salmonellosis, anthrax, ophthalmia, shigellosis, typhoid fever, tuberculosis, cholera, and infantile diarrhea; protozoan infections such as amebic dysentery; helminthic infections such as pinworms, roundworms, hookworms, and tapeworms; and viral and rickettsial infections. Recently house flies were shown to spread a deadly strain of *Escherichia coli* in Japan (Sasaki et al. 2000). Flies also transmit pathogens responsible for eye diseases such as trachoma and epidemic conjunctivitis and infect wounds or skin with diseases such as cutaneous diphtheria, mycoses, yaws, and leprosy (Keiding 1986). Fly-transmitted trachoma alone causes six million cases of childhood blindness each year (World Health Organization 2004). Considering that house

flies are highly mobile, come into contact with excreta, carcasses, garbage, and other septic matter, and that they are intimately associated with humans, our food, and utensils, it is not surprising that they are involved in transmission of so many serious and widespread diseases (Scott and Lettig 1962, Keiding 1986). Most recently, house flies have been shown to transmit life-threatening antibiotic-resistant bacteria (Rahuma et al. 2005, Macovei and Zurek 2006), which are an ever increasing problem in hospitals and other health care facilities (Sundin 1996, Graczyk et al. 2001, Maisnier-Patin and Andersson 2004, Boulesteix et al. 2005).

House flies are always found in association with humans and human activities. In fact, house flies and humans have evolved together, with house flies following the spread of *Homo sapiens* across the planet (Münder 1994). House flies are also one of the most serious pests at dairy, horse, hog, sheep, and poultry facilities worldwide. Exposure to debilitating disease-causing agents, public health and nuisance concerns, lowered levels of milk and egg production, and reduced feed conversion all result from house fly activity. Economic losses and the cost associated with fly suppression are difficult to quantify, but costs of pesticides for fly control at poultry facilities alone are estimated at more than \$200 million annually in the United States (Geden et al. 1994). With the loss of available insecticides (caused by governmental can-

¹ Department of Entomology, Comstock Hall, Cornell University, Ithaca, NY 14853 (e-mail: jgs5@cornell.edu).

² Department of Entomology and Plant Pathology, Auburn University, Auburn, AL 36849.

³ Danish Pest Infestation Laboratory, Department of Integrated Pest Management, Faculty of Agricultural Sciences, University of Aarhus, DK-2800 Lyngby, Denmark.

⁴ Department of Molecular Biology and Genetics, Comstock Hall, Cornell University, Ithaca, NY 14853.

cellations, the high cost of reregistration in the absence of patent protection, and the development of insecticide resistance), there is a pressing need for the development of new insecticides and control strategies.

Many species of arthropods are the sources of potent allergens that sensitize and induce IgE-mediated allergic reactions in humans. Most of these arthropod allergens are proteins, and the allergic response mechanism is the same as those from other sources such as plant pollens, molds, and foods. Allergies to house flies are rare, but cases of respiratory allergy from occupational exposure (farmers) have been reported (Wahl and Fraedrich 1997, Focke et al. 2003). Identification of house fly allergens could lead to recombinant allergens with a potential use in diagnosis and immunotherapy.

Given the importance of house flies in the transmission of human and animal diseases, there has been substantial effort to control fly populations, primarily with insecticides. Generally, house fly control has involved DDT and methoxychlor, as well as other chlorinated hydrocarbons (e.g., lindane, and chlordane), organophosphates (e.g., malathion, diazinon, and dimethoate), carbamates (e.g., methomyl), pyrethrins (usually with piperonyl butoxide), pyrethroids (e.g., permethrin, fenvalerate, and cyfluthrin), and most recently spinosad (limited use) and neonicotinoid baits (e.g., imidacloprid). House flies have shown a remarkable ability to rapidly evolve resistance to each of the insecticides used against it (Keiding 1999). Identification of the gene involved in a specific resistance is hampered by the lack of a genome sequence. House fly control is also practiced using biological control agents such as pteromalid wasps (Axtell 1990). The genome for one of these important biological control agents, *Nasonia vitripennis*, has recently been completed.

Suitability as a Study Organism. The house fly has many advantages as an experimental organism. It is easy to rear on standard media, and thousands of house flies can be produced in a matter of weeks. Under normal laboratory conditions, it takes ≈ 10 d to develop from egg to adult.

Insect transgenesis is critically important for both practical applications and for addressing basic scientific questions. The generation of transgenic lines of insects has proven to be perhaps the most powerful method for showing the functional role of genes, both by overexpression studies and by mutation-rescue studies. Transgenic insects have resulted in significant progress in understanding the genes involved in disease transmission and in understanding the biological and physiological roles of numerous genes. Germ line transformation of house flies has been successfully carried out by various methods (Atkinson et al. 1993, O'Brochta et al. 1994, Warren et al. 1994, Hediger et al. 2001). More recently, one of the authors (N. Liu, Auburn University) has carried out *Musca* transformations to study the genes associated with insecticide resistance. Similar studies will also facilitate the identification of new target sites that

could lead to the development of novel insecticides with new modes of action and low toxicity to non-target species. The availability of transgenic technology and completion of the house fly genome will open numerous areas of study that were previously not approachable.

House Fly Genetics. *Musca domestica* has a well-described linkage map for the five autosomes and two sex chromosomes (X and Y) (Hiroyoshi 1960, Tsukamoto et al. 1961, Milani et al. 1967, Nickel and Wagoner 1974, Hiroyoshi 1977). Crossing over is very rare in male house flies (Hamm et al. 2005), a feature that has been used to advantage in genetic analysis of *Musca*, just as it has for *Drosophila*. Dozens of genetically defined house fly strains are available globally for identification of genes responsible for various traits, as well as for identification of polymorphisms. House flies do not suffer from severe inbreeding depression (Reed and Bryant 2004), and many highly inbred strains (which are preferred for genome sequencing projects) are available.

In the house fly, sex is determined by a dominant factor, M, which is located on the Y chromosome in "standard" populations. Thus, males are XY^M and females are XX (Hiroyoshi 1964, Dübendorfer et al. 2002). This is believed to be the ancestral state of sex determination in house flies (Bull and Charnov 1977, Denholm et al. 1983). However, there are "autosomal male" (A^M) strains in which the M factor is located on one or more of the five autosomes (I-V) (Franco et al. 1982, Inoue et al. 1983, Tomita and Wada 1989, Hamm et al. 2005, Hamm and Scott 2009, Kozielska et al. 2008) or even rarely on X (Schmidt et al. 1997). The M factor located on Y functions biologically in a way identical to the M located on any of the other autosomes (Tomita and Wada 1989, Schmidt et al. 1997). The sequence of M is unknown. In the A^M strains, females are XX and males are also XX (or XO) (Hiroyoshi 1964; Wagoner 1969; Franco et al. 1982; Denholm et al. 1983, 1990). Populations are found in which males are A^M/A^M (Tomita and Wada 1989, Hamm and Scott 2009). Such populations have females with F (feminizing factor located on autosome 4), which is epistatic to M, as a means to produce female offspring. In these populations, females have become the heterogametic sex. F has recently been sequenced (D. Bopp, personal communication).

Quantitative real-time polymerase chain reaction (qRT-PCR) has been shown to be a reliable method for determination of genome size (Wilhelm et al. 2003), and the genome sizes of *D. melanogaster* and *M. domestica* were recently compared using this method. The size of the *D. melanogaster* genome was found to be 180–181 Mbp (in agreement with the published genome size; Adams et al. 2000), and the size of the *M. domestica* genome was found to be 309–312 Mbp (Gao and Scott 2006) or ≈ 1.7 -fold larger than *D. melanogaster*. This indicates that the size of the house fly genome makes it an excellent candidate for whole genome sequencing.

Rationale for Sequencing of the House Fly Genome

Expanding Our Understanding of Basic House Fly Biology to Develop New Control Strategies. Given the tremendous importance of house flies in the transmission of human and animal diseases, substantial effort has been made to control this pest. Availability of the house fly genome will allow for identification of important target sites and will allow for the development of selective new insect control agents. Identification of novel target sites in the house fly will also aid in the development of new insecticides for control of house flies, as well as of agricultural pests that limit the supply (and quality) of human foods. A genome sequence would also provide an opportunity to explore biological control in novel ways, including disruption of the unusual autosome-based sex determination system, sterile male release, confounding signals for mate recognition, etc. Such approaches may be safer than insecticides, given the proximity of house flies to humans, animals, and many of their important food sources, and would offer important alternative control measures for organic farmers.

The biochemistry and genetics of insecticide resistance have been well studied in the house fly, arguably more widely than in any other insect. This is because of the medical and economic importance of house flies, that they are direct targets of insecticide control, the relatively rapid rate at which they develop resistance, the availability of strains resistant to almost every class of insecticide, the well-understood biology of the house fly, and that the house fly has proven to be a useful model for understanding and predicting resistance in other insect species (Scott 1990, 1991, 1999). Availability of the house fly genome would allow for more rapid identification of the genes and regulatory sequences involved in resistance to insect control agents.

The house fly has been, and continues to be, a major insect for studies of environmental toxicants. It has had a preeminent role in insect toxicology studies, especially with focus on comparative toxicity between insects and mammals, in the development of new insecticides (Casida and Quistad 2004) and insecticide resistance (see above). The house fly has been a model insect for these scientific areas of inquiry. Completion of the house fly genome will facilitate this research by the identification of novel target sites, further elucidation of differences in target sites (ion channels, neuroreceptors, hormones, etc.) between insects and mammals, and by facilitating identification of genes involved in insecticide resistance. The neonicotinoids serve as an excellent example of the payoff that comes from this comparative biochemical approach. Neonicotinoids are the fastest growing class of insecticides and were developed specifically by the process of selecting agents that interact with insect and not mammalian receptors (Matsuda et al. 2001, Nauen et al. 2003, Tomizawa and Casida 2003, Wakita et al. 2003). Studies of house fly nicotinic acetylcholine receptor subunits have recently identified novel RNA editing sites (Gao et al. 2007a, b, c), but lack of a genome

sequence hampers efforts to understand the scope of RNA editing in this species.

Despite efforts by developmental biologists, the molecular identity of M has remained elusive. In *Drosophila*, *Sex-lethal* (*Sxl*) integrates information about the dose of X and autosomes and provides the initial switch for the sex determination cascade. In *M. domestica*, M is not a signal for *Sxl*, and in fact, *Sxl* is not involved in sex determination (Meise et al. 1998). Most intriguingly, in *Drosophila*, sex determination and dosage compensation are tied to the same pathway, whereas these processes are decoupled in *Musca*. It has been speculated that it is this decoupling that gives *Musca* the impressive flexibility and polymorphism in sex determination mechanisms (Dübendorfer et al. 2002, Hediger et al. 2004). A full genome sequence of *M. domestica* would allow immediate identification of all homologs to the *Drosophila* sex determination cascade and would greatly accelerate discovery of the genes that cause the radical divergence in the fundamental processes of sex determination found in most Diptera compared with dosage compensation found in *Drosophila* (Dübendorfer et al. 2002, Hediger et al. 2004).

The house fly has been a model system for studies of insect olfaction (Kelling et al. 2002, 2003), and (Z)-9-tricosene plays an important role in intersex communication and mate selection in house flies. Sequencing of the house fly genome will identify receptor molecules (in antennal and palpal olfactory cells) that will aid olfaction studies, and will facilitate development of attractants for house flies to baits in management systems (Darbro and Mullens 2004, Hanley et al. 2004).

Improved Understanding of Insect Immunity. The house fly thrives in a virtual sea of animal pathogens. Sequencing of the house fly genome will shed light on the immune defense systems of this important species and provide valuable information about how it is able to flourish, despite living in intimate contact with septic flora and fauna. Comparison between the innate immune systems of *Musca*, *Drosophila*, and *Anopheles*, which face different ecological pressures and pathogens, will be informative, just as the *Drosophila*-*Anopheles* comparison has been (Christophides et al. 2002). The relatively close relationship to *Drosophila* has already greatly expedited this analysis, because >30 individual innate immunity genes have been sequenced in *Musca*. The advantage of a genome sequence is that it will allow discovery of genes unique to *Musca* and regulatory systems that allow it to survive in a far more septic environment.

Completion of the *Drosophila* and *Anopheles* genomes provided unprecedented opportunities to study insect-pathogen interactions (Christophides et al. 2002, Lazzaro and Clark 2003, Schlenke and Begun 2003, Lazzaro et al. 2004, Osta et al. 2004, Srinivasan et al. 2004). The house fly will also be of great value for two reasons. First, house flies live in intimate association with vertebrate pathogens such as *Helicobacter pylori* (causative agent of gastric ulcer; Li and Stutzenberger 2000), *Salmonella*, *Campylobacter jejuni*

(Hald et al. 2004), *E. coli* (including toxin producing strains E105 and O157:H7 that cause food poisoning; Moriya et al. 1999, de Jesus et al. 2004), and trachoma (i.e., transmission of *Chlamydia trachomatis*; Emerson et al. 1999, 2000). However, house flies are remarkably resilient to pathogens. Understanding the basis for their refractoriness to many pathogens would offer important insights into ways to improve human health. Second, house fly populations in temperate climates are occasionally decimated by Entomophthora, an entomopathogenic fungus (Zygomycetes, Entomophthoraceae). A genome sequence would expedite the study of why certain populations of *Musca* are sensitive to this fungus, whereas others are refractory. Microarray studies using the house fly genome to investigate genes associated with pathogen exposure will be a cornerstone in future studies in this field (Jensen et al. 2001, Kalsbeek et al. 2001, Zurek et al. 2002) and would become a model system for biological control (entomopathogenic fungi; parasitic hymenoptera; microsporidia) of insects.

Comparative Genomics of a Host-Parasitoid System. Sequencing of the parasitoid wasp *N. vitripennis* has been completed. *Nasonia* is a parasitoid of the house fly (*Nasonia* is sold commercially for fly control). Having the genome of both the parasitoid (*Nasonia*) and the host (*M. domestica*) will allow the first opportunity to study the interactions between a pest insect and its parasitoid at the whole genome level. How do *Nasonia* eggs evade the immune response of the house fly? What is the response of the house fly to *Nasonia* venoms? What immunological factors permit some flies but not others to destroy the developing *Nasonia* embryo? Sequencing of the house fly genome is strongly supported by the *Nasonia* community (Scott et al. 2008).

Improving Genome Annotation, Especially Within Insects and Diptera. The Dipteran clade has radiated into >120,000 known species since its origin in the late Jurassic. *M. domestica* is well placed within the Diptera to maximize the utility of sequence data for comparison between existing Dipteran genomes. Although systematic/phylogenetic research on Diptera has been carried out for more than a century, a well-supported tree for the entire order has not been completed (Yeates and Wiegmann 1999). However, it is clear that house fly and *Drosophila* represent a different suborder than *Anopheles*, and house fly represents a different Section (Calyptrate) than *Drosophila* (Acalyptrate) (note: some classifications differ in the taxonomic level where they split calyptrate and acalyptrate flies). Multiple, deeply divergent comparisons within the order allows identification of lineage effects on rates and patterns of genomic diversity. These comparisons become more powerful in elucidating genome evolution as the phylogenetic context is broadened. Given the well-centered position between *Drosophila* and mosquitoes, the *Musca* genome would be nearly ideal for leveraging analysis and annotation of the *Anopheles* and *Aedes* genomes by bridging this gap. In addition, the house fly genome will provide a valuable outgroup for analyses of *Drosophila*

genomes, given their more recent common ancestor (compared with *Drosophila* versus *Anopheles* or *Aedes*). The deepest common ancestor to the set of *Drosophila* species whose genomes were sequenced is estimated to be 60–40 million years ago (MYA), and the common ancestor between *D. melanogaster* and *M. domestica* has been estimated to be \approx 100 MYA (Beverley and Wilson 1984). This places it remarkably well in the gap between *Drosophila* and *Anopheles* and will allow a very broad evolutionary analysis across the Dipteran order.

The *Glossina* genome has recently been suggested as worthy for having its genome sequenced (Aksoy et al. 2005), and this effort is underway. However, this does not disqualify the house fly for a number of reasons. First, the *Glossina* genome is quite large (500–600 Mbp) and contains numerous repeat sequences (Aksoy et al. 2005) that will make the sequencing effort quite difficult. Second, *Glossina* and *Musca* would both provide useful outgroups for *Drosophila* and mosquito genomes (without being redundant). At the same time, they would provide useful comparisons with each other at a level of difference similar to a comparison of *Anopheles* and *Aedes*. In addition, house flies have many unique features (including their profound impact on human and animal health) and have entirely different lifestyles, habitats, and behaviors from *Glossina*, all of which justify sequencing the house fly genome, whether or not *Glossina* is sequenced. Furthermore, house fly has far better genetics; more is known about its sex determination, physiology, biochemistry, neurobiology, and evolution. The house fly is a global pest, whereas *Glossina* is a pest only in Africa.

The completed genome sequences of *Drosophila melanogaster* and *Anopheles gambiae* have been extremely valuable for deductions about the evolutionary origins, structure, and even the function of many human genes (Kortschak et al. 2003). Nevertheless, a significant number of gene modifications and extensive gene loss has occurred in *Drosophila*. Although the genomes and proteomes of *An. gambiae* and *D. melanogaster*, which diverged \approx 220–240 MYA (Wiegmann et al. 2003), show considerable similarities, both lineages have experienced multiple gene acquisitions and losses, especially through expansions and contractions of gene families (Zdobnov et al. 2002). Sequences of orthologous genes in these two insect species have diverged to the point that synonymous positions are virtually randomized (Zdobnov et al. 2002). It was hoped that regulatory regions of genes would become clear by comparison of 5' regions of genes in *Drosophila* and *Anopheles*, but this has proven to be much more difficult. For example, the 5' and 3' regulatory flanking regions of many genes in house flies are virtually unalignable to the orthologous sequence in *Drosophila* (Shaw et al. 2001); the genetic cascades regulating sex determination of the house fly and *D. melanogaster* seem strikingly different, and the upstream regulators of sex determination genes are different between these two insect species (Dübendorfer et al. 2002). Furthermore, 24% of *Apis* expressed

sequence tags (ESTs) showed better matches to Chordata than to *Drosophila* genes (Whitfield et al. 2002). Some *Apis* ESTs showed significant matches to human sequences, but no matches to the *Drosophila* genome (inferred to be genes that were lost from *Drosophila*). Similar results have also been identified in the current house fly EST sequences (N.L., unpublished data). Although either *Drosophila* or *An. gambiae* (or both) homologs could be recognized for more than one half of the house fly EST sequences, some of these EST sequences showed better matches to other more distant species, such as *Plasmodium falciparum*, *Carassius auratus* (goldfish), and *Homo sapiens*, than to *Drosophila* and/or *An. gambiae* homologs. Some of the *Musca* EST sequences showed no matches to the *Drosophila* and/or *An. gambiae* genome. These results indicate that the genomic sequences from other insect species will be extremely important for linking human genes to their *Drosophila* or *An. gambiae* homologs.

Understanding the evolution of *cis*-regulatory sequences in *Drosophila* has proven difficult in some cases (e.g., *achaete-scute* genes), because the patterns of expression are not substantially different between *Drosophila* species but are so extremely diverged in *Anopheles* that analysis is difficult. Thus, the house fly genome would provide a critical resource for the analysis of *cis*-regulatory sequences in *Drosophila*.

Conclusions

A thorough understanding of the biology of complex organisms requires complete sequencing information and identification of all functional elements from the genomes of these organisms. The "whole genome" approach has vastly improved comparative and evolutionary studies, as well as physical map building. It has addressed several important scientific questions about genome evolution, such as evolutionary rates, speciation, genome reorganization, and origins of variation. The approach has also been important for identification of conserved sequences involved in gene regulation and other genomic functions, identification of specific functional sequences (i.e., those that have been substituted or modified during evolution, and which have undergone recent selection; Vandahl et al. 2004), and elucidation of sequence variation in the population of organisms (such as alternative splicing in the regulation of gene function; Tan et al. 2002). The whole genome approach will also be important for identification of sequences that are broadly conserved across insect genomes to provide insight into the unique features in the genome and for obtaining a broader and more complete assessment of the extent of genetic variation in the population of organisms; identification of variation in gene expression; and understanding the evolution (Yan et al. 2002). The whole genome is also necessary to understand the interactions of house flies with the parasitoid wasp *N. vitripennis*. Although some of the genes that are expected to be regulated by parasitoid venoms and egg laying could be inferred from other studies, only a whole genome will provide comprehensive insight

into the genes involved in host/parasitoid interactions.

The calyptrate flies, with *M. domestica* as the most prominent experimental organism, includes a large number of important vectors of human and veterinary diseases, as well as important species for forensic entomology: dog dung fly (*Musca sorbens*), face fly (*Musca autumnalis*), blow flies (*Lucilia*, *Calliphora*, *Chrysomya*), flesh flies (*Sarcophaga*), screwworm (*Cochliomyia*), tsetse fly (*Glossina*), the little house fly (*Fannia*), warble flies (*Hypoderma*), yellow dung flies (*Scathophaga*), and the root maggot fly (*Anthomyiida*). By using genetic manipulations of *M. domestica* to place function of novel genes in its genome, we anticipate that it will be easy to transfer the knowledge gained to other synanthropic flies.

Many genes, especially regulatory genes, are often expressed at a very low level, and they would be rare in EST libraries. The entire house fly genome sequence will, especially when compared with the *Drosophila* and mosquito genome sequences, facilitate the identification of homologous genes expressed at low levels or in a specific tissue. Expression patterns can be validated with high-throughput real-time PCR systems for use in either general population or micro-evolutionary studies (e.g., the spread and fitness of resistance genes).

Currently there are ≈ 40 laboratories worldwide whose primary research focus is the house fly. About one half of these are engaged in studies of molecular biology that would immediately benefit from a complete genome sequence. Most of the others are studying aspects of toxicology and pest control, and immediate access to design of primers for PCR analysis would open the door to simple but powerful molecular approaches to this group. *Drosophila* researchers would benefit from, and are strongly supportive of (Scott et al. 2008), sequencing of the house fly genome. The white papers that resulted in funding to sequence an additional 11 genomes of *Drosophila* species failed to include an outgroup to the set of *Drosophila* species, and *Anopheles gambiae* is just too distantly related for optimal analysis (in most cases).

Suitable tools are available to facilitate sequencing of the house fly genome, and the scientific community is solidly behind this effort. Several cDNA and genomic libraries (various tissues, strains, and life stages) have been prepared, and at least two pilot EST projects are underway. More than 450 nucleotide and >750 EST sequences from *M. domestica* can be found in GenBank. As a part of USDA NRI and Auburn University Biogrant funded project, a house fly normalized cDNA library has been constructed from the mRNA of house flies. More than 300 ESTs have been generated, resulting in 292 high-quality cDNA sequence reads. Thirty-nine ESTs were assembled into eight contigs. The remaining 253 ESTs are unique, suggesting a 15% redundancy in the house fly sequence set. This EST sequencing effort, combined with other larger EST projects, will be excellent resources for the genomic library (BAC library) screening and building contig maps for comparative genomic

studies. House flies can be readily transformed with mobile elements such as piggyback, hermes, or hobo (Atkinson et al. 1993; O'Brochta et al. 1994, 1996; Warren et al. 1994; O'Brochta and Atkinson 1996, 1997; Sarkar et al. 1997; Hediger et al. 2001), and some genes can be silenced using RNAi (McGregor et al. 2001, Burghardt et al. 2005).

Letters of support written for the house fly white paper (Scott et al. 2008) eloquently showed how researchers from diverse scientific areas (genomics, proteomics, developmental biology, population genetics, evolutionary biology, etc.) that use a wide range of study animal (mosquitoes, *Drosophila*, *Nasonia*, *Tribolium*, *Musca*, etc.) would make immediate use of the *M. domestica* genome sequence to accelerate their research programs on fundamental aspects of genetics (sex determination, dosage compensation, olfaction, immunology, etc.), as well as practical problems of pest control. It is clear that the scientific community considers sequencing of the house fly genome to be an extremely high priority.

References Cited

- Adams, M. D., S. E. Celniker, R. A. Holt, C. A. Evans, J. D. Gocayne, P. G. Amanatides, S. E. Scherer, P. W. Li, R. A. Hoskins, R. F. Galle, et al. 2000. The genome sequence of *Drosophila melanogaster*. *Science* 287: 2185–2195.
- Aksoy, S., M. Merriman, N. Hall, H. Masahira, W. Hide, and M. J. Lehane. 2005. A case for a Glossina genome project. *Trends Parasitol.* 21: 107–111.
- Atkinson, P. W., W. D. Warren, and D. A. O'Brochta. 1993. The hobo transposable element of *Drosophila* can be cross-mobilized in houseflies and excises like the Ac element of maize. *Proc. Natl. Acad. Sci. U.S.A.* 90: 9693–9697.
- Axtell, R. C. 1990. Potential of biocontrol for livestock and poultry pests, pp. 293. In D. A. Rutz and R. S. Patterson (eds.), *Biocontrol of arthropods affecting livestock and poultry*. Westview Press, Boulder, CO.
- Beverley, S. M., and A. Wilson. 1984. Molecular evolution in *Drosophila* and the higher Diptera II. A time scale for fly evolution. *J. Molec. Evol.* 21: 1–13.
- Boulesteix, G., P. Le Dantec, B. Chevalier, M. Dieng, B. Niang, and B. Diatta. 2005. Role of *Musca domestica* in the transmission of multiresistant bacteria in the centres of intensive care setting in sub-Saharan Africa. *Réanimation* 24: 361–365.
- Bull, J. J., and E. L. Charnov. 1977. Changes in the heterogametic mechanism of sex determination. *Heredity* 39: 1–14.
- Burghardt, G., M. Hediger, C. Siegenthaler, M. Moser, A. Dübendorfer, and D. Bopp. 2005. The *transformer2* gene in *Musca domestica* is required for selecting and maintaining the female pathway of development. *Dev. Genes Evol.* 215: 165–176.
- Casida, J. E., and G. B. Quistad. 2004. Why insecticides are more toxic to insects than people: the unique toxicology of insects. *J. Pestic. Sci.* 29: 81–96.
- Christophides, G. K., E. M. Zdobnov, C. Barillas-Mury, E. Birney, S. Blandin, C. Blass, P. T. Brey, F. H. Collins, A. Danielli, G. Dimopoulos, et al. 2002. Immunity-related genes and gene families in *Anopheles gambiae*. *Science* 298: 159–165.
- Darbro, J. M., and B. A. Mullens. 2004. Assessing insecticide resistance and aversion to methomyl-treated toxic baits in *Musca domestica* L. (Diptera: Muscidae) populations in southern California. *Pest. Manag. Sci.* 60: 901–908.
- de Jesus, A. J., A. R. Olsen, J. R. Bryce, and R. C. Whiting. 2004. Quantitative contamination and transfer of *Escherichia coli* from foods by houseflies, *Musca domestica* L. (Diptera: Muscidae). *Int. J. Food Microbiol.* 93: 259–262.
- Denholm, I., M. G. Franco, P. G. Rubini, and M. Vecchi. 1983. Identification of a male determinant on the X chromosome of housefly (*Musca domestica* L.) populations in South-East England. *Genet. Res. Camb.* 42: 311–322.
- Denholm, I., P. G. Rubini, C. Rovati, and M. Vecchi. 1990. Genetic basis of sex determination in two South African strains of house fly. *S. African J. Sci.* 86: 41–43.
- Dübendorfer, A., M. Hediger, G. Burghardt, and D. Bopp. 2002. *Musca domestica*, a window on the evolution of sex-determining mechanisms in insects. *Int. J. Dev. Biol.* 46: 75–79.
- Emerson, P. M., S. W. Lindsay, G.E.L. Walraven, H. Faal, C. Bogh, K. Lowe, and R. L. Bailey. 1999. Effect of fly control on trachoma and diarrhoea. *Lancet* 353: 1401–1403.
- Emerson, P. M., R. L. Bailey, and O. S. Mahdi. 2000. Transmission ecology of the fly *Musca sorbens*, a putative vector of trachoma. *Trans. R. Soc. Trop. Med. Hyg.* 94: 28–32.
- Focke, M., W. Hemmer, and S. Wohrl. 2003. Specific sensitization to the common housefly (*Musca domestica*) not related to insect panallergy. *Allergy* 58: 448–451.
- Franco, M. G., P. G. Rubini, and M. Vecchi. 1982. Sex-determinants and their distribution in various populations of *Musca domestica* L. of Western Europe. *Genet. Res. Camb.* 40: 279–293.
- Gao, J., and J. G. Scott. 2006. Use of quantitative real-time PCR to estimate the size of the house fly (*Musca domestica*) genome. *Insect Molec. Biol.* 15: 835–837.
- Gao, J.-R., J. M. Deacutis, and J. G. Scott. 2007a. Characterization of the nicotinic acetylcholine receptor subunit gene *Mda2* from the housefly, *Musca domestica*. *Arch. Insect Biochem. Physiol.* 64: 30–42.
- Gao, J.-R., J. M. Deacutis, and J. G. Scott. 2007b. The nicotinic acetylcholine receptor subunit *Mda6* from *Musca domestica* is diversified via post transcriptional modification. *Insect Molec. Biol.* 16: 325–334.
- Gao, J.-R., J. M. Deacutis, and J. G. Scott. 2007c. Characterization of the nicotinic acetylcholine receptor subunits *Mdalpha5* and *Mdbeta3* on autosome 1 of *Musca domestica* indicate they are not involved in spinosad resistance. *Insect Molec. Biol.* 16: 691–701.
- Geden, C. J., J. J. Arends, R. C. Axtell, D. R. Barnard, D. M. Gaydon, L. A. Hickie, J. A. Hogsette, W. F. Jones, B. A. Mullens, M. P. Nolan, Jr., M. P. Nolan, III, J. J. Petersen, and D. C. Sheppard. 1994. Economic significance of poultry. In C. J. Geden and J. A. Hogsette (eds.), *Research and extension needs for integrated pest management for arthropods of veterinary importance*. (<http://cmave.usda.uff.edu/lincoln.html>).
- Graczyk, T. K., R. Knight, R. H. Gilman, and M. R. Cranfield. 2001. The role of non-biting flies in the epidemiology of human infectious diseases. *Microbes Infect.* 3: 231–235.
- Greenberg, B. 1965. Flies and disease. *Sci. Am.* 213: 92–99.
- Hald, B., H. Skovgard, D. D. Bang, K. Pedersen, J. Dybdahl, J. B. Jespersen, and M. Madsen. 2004. Flies and Campylobacter infection of broiler flocks. *Emerg. Infect. Dis.* 10: 1490–1492.
- Hamm, R. L., and J. G. Scott. 2009. A high frequency of male determining factors in male *Musca domestica* (Diptera: Muscidae) from Ipswich, Australia. *J. Med. Entomol.* 46: 169–172.
- Hamm, R. L., T. Shono, and J. G. Scott. 2005. A cline in frequency of autosomal males is not associated with in-

- secticide resistance in house fly (Diptera: Muscidae). *J. Econ. Entomol.* 98: 171–176.
- Hanley, M. E., D. W. Dunn, S. R. Abolins, and D. Goulson. 2004. Evaluation of (Z)-9-tricosene baited targets for control of the housefly (*Musca domestica*) in outdoor situations. *J. Appl. Entomol.* 128: 478–482.
- Hediger, M., M. Niessen, E. A. Wimmer, A. Dübendorfer, and D. Bopp. 2001. Genetic transformation of the housefly *Musca domestica* with the lepidopteran derived transposon piggyBac. *Insect Molec. Biol.* 10: 113–119.
- Hediger, M., G. Burghardt, C. Siegenthaler, N. Buser, D. Hilfiker-Kleiner, A. Dübendorfer, and D. Bopp. 2004. Sex determination in *Drosophila melanogaster* and *Musca domestica* converges at the level of the terminal regulator *doublesex*. *Dev. Genes Evol.* 214: 29–42.
- Hiroyoshi, T. 1960. Some new mutants and linkage groups of the house fly. *J. Econ. Entomol.* 53: 985–990.
- Hiroyoshi, T. 1964. Sex-limited inheritance and abnormal sex ratio in strains of the housefly. *Genetics* 50: 373–385.
- Hiroyoshi, T. 1977. Some new mutants and revised linkage maps of the housefly, *Musca domestica* L. *Jpn. J. Genet.* 52: 275–288.
- Inoue, H., Y. Fukumori, and T. Hiroyoshi. 1983. Mapping of autosomal male-determining factors of the housefly, *Musca domestica* L., by means of sex reversal. *Jpn. J. Genet.* 58: 451–461.
- Jensen, A. B., L. Thomsen, and J. Eilenberg. 2001. Intraspecific variation and host specificity of *Entomophthora muscae sensu stricto* isolates revealed by random amplified polymorphic DNA, universal primed PCR, PCR-restriction fragment length polymorphism, and conidial morphology. *J. Invertebr. Pathol.* 78: 251–259.
- Kalsbeek, V., B. A. Mullens, and J. B. Jespersen. 2001. Field studies of *Entomophthora* (Zygomycetes: Entomophthorales)-induced behavioral fever in *Musca domestica* (Diptera: Muscidae) in Denmark. *Biol. Control* 21: 264–273.
- Keiding, J. 1986. The house fly: biology and control. WHO Vector Control Series 63.
- Keiding, J. 1999. Review of the global status and recent development of insecticide resistance in field populations of the housefly, *Musca domestica* (Diptera: Muscidae). *Bull. Entomol. Res.* 89: S7–S67.
- Kelling, F. J., G. Biancaniello, and C. J. den Otter. 2002. Electrophysiological characterization of olfactory cell types in the antennae and palps of the housefly. *J. Insect Physiol.* 48: 997–1008.
- Kelling, F. J., G. Biancaniello, and C. J. den Otter. 2003. Effect of age and sex on the sensitivity of antennal and palpal olfactory cells of houseflies. *Entomol. Exp. Appl.* 106: 45–51.
- Kortschak, R. D., G. Samuel, R. Saint, and D. J. Miller. 2003. EST analysis of the cnidarian *Acropora millepora* reveals extensive gene loss and rapid sequence divergence in the model invertebrates. *Curr. Biol.* 13: 2190–2195.
- Kozielska, M., B. Feldmeyer, I. Pen, F. J. Weissing, and L. W. Beukebook. 2008. Are autosomal sex-determining factors of the housefly (*Musca domestica*) spreading north? *Genet. Res. Camb.* 90: 157–165.
- Lazzaro, B. P., and A. G. Clark. 2003. Molecular population genetics of inducible antibacterial peptide genes in *Drosophila melanogaster*. *Molec. Biol. Evol.* 20: 914–923.
- Lazzaro, B. P., B. K. Scurman, and A. G. Clark. 2004. Genetic basis of natural variation in *D. melanogaster* antibacterial immunity. *Science* 303: 1873–1876.
- Li, S. E., and F. J. Stutzenberger. 2000. The housefly (*Musca domestica*) as a possible vector for *Helicobacter pylori* at agricultural sites. *Intern. J. Environ. Health Res.* 10: 141–152.
- Macovei, L., and L. Zurek. 2006. Ecology of antibiotic resistance genes: characterization of enterococci from houseflies collected in food settings. *Appl. Environ. Microbiol.* 72: 4028–4035.
- Maisnier-Patin, S., and D. I. Andersson. 2004. Adaptation to the deleterious effects of antimicrobial drug resistance mutations by compensatory evolution. *Res. Microbiol.* 155: 360–369.
- Matsuda, K., S. D. Buckingham, D. Kleier, J. J. Rauh, M. Grauso, and D. B. Sattelle. 2001. Neonicotinoids: insecticides acting on insect nicotinic acetylcholine receptors. *Trends Pharmacol. Sci.* 22: 573–580.
- McGregor, A. P., P. J. Shaw, J. M. Hancock, D. Bopp, M. Hediger, N. S. Wratten, and G. A. Dover. 2001. Rapid restructuring of bicoid-dependent hunchback promoters within and between Dipteran species: implications for molecular coevolution. *Evol. Develop.* 3: 397–407.
- Meise, M., D. Hilfiker-Kleiner, A. Dübendorfer, C. Brunner, R. Nothiger, and D. Bopp. 1998. *Sex-lethal*, the master sex-determining gene in *Drosophila*, is not sex-specifically regulated in *Musca domestica*. *Development* 125: 1487–1494.
- Milani, R., P. G. Rubini, and M. G. Franco. 1967. Sex-determination in the housefly. *Genetica Agaria* 21: 385–411.
- Moriya, K., T. Fujibayashi, T. Yoshihara, A. Matsuda, N. Sumi, N. Umezaki, H. Kurahashi, N. Agui, A. Wada, and H. Watanabe. 1999. Verotoxin-producing *Escherichia coli* O157: H7 carried by the housefly in Japan. *Med. Vet. Entomol.* 13: 214–216.
- Mündi, K. 1994. Housefly, an everyday monster. Fox Lorber, New York.
- Nauen, T., U. Ebbinghaus-Kintscher, V. Salgado, and M. Kaussmann. 2003. Thiamethoxam is a neonicotinoid precursor converted to clothianidin in insects and plants. *Pestic. Biochem. Physiol.* 76: 55–69.
- Nickel, C. A., and D. E. Wagoner. 1974. Mutants on linkage groups 3 and 4 of the house fly. *Ann. Entomol. Soc. Am.* 67: 775–776.
- O'Brochta, D. A., and P. Atkinson. 1996. Transposable elements and gene transformation in non-Drosophilid insects. *Insect Biochem. Molec. Biol.* 26: 739–753.
- O'Brachta, D. A., and P. W. Atkinson. 1997. Recent developments in transgenic insect technology. *Parasitol. Today* 13: 99–104.
- O'Brachta, D. A., W. D. Warren, K. J. Saville, and P. W. Atkinson. 1996. *Hermes*, a functional non-Drosophilid insect gene vector from *Musca domestica*. *Genetics* 142: 907–914.
- O'Brochta, D. A., W. D. Warren, K. J. Saville, and P. W. Atkinson. 1994. Interplasmid transposition of *Drosophila* hobo elements in non-drosophilid insects. *Mol. Gen. Genet.* 244: 9–14.
- Osta, M. A., G. K. Christophides, D. Vlachou, and F. C. Kafatos. 2004. Innate immunity in the malaria vector *Anopheles gambiae*: comparative and functional genomics. *J. Exp. Biol.* 207: 2551–2563.
- Rahuma, N., K. S. Ghenghesh, R. Ben-Aissa, and A. Elamaari. 2005. Carriage by the housefly (*Musca domestica*) of multiple-antibiotic-resistant bacteria that are potentially pathogenic to humans, in hospital and other urban environments in Misurata, Libya. *Ann. Trop. Med. Parasitol.* 99: 795–802.
- Reed, D. H., and E. H. Bryant. 2004. Phenotypic correlations among fitness and its components in a population of the housefly. *J. Evol. Biol.* 17: 919–923.

- Sarkar, A., C. J. Coates, S. Whyard, U. Willhoeft, P. W. Atkinson, and D. A. O'Brachta. 1997. The *Hermes* element from *Musca domestica* can transpose in four families of cyclorrhaphan flies. *Genetica* 99: 15–29.
- Sasaki, T., M. Kobayashi, and N. Agui. 2000. Epidemiological potential of excretion and regurgitation by *Musca domestica* (Diptera: muscidae) in the dissemination of *Escherichia coli* O157: H7 to food. *J. Med. Entomol.* 37: 945–949.
- Schlenke, T. A., and D. J. Begun. 2003. Natural selection drives *Drosophila* immune system evolution. *Genetics* 164: 1471–1480.
- Schmidt, R., M. Hediger, S. Roth, R. Nothiger, and A. Dübendorfer. 1997. The Y-chromosomal and autosomal male-determining *M* factors of *Musca domestica* are equivalent. *Genetics* 147: 271–280.
- Scott, H. G., and K. S. Lettig. 1962. Flies of public health importance and their control. U.S. Government Printing Office, Washington, DC.
- Scott, J. G. 1990. Investigating mechanisms of insecticide resistance: methods, strategies and pitfalls, pp. 39–57. In R. T. Roush and B. Tabashnik (eds.), *Pesticide resistance in arthropods*. Chapman & Hall, New York.
- Scott, J. G. 1991. Insecticide resistance in insects, pp. 663. In D. Pimentel (ed.), *Handbook of pest management in agriculture*, 2nd ed. CRC, Boca Raton, FL.
- Scott, J. G. 1999. Cytochromes P450 and insecticide resistance. *Insect Biochem. Molec. Biol.* 29: 757–777.
- Scott, J. G., N. Liu, and M. Kristensen. 2008. Rationale for sequencing the genome of the house fly, *Musca domestica*. (http://www.entomology.cornell.edu/Faculty_Staff/Scott/whitepaper.pdf).
- Shaw, P. J., A. Salameh, A. P. McGregor, S. Bala, and G. A. Dover. 2001. Divergent structure and function of the bicoid gene in Muscoidea fly species. *Evol. Develop.* 3: 251–259.
- Srinivasan, P., E. G. Abraham, A. Ghosh, K., J. Valenzuela, J.M.C. Ribeiro, G. Dimopoulos, F. C. Kafatos, J. H. Adams, H. Fujioka, and M. Jacobs-Lorena. 2004. Analysis of the *Plasmodium* and *Anopheles* transcriptomes during oocyst differentiation. *J. Biol. Chem.* 279: 5581–5587.
- Sundin, G. W. 1996. Evolution and selection of antibiotic and pesticide resistance: a molecular genetic perspective, pp. 97–105. In T. M. Brown (ed.), *Molecular genetics and evolution of pesticide resistance*, 645th ed. American Chemical Society, Washington, DC.
- Tan, J., Z. Z. Liu, Y. Nomura, A. L. Goldin, and K. Dong. 2002. Alternative splicing of an insect sodium channel gene generates pharmacologically distinct sodium channels. *J. Neurosci.* 22: 5300–5309.
- Tomita, T., and Y. Wada. 1989. Multifactorial sex determination in natural populations of the housefly (*Musca domestica*) in Japan. *Jpn. J. Genet.* 64: 373–382.
- Tomizawa, M., and J. E. Casida. 2003. Selective toxicity of neonicotinoids attributable to specificity of insect and mammalian nicotinic receptors. *Annu. Rev. Entomol.* 48: 339–364.
- Tsukamoto, M., Y. Baba, and S. Hiraga. 1961. Mutations and linkage groups in Japanese strains of the housefly. *Jpn. J. Genet.* 36: 168–174.
- Vandahl, B.B.S., S. Birkelund, and G. Christiansen. 2004. Genomic and proteome analysis of *Chlamydia*. *Proteomics* 4: 2831–2842.
- Wagoner, D. E. 1969. Presence of male determining factors found on three autosomes in the house fly, *Musca domestica*. *Nature* 223: 187–188.
- Wahl, R., and J. Fraedrich. 1997. Occupational allergy to the housefly (*Musca domestica*). *Allergy* 52: 236–238.
- Wakita, T., K. Kinoshita, E. Yamada, N. Yasui, N. Kawahara, A. Naoi, M. Nakaya, K. Ebihara, H. Matsuno, and K. Kodaka. 2003. The discovery of dinotefuran: a novel neonicotinoid. *Pest. Manag. Sci.* 59: 1016–1022.
- Warren, W. D., P. W. Atkinson, and D. A. O'Brachta. 1994. The *Hermes* transposable element from the house fly, *Musca domestica*, is a short inverted repeat-type element of the hobo, Ac, Tam3 (hAT) element family. *Genet. Res. Camb.* 64: 87–97.
- Whitfield, C. W., M. R. Band, M. F. Bonaldo, C. G. Kumar, L. Liu, J. R. Pardinis, H. M. Robertson, M. B. Soares, and G. E. Robinson. 2002. Annotated expressed sequence Tags and cDNA microarrays for studies of brain and behavior in the honey bee. *Genome Res.* 12: 555–566.
- Wiegmann, B. M., D. K. Yeates, J. L. Thorne, and H. Kishino. 2003. Time flies, a new molecular time-scale for brachyceran fly evolution without a clock. *Syst. Biol.* 52: 745–756.
- Wilhelm, J., A. Pingoud, and M. Hahn. 2003. Real-time PCR-based method for the estimation of genome sizes. *Nucleic Acids Res.* 31: e56.
- World Health Organization. 2004. Trachoma. (<http://www.who.int/pbd/blindness/trachoma/en/>).
- Yan, H., W. Yuan, V. E. Velculescu, B. Vogelstein, and K. W. Kinzler. 2002. Allelic variation in human gene expression. *Science* 1143.
- Yeates, D. K., and B. M. Wiegmann. 1999. Congruence and controversy: toward a higher-level phylogeny of Diptera. *Annu. Rev. Entomol.* 44: 397–428.
- Zdobnov, E. M., C. von-Mering, I. Letunic, D. Lorrents, M. Suyama, R. R. Copley, G. K. Christophides, D. Thomasova, R. A. Holt, G. M. Subramanian, et al. 2002. Comparative genome and proteome analysis of *Anopheles gambiae* and *Drosophila melanogaster*. *Science* 298: 149–159.
- Zurek, L., D. W. Watson, S. B. Krasnoff, and C. Schal. 2002. Effect of the entomopathogenic fungus, *Entomophthora muscae* (Zygomycetes : Entomophthoraceae), on sex pheromone and other cuticular hydrocarbons of the house fly, *Musca domestica*. *J. Invertebr. Pathol.* 80: 171–176.

Received 14 June 2008; accepted 4 November 2008.