HOMOPLASIOUS CHARACTER COMBINATIONS AND GENERIC DELIMITATION: A CASE STUDY FROM THE INDO-PACIFIC ARECOID PALMS (ARECACEAE: ARECEAE)¹

MARIA VIBE NORUP,^{2,3,5} JOHN DRANSFIELD,³ MARK W. CHASE,³ ANDERS S. BARFOD,²
EDWINO S. FERNANDO,⁴ AND WILLIAM J. BAKER³

²Department of Systematic Botany, University of Aarhus, Ny Munkegade, Building 540, 8000 Aarhus, Denmark; ³Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AB, UK; and ⁴Department of Forest Biological Sciences, The University of the Philippines, Los Baños, College, 4031 Laguna, Philippines

The complex distributions of morphological character states in the Indo-Pacific palm tribe Areceae (Arecaceae; Arecoideae) are potentially challenging for the delimitation of its genera. In the first exhaustive sampling of all 65 genera of the Areceae, we examined relationships of two of the tribe's most problematic genera, *Heterospathes* and *Rhopaloblastes*, using portions of the low-copy nuclear genes phosphoribulokinase (*PRK*) and RNA-polymerase II subunit B (*RPB2*). Both genera fell within a highly supported clade comprising all Areceae genera, but are clearly unrelated. *Rhopaloblastes* was strongly supported as monophyletic and is most closely related to Indian Ocean genera. *Heterospathes* was resolved with strong support within a clade of western Pacific genera, but with the monotypic *Alsmithia* nested within it. *Ptychosperma micranthum*, which has previously been included in both *Heterospathes* and *Rhopaloblastes*, is excluded from these and from *Ptychosperma*, supporting its recent placement in a new genus *Dransfieldia*. Morphological comparisons indicate that the crownshaft is putatively synapomorphic for the Areceae with numerous reversals within the clade and some independent origins elsewhere. The putative diagnostic characters of *Heterospathes* show high levels of homoplasy, and the genus can only be distinguished by a suite of characters, whereas *Rhopaloblastes* is more clearly defined. Our results have implications not only for the two genera in focus, but have also been influential for the new classification of the Areceae.

Key words: Arecaceae; biogeography; *Heterospathes*; low-copy nuclear DNA; molecular phylogenetics; morphology; pseudomonomy; *Rhopaloblastes*.

Extremely high levels of morphological homoplasy in large and species-rich plant groups (e.g., Baker et al., 2000a; Moylan et al., 2004; Tam et al., 2004) are frequently responsible for problems in generic delimitation. Furthermore, while genus circumscriptions may be strongly supported by evidence from DNA sequence data, unique morphological synapomorphies may be lacking, the group being defined on combinations of homoplasious character states (e.g., Hughes et al., 2004). Within the palm family, generic limits are especially challenging in tribe Areceae (Arecoideae), a species-rich group comprising more than 660 species in 65 genera in the latest classification of palms (Dransfield et al., 2005; Govaerts and Dransfield, 2005). However, a detailed and well-resolved phylogenetic study of the group with which generic limits may be tested is not yet available.

The Areceae is an Old World tribe distributed throughout the Indo-Pacific region from Pemba in the west to Samoa in the

east, with centers of diversity in Madagascar, the Sunda Shelf, Papuasias, and the western Pacific region (Dransfield et al., 2005; Govaerts and Dransfield, 2005). The great diversification of Areceae in the area is paralleled by other higher plants and has most likely been spurred on by the complex geological and climatic history of the region and the many niches presented by the numerous islands and archipelagos of the Indo-Pacific region (Turner, 1995; Ridder-Numan, 1996; Dransfield, 1999; Morley, 2000, 2001). Most members of the tribe are medium-sized to large, monoecious palms characterized by reduplicate pinnate leaves and a crownshaft (a conspicuous cylinder of tubular, neatly abscising leaf sheaths at the stem tip). Inflorescences in bud are usually enclosed within a prophyll and typically one conspicuous peduncular bract. In the pistillate flower, the gynoeceum comprises one fertile carpel with a single ovule and remnants of two aborted carpels and is thus termed pseudomonomerous.

The delimitation of the Areceae has recently been revised in the new phylogenetic classification of palms that is followed here (Dransfield et al., 2005). Two recent molecular phylogenetic studies focusing on relationships within Arecoideae have indicated that the Areceae as defined in the previous classification (Uhl and Dransfield, 1987; Dransfield and Uhl, 1998) is not monophyletic (Hahn, 2002; Lewis and Doyle, 2002). These authors also identified a strongly supported clade of Indo-Pacific, pseudomonomerous genera that now equates to tribe Areceae sensu Dransfield et al. (2005). This clade was also recovered in family-wide phylogenetic studies, although without bootstrap support greater than 50% (Baker et al., 1999; Asmussen et al., 2000; Asmussen and Chase, 2001; Asmussen et al., in press). The majority of genera of Areceae sensu Uhl

¹ Manuscript received 21 May 2005; revision accepted 27 April 2006.

The authors thank V. Savolainen, A. Loo, C. Lewis, and J. Roncal for sharing unpublished sequences for this paper, D. Springate, O. Maurin, and T. Wilmot for helping with sequence generation, the staff at the Jodrell Laboratory, RBG Kew, for technical assistance, C. Lewis, C. Asmussen, J.-C. Pintaud, R. Banka, R. Maturbongs, J. Wanggai, C. Heatubun, A. Loo, P. Forster, and S. Zona for supplying DNA samples, and C. Asmussen and two anonymous reviewers for providing constructive comments on earlier versions of the manuscript. M. V. N. was able to conduct her studies at the Royal Botanic Gardens, Kew, thanks to support from the Oticon Foundation, the Faculty of Science, University of Aarhus, and the Royal Botanic Gardens, Kew. DNA sequencing was in part funded by grants from the Royal Society, the Leverhulme Trust, and the British American Tobacco Biodiversity Partnership.

⁵ Author for correspondence (e-mail: mnorup@snm.ku.dk)

and Dransfield (1987) are included within the limits of the revised concept of the tribe, the remainder being placed in other tribes such as Euterpeae, Oranieae, and Roystoneae. Surprisingly, two Pacific genera, *Pelagodoxa* Becc. and *Sommieria* Becc., which conform to the Areceae both in their biogeography and in their pseudomononamous gynoeceia (Stauffer et al., 2004), are also excluded from the Areceae according to the analysis of Lewis and Doyle (2002) and have consequently been placed in their own tribe Pelagodoxeae (Dransfield et al., 2005).

Generally, the Areceae is characterized by a complex yet limited range of morphological diversity with genera distinguished on combinations of widespread character states. This situation is highly problematic for identification and delimitation of genera. The current paper focuses on two such genera within the Areceae (Lewis and Doyle, 2002), *Heterospatha* Scheff. and *Rhopaloblade* Scheff. (Areceae, subtribe Iguanurinae; Uhl and Dransfield, 1987), the circumscription of which has caused considerable confusion in the past. *Heterospatha* contains 38 species with a center of species diversity in New Guinea that extends into the western Pacific, particularly the Solomon Islands, the Philippines, and the Moluccas (Govaerts and Dransfield, 2005). The six species of *Rhopaloblade* are disjunctly distributed across Wallace's Line with one species in the Nicobar Islands, one in Peninsular Malaysia and Singapore, and the remaining four species from the Moluccas to New Guinea and the Solomons (Banka and Baker, 2004). Like many other Areceae, *Heterospatha* and *Rhopaloblade* appear to be defined by potentially homoplasious characters. For example, inflorescence morphology of *Heterospatha* is highly reminiscent of that in *Alsmithia* H. E. Moore, and there are numerous examples of other Areceae that, like *Heterospatha*, do not possess a crownshaft (e.g., *Lepidorrhachis* (H. Wendl. & Drude) O.F. Cook and *Cyphosperma* H. Wendl.). A recent revision of *Rhopaloblade* (Banka and Baker, 2004) has to some extent clarified the diagnostic characters of this genus, although the uniqueness of any of these features has not been established. For example, several other palm genera, such as *Cyrtostachys* Blume, display the strongly recurving primary inflorescence branches that are said to be distinctive in *Rhopaloblade*.

Heterospatha and *Rhopaloblade* are currently accepted as distinct genera, but their taxonomic histories are intertwined with several other genera, a common situation throughout the Areceae. The two genera were originally described by Scheffer (1876) in the same publication. Today, a number of species currently accepted in *Heterospatha* were originally described in *Rhopaloblade* (e.g., *R. macgregorii* Becc., transferred to *H. macgregorii* (Becc.) H. E. Moore; Moore, 1970). In addition, several species currently placed in both *Heterospatha* and *Rhopaloblade* were originally described in another genus, *Ptychoraphis* Becc. (Beccari, 1885). *Ptychoraphis*, however, is problematic because the distinguishing characters are reminiscent of characters used to define *Heterospatha* and *Rhopaloblade* (Moore, 1970). Beccari himself later transferred members of *Ptychoraphis* to both *Heterospatha* (Beccari, 1909) and *Rhopaloblade* (Beccari ex Martelli, 1934, 1935). In his revision of *Rhopaloblade*, Moore (1970) finally sank *Ptychoraphis* and split the remaining species between *Heterospatha* and *Rhopaloblade*. One species remains strikingly problematic. Originally described as *Ptychosperma micranthum* Becc., it combines the features of several genera of Areceae, and its taxonomic position has been disputed ever

since it was described by Beccari (1877). Hooker (1883) referred it to *Rhopaloblade* (although the combination was not validly published until Jackson (1895), but Moore (1970) later decided to include the species in *Heterospatha*. The species does not conform to the modern delimitation of any of the three genera to which it has been ascribed and its correct placement remains unclear.

The tortuous taxonomic history and the lack of clear diagnostic features described above call into question the monophyly of *Heterospatha* and *Rhopaloblade*. Furthermore, a clarification of generic limits is required for on-going taxonomic work in Papuasia, the center of diversity for both genera (Baker, 2002). In a biogeographical context, *Heterospatha* and *Rhopaloblade* are particularly significant because they both have wide distributions. *Rhopaloblade* is in fact one of the few palm genera to occur on islands in both the Indian and Pacific Oceans. To assess the monophyly of *Heterospatha* and *Rhopaloblade* and establish their relationships and the biogeographic patterns within the Areceae, it is vital to include all 65 genera of Areceae in this study principally because the characters defining *Heterospatha* and *Rhopaloblade* are shared with several other genera in the tribe. Thus, in this paper, we present not only a phylogenetic assessment of *Heterospatha* and *Rhopaloblade*, but also the first complete generic-level molecular analysis of the entire tribe.

Due to an exceptionally slow rate of plastid DNA evolution in palms in general (Wilson et al., 1990; Asmussen and Chase, 2001), resolution among genera of the Areceae has only been achieved by combining a number of low-copy nuclear markers (Lewis and Doyle, 2002). Low-copy nuclear regions provide some of the most variable and phylogenetically informative molecular markers available, with up to five times the evolutionary rate of plastid DNA, and are especially advantageous in obtaining resolution among rapidly diversifying lineages or at low taxonomic levels (Mort and Crawford, 2004; Small et al., 2004). Although these loci are still not widely used, they promise to be increasingly important in obtaining phylogenetic results at low taxonomic levels, despite potential obstacles relating to molecular evolutionary dynamics, e.g., gene duplications (Mort and Crawford, 2004; Pfeil et al., 2004). In palm systematics, low-copy nuclear loci have been employed with considerable success at a range of taxonomic levels (Lewis and Doyle, 2001, 2002; Norup et al., 2003; Gunn, 2004; Roncal et al., 2005; Thomas et al., 2006; Loo et al., in press).

Intron 4 of *PRK* and intron 23 of *RPB2* were expected to be useful in reconstructing relationships within the Areceae based on promising results from tribes Areceae, Geonomeae and Hyophorbeae (Lewis and Doyle, 2002; Roncal et al., 2005; Thomas et al., 2006; Loo et al., in press). Phosphoribulokinase is a key regulatory enzyme of the Calvin cycle of photosynthetic carbon dioxide assimilation. RNA-polymerase II subunit B codes for the second largest subunit in the RNA-polymerase-II complex, the enzyme complex responsible for DNA transcription of protein-coding genes, and is found in all eukaryotes. Across organisms as diverse as humans, *Drosophila*, fungi, and plants, exon sequences have been shown to be highly conserved and proposed to be valuable in phylogenetic studies (Kawagishi et al., 1993; Denton et al., 1998; Oxelman and Bremer, 2000; Oxelman et al., 2004; Pfeil et al., 2004; Popp and Oxelman, 2004).

The objectives of our study were (1) to evaluate monophyly of *Heterospatha* and *Rhopaloblade* by sampling extensively

within the two genera and including a complete genus-level sample of the Areceae using the low-copy nuclear DNA regions, intron 4 of *PRK* and intron 23 of *RPB2*; (2) to resolve character transformation patterns in key morphological features used to circumscribe *Heterospathe* and *Rhopaloblaste*; and (3) to assess biogeographic implications of the phylogenetic relationships of the two genera.

MATERIALS AND METHODS

Taxon sampling—DNA sequences of 101 taxa were analyzed in this study, including sequences from all 65 genera of tribe Areceae. We sampled down to species level when appropriate to the focus of the paper (sometimes with more than one collection per species); thus we included 16 collections of *Heterospathe*, five collections of *Rhopaloblaste* and two species of *Ptychosperma*, the problematic *P. micranthum*, and the more typical *P. salomonense* Burret. Our sampling was limited only by the remoteness of many of their provenances. The unnamed accessions of *Heterospathe* collected by Baker represent a range of distinct species, but the inadequacies of available taxonomic treatments preclude assignment of species epithets at this time. Outgroup species consisted of a number of putative near-relatives of the Areceae: *Asterogyne martiana* (H. Wendl.) H. Wendl., *Calyptrogyne costatifrons* (L. H. Bailey) Nevers, *Geonoma deversa* (Poi.) Kunth., *Leopoldinia pulchra* Mart., *Manicaria saccifera* Gaertn., *Neonicholsonia watsonii* Dammer, *Pelagodoxa henryana* Becc., *Pholidostachys pulchra* H. Wendl., *Sommieria leucophylla* Becc.; as well as more distantly related taxa, *Podococcus barteri* G. Mann & H. Wendl., *Socratea exorrhiza* (Mart.) H. Wendl., and *Chamaerops humilis* L., all selected on the basis of a larger molecular data matrix (W. Baker et al., unpublished data.) and previous studies (Lewis and Doyle, 2002; Norup et al., 2003). We assigned *Chamaerops humilis* as the ultimate outgroup with which phylogenetic trees were rooted. Complete taxon, voucher, and distribution information (including GenBank accession numbers) is presented in Appendix.

Morphological characters—Ten macromorphological characters pertinent to the delimitation of *Heterospathe* and *Rhopaloblaste* were scored at the generic level across the entire taxon sample (Table 1). Characters were taken from descriptions in the literature (Moore et al., 1982; Uhl and Dransfield, 1987; Dowe and Uhl, 1989; Dransfield, 1991; Dransfield and Beentje, 1995; Dransfield and Uhl, 1998) and herbarium specimens, living specimens, and photographs at the Royal Botanic Gardens, Kew. These characters are described in the following section. In some cases, continuous characters could not be readily demarcated objectively (e.g., character 7, peduncular bract length, which is arbitrarily defined). However, these characters have been used in taxonomic descriptions of these taxa, and therefore it was appropriate to explore them here. Using MacClade, version 4.06 (Maddison and Maddison, 2003) character states were optimized onto one of the most parsimonious trees from the combined analysis.

Leaf character—1. Leaf sheath splitting: splitting to the base (0); remaining entire and tubular for at least three quarters until abscission or marcescence (1). Leaf sheath morphology plays a prominent role in the delimitation of palm genera. In subfamilies Arecoideae and Ceroxyloideae in particular, a distinction is made between species and genera with and without a crownshaft. A crownshaft is present when the leaf sheath is tubular and remains entire, only splitting slightly at the top until senescence, whereupon the sheath splits opposite the petiole along a predetermined line and abscises at the base. The alternative condition is more varied, but typically involves the leaf sheath splitting throughout its length and being open at maturity. A crownshaft is present in most Areceae genera and typically forms part of a broader suite of character states that presumably are functional consequences of strictly tubular leaf sheaths (character 1): inflorescence infrafoliar (character 2), peduncle shorter than the rachis (character 4), prophyll and peduncular bracts caducous (characters 5 and 6), and peduncular bract similar to prophyll or smaller (character 7). These states are not, however, logically dependent on the presence of tubular leaf sheaths, as indicated by the numerous exceptional taxa that do not conform to the typical character state suite.

Inflorescence characters—2. Inflorescence position: interfoliar (at least in bud) (0); infrafoliar (1). In palms, the most common position of the

inflorescence is interfoliar (between the leaves). Within the Areceae, inflorescence position is either interfoliar (as in *Heterospathe*) or infrafoliar (below the leaves) as in *Rhopaloblaste*. The infrafoliar condition most often occurs when the inflorescence in bud is exposed by the fall of the subtending leaf and is frequently associated with the presence of a crownshaft.

3. Inflorescence branches in bud: not convoluted (0); convoluted (1). Within the bud, inflorescence branches are typically packed in a linear fashion, but in a few rare cases (e.g., in *Rhopaloblaste*) in which the inflorescence appears to elongate before it emerges, branches become convoluted.

4. Peduncle length: shorter than rachis (0); longer than (or similar to) rachis (1). Peduncle length is often, but not always correlated with the position of the inflorescence. Thus it is typically elongate when the inflorescence is interfoliar (and must be exerted from the leaf sheaths), and usually short when the inflorescence is infrafoliar. Sometimes (e.g., in some species of *Dypsis* Noronha ex Mart.) elongate peduncles are found in combination with crownshafts (Dransfield and Beentje, 1995).

5. Prophyll persistence: persistent (0); caducous (1). The prophyll is the first bract on the primary inflorescence axis in palms. It is usually two-keeled and completely encloses the inflorescence during early development. Generally, prophylls of Areceae members abscise at anthesis. However, in a few genera the prophyll persists, e.g., in *Brongniartikentia* Becc., *Heterospathe*, and *Alsmithia*.

6. Peduncular bract persistence: persistent (0); caducous (1). Peduncular bracts are sterile bracts borne above the prophyll, and in most members of Areceae they share a role in protection of the inflorescence in early stages. As with prophylls, peduncular bracts in Areceae members most often abscise at anthesis. However, sometimes the peduncular bract persists, e.g., in species of *Heterospathe* and *Brongniartikentia*.

7. Peduncular bract length: longer than prophyll (0); similar to prophyll (or smaller) (1). In most members of tribe Areceae the peduncular bract is enlarged (compared to most other palms) and equals the prophyll, together the two bracts enclosing the inflorescence in bud. In some genera of Areceae, e.g., *Heterospathe* and *Alsmithia*, the peduncular bract is much longer than the prophyll.

Fruit and seed characters—8. Endocarp sculpturing: none (0); conspicuous flanges, grooves, ridges (1). Generally the endocarps of Areceae are thin and lack ornamentation. In genera such as *Alsmithia*, *Burretioakentia* Pic. Serm., and *Veillonina* H. E. Moore, however, the endocarp is thickened and intricately sculptured. Although this character is not relevant to the current circumscriptions of *Heterospathe* and *Rhopaloblaste*, preliminary analyses of our DNA data sets indicated that it may be of great significance to a revised generic concept in *Heterospathe*.

9. Endosperm type: homogeneous (0); ruminant (1). Endosperm condition has been considered to have considerable diagnostic value. Palm seeds contain abundant endosperm that can be homogeneous or ruminant (i.e., with folds and invaginations from the seed coat). Rumination can be very variable and, in some cases, polymorphic within species (Zona, 2003).

Seedling character—10. Eophyll shape: bifid (0); pinnate (1); entire (2). The eophyll is the first expanded leaf produced by a seedling and is in its simplest form undivided. However, in palms with reduplicate leaves (such as members of the Areceae) a bifid eophyll is the most common form. Pinnate eophylls occur in, e.g., *Rhopaloblaste* and *Acanthophoenix* H. Wendl.; in the latter, bifid eophylls are also known in some species.

Three characters states, which according to Banka and Baker (2004) are useful in delimitation of *Rhopaloblaste*, were not incorporated in this analysis, namely, strongly divaricate first-order inflorescence branches, a club-shaped embryo, and an indumentum on the leaf petiole and rachis consisting of multiple, laminar, membranous brownish-black scales. They were excluded due to character-coding difficulties and lack of adequate comparative data across our samples.

DNA extractions, amplification, and sequencing—DNA was extracted using 0.3 g silica-gel-dried (Chase and Hills, 1991) or 1.0 g fresh material and a version of the 2 × CTAB extraction method modified from that of Doyle and Doyle (1987). Before precipitation, a sample was purified with the Qiagen PCR purification kit (Qiagen, Crawley, West Sussex, UK) following the manufacturer's protocol; this was used immediately for PCR. The remaining DNA was then purified on cesium chloride/ethidium bromide gradients (1.55 g · L⁻¹ density) and dialyzed. These DNA samples are available from the DNA Bank of the Royal Botanic Gardens, Kew (<http://www.rbgekew.org.uk/data/dnaBank>).

TABLE 1. Matrix of morphological character states for tribe Areceae and outgroups. Description of characters and states are given in Materials and Methods, subsection Morphological characters. ? = missing data; - = state not applicable; nos. surrounded by brackets signify polymorphic state assignments. Taxa are followed by a voucher number if there is more than one collection per species or if the species is undetermined.

Taxa	Character states for characters 1–10
<i>Acanthophoenix rubra</i>	110011100{01}
<i>Actinokentia huerlimannii</i>	110011100?
<i>Actinorhytis calapparia</i>	1100111010
<i>Adonidia merrillii</i>	1100111010
<i>Alloschmidia glabrata</i>	1000111000
<i>Alsmithia longipes</i> Baker 1180	0001000100
<i>Alsmithia longipes</i> Zona 1039	0001000100
<i>Archontophoenix purpurea</i>	1100111010
<i>Areca catechu</i>	11001–010
<i>Asterogyne martiana</i>	0001010000
<i>Balaka longirostris</i>	1101110000
<i>Basselintia velutina</i>	1100111000
<i>Bentinckia condapanna</i>	1100111100
<i>Brassiophoenix drymophloeoides</i>	1100111100
<i>Brongniartikentia lanuginosa</i>	1{01}01110000
<i>Burriokentia vieillardii</i>	1100111100
<i>Calyptrocalyx hollrungii</i>	000–000000
<i>Calyptrogyne costatifrons</i>	000–010000
<i>Campecarpus fulcitus</i>	1100111100
<i>Carpentaria acuminata</i>	1100111000
<i>Carposylon macrospermum</i>	1100111000
<i>Chamaerops humilis</i>	00000–012
<i>Chambeyronia macrocarpa</i>	1100111000
<i>Clinosperma bracteale</i>	110011100?
<i>Clinostigma savoryanum</i>	1100111000
<i>Cyphokentia macrostachya</i>	110011100?
<i>Cyphophoenix nucele</i>	1100111000
<i>Cyphosperma balansae</i>	0001000100
<i>Cyrtostachys renda</i>	1100111000
<i>Deckenia nobilis</i>	1100111000
<i>Dictyosperma album</i>	1100111010
<i>Drymophloeus litigiousus</i>	1100111010
<i>Dypsis lutescens</i>	1{01}01011000
<i>Geonoma deversa</i>	0{01}10?{01}1000
<i>Hedyscepe canterburyana</i>	1100111000
<i>Heterospathe cagayanensis</i>	0001000010
<i>Heterospathe delicatula</i>	0001000010
<i>Heterospathe elata</i> Houg. & Hubb. 1024	0001000010
<i>Heterospathe elata</i> Chapin 55	0001000010
<i>Heterospathe elata</i> Lewis 99-034	0001000010
<i>Heterospathe humilis</i>	0001000010
<i>Heterospathe macgregorii</i>	0001000010
<i>Heterospathe philippinensis</i>	0001000010
<i>Heterospathe phillipsii</i>	0001000010
<i>Heterospathe scitula</i>	0001000010
<i>Heterospathe sibuyanensis</i>	0001000010
<i>Heterospathe</i> sp. Baker 1115	0001000010
<i>Heterospathe</i> sp. Baker 1142	0001000010
<i>Heterospathe</i> sp. Banka 2008	0001000010
<i>Heterospathe</i> sp. nov. Fernando 1624	0001000010
<i>Howea belmoreana</i>	000–000000
<i>Hydriastele beguinii</i>	1100111010
<i>Hydriastele brassii</i>	1100111010
<i>Hydriastele costata</i>	1100111000
<i>Hydriastele microspadix</i>	1100111010
<i>Iguanura wallichiana</i>	0001000010
<i>Kentopsis oliviformis</i>	1100111000
<i>Laccospadix australasicus</i>	000–000010
<i>Lavoixia macrocarpa</i>	110011100?
<i>Lemurophoenix hallexii</i>	1100111010
<i>Leopoldinia pulchra</i>	0001111000
<i>Lepidorrhachis mooreana</i>	0100111000

TABLE 1. Continued.

Taxa	Character states for characters 1–10
<i>Linospadix albertisiana</i>	000–010000
<i>Loxococcus rupicola</i>	1100111010
<i>Manicaria saccifera</i>	0001001000
<i>Marojejya darianii</i>	0001001100
<i>Masoala madagascariensis</i>	0001001000
<i>Moratia cerifera</i>	1100111000
<i>Nenga pumila</i> var. <i>pachystachya</i>	11001–010
<i>Neonicholsonia watsonii</i>	000–000010
<i>Neoveitchia storckii</i>	1100110000
<i>Nephrosperma vanhoutteanum</i>	0001010010
<i>Normanbya normanbyi</i>	1100111010
<i>Oncosperma tigillarum</i>	1100111010
<i>Pelagodoxa henryana</i>	0001010000
<i>Phoenicophorium borsigianum</i>	0001000012
<i>Pholidostachys pulchra</i>	000–{01}10000
<i>Physokentia rosea</i>	1100111100
<i>Pinanga coronata</i>	11001–010
<i>Podococcus barteri</i>	0001001001
<i>Ponapea palauensis</i>	1100111100
<i>Ptychococcus paradoxus</i>	11001111{01}0
<i>Ptychosperma micranthum</i>	11010{01}1010
<i>Ptychosperma salomonense</i>	1100111000
<i>Rhopaloblaste augusta</i>	1110111011
<i>Rhopaloblaste ceramica</i>	1110111011
<i>Rhopaloblaste ledermanniana</i> Heatubun 191	1110111011
<i>Rhopaloblaste ledermanniana</i> Maturbongs 650	1110111011
<i>Rhopaloblaste singaporensis</i>	1120111011
<i>Rhopalostylis baueri</i>	1100111000
<i>Roscheria melanochaetes</i>	1{01}01000010
<i>Satakentia liukuensis</i>	1100111000
<i>Socratea exorrhiza</i>	1101110000
<i>Solfia samoensis</i>	1101111000
<i>Sommieria leucophylla</i>	0001001000
<i>Tectiphiala ferox</i>	110011100?
<i>Veillonina alba</i>	110011010?
<i>Veitchia spiralis</i>	1100111000
<i>Verschaffeltia splendida</i>	0001010110
<i>Wodyetia bifurcata</i>	1100111000

Primers for amplification of *PRK* (entire intron 4, partial exons 4 and 5) followed Lewis and Doyle (2002); 717F (5'-GTGATATGGAAGAACGTGG-3') and 969R (5'-ATTCCAGGGTATGAGCAGC-3') are designed to target only paralogue 2 within *PRK*. Lewis and Martínéz (2000) found two paralogues of *PRK* in a species-level analysis and reported paralogue 2 to be informative. This paralogue was also employed in the study of Lewis and Doyle (2002). Primers for *RPB2* intron 23 followed Roncal et al. (2005): INT23F (5'-CAACTTATT GAGTGCATCATGG-3') and INT23R (5'-CCACGCATCT-GATATCCAC-3'). For PCR amplification, a high-magnesium PCR buffer was used for both *RPB2* and *PRK* (1.1× ReddyMix PCR Master Mix, 2.5mM MgCl₂). Reactions included the following: 1–2 µL of DNA template, 45 µL of 1.1× ReddyMix (Abgene, Epsom, UK), 0.5 µL of both forward and reverse primers, and 1 µL of bovine serum albumin (BSA). For the amplification of *PRK*, 1 µL of 2% dimethylsulfoxide (DMSO) was added to improve amplification success.

Thermocycler conditions employed for *PRK* amplification were an initial cycle at 94°C for 4 mins, 32 cycles of 94°C for 1 min, 56.5°C for 1 min, and 72°C for 2 mins. Following the last cycle, we used a final extension at 72°C for 7 mins. Amplification of *RPB2* was conducted using conditions similar to the above, except for an annealing temperature of 54.5°C. Some taxa that did not readily amplify were run with a lower annealing temperature (52°C for *PRK*, 53°C for *RPB2*). All PCR and cycle sequencing reactions were run on a Perkin-Elmer GenAMP model 9600 or 9700 (Perkin-Elmer, Boston, Massachusetts, USA). Following amplification, samples were purified using Qiagen PCR purification kit (Qiagen) following the manufacturer's protocol. Purified DNA fragments were sequenced using the PCR primers and the BigDye Terminator Mix v. 3.0 (Applied Biosystems, ABI, Warrington, Cheshire, UK) following the manufacturers protocol. The sequencing protocol consisted of 26 cycles of

10 sec denaturation (96°C), 5 sec annealing (50°C), and 4 min elongation (60°C). The cleaned cycle-sequencing products were analyzed on an ABI 377 automated sequencer, according to the manufacturer's protocols. Both strands were sequenced for each region for all taxa.

In six taxa, two for *PRK* and four for *RPB2*, amplification results were poor or yielded electropherograms that indicated the presence of multiple copies of the target sequence. Such PCR products were cloned using pGEM-T Easy Vector System (Promega, Chilworth Science Park, Southampton, UK) following the manufacturer's protocol. Transformed cells were grown on solid LB medium, and five transformed colonies per cloned taxon were sampled for PCR by inoculating PCR reactions with transformed cells as template.

Alignment and phylogenetic analysis—Sequences were edited and assembled using Sequencer 4.1.2 software (Gene Codes, Ann Arbor, Michigan, USA). All sequences, including some previously published by Lewis and Doyle (2002), were aligned by eye in PAUP, version 4.0b10 for Macintosh (Swofford, 2002). Ambiguous parts of the alignment (e.g., some rather large homopolymer regions) and parsimony uninformative characters were excluded from the analysis. Matrices can be downloaded from TreeBase (<http://www.treebase.org>). Cladistic analyses were conducted with PAUP* using maximum parsimony. All character changes were treated as unordered and equally weighted (Fitch, 1971). Indels were treated as missing data.

The data partitions were analyzed separately and simultaneously. Preliminary searches yielded extremely large numbers of equally parsimonious resolutions, a problem exacerbated by the presence of zero length branches. These searches could not, in practice, be run to completion. Thus, a pragmatic strategy similar to that used by Asmussen and Chase (2001) was followed: 1000 replicates of random taxon addition were performed using tree-bisection-reconnection (TBR) branch swapping with 25 trees held at each step; trees found in these 1000 replicates were then used as starting trees for a second search using TBR branch swapping until a pre-set maximum of 20 000 trees were reached. These trees were then swapped to completion. Support for clades was assessed using the bootstrap (Felsenstein, 1985) as implemented in PAUP* using 500 replicates of simple taxon addition, TBR swapping, and a limit of 10 trees saved in each replicate (Salamin et al., 2003).

The *PRK* and *RPB2* data sets were examined for congruence using the partition homogeneity test implemented in PAUP* (Farris et al., 1995; 100 replicates, saving 10 trees per replicate). However, because this method has been shown to be unreliable (Dolphin et al., 2000; Reeves et al., 2001; Yoder et al., 2001), a "hard" incongruence test was also performed by directly comparing topologies from the separate analyses with respect to resolution and bootstrap support (i.e., looking for incongruent clades with bootstrap percentages (BP) >85; Wiens, 1998; Sheahan and Chase, 2000).

RESULTS

***PRK* and *RPB2* Clones**—Five or more clones were sequenced for two of the six taxa for which heterogeneity was observed, namely *Brongniartikentia lanuginosa* H. E. Moore and *Veillonina alba* H. E. Moore for *PRK*. Comparisons between the clones revealed that, although different versions of the target sequence occurred within species, they differed in only a very small number of substitutions and a few short indels; these most likely represented alleles or recent paralogy because they all resolved as monophyletic groups in preliminary analyses. For the purposes of this study, therefore, one clone was selected at random for inclusion in the analyses. For the remaining four taxa for *RPB2*, *Cyphokentia macrostachya* Brongn., *Heterospatha phillipsii* D. Fuller & Dowe, *Marojejya darianii* J. Dransf. & N. Uhl and *Roscheria melanochaetes* H. Wendl., only one clone was obtained.

***PRK* analysis**—The final alignment included 800 positions; 119 positions were excluded as ambiguously aligned because they consisted of microsatellite motifs or long homopolymer regions. Of the remaining characters, 206 (30.2%) were potentially parsimony informative (Table 2). In *Heterospatha*,

TABLE 2: Statistics from phylogenetic analyses of *PRK*, *RPB2*, and combined matrices for tribe Areceae and outgroups. CI = consistency index; RI = retention index.

DNA region	<i>PRK</i>	<i>RPB2</i>	Combined
No. of taxa	101	100	100
Aligned length	800	1301	2101
No. potentially informative characters	206 (30.2%)	257 (22.3%)	463 (25.2%)
No. trees	20 000	20 000	20 000
Tree length	581	674	1291
CI	0.51	0.58	0.53
RI	0.71	0.71	0.69
No. of clades with ≥85% bootstrap	19	26	38
No. of clades with ≥50% bootstrap	44	47	62

the *PRK* intron was almost consistently about half the length of the region in other palms. Although this feature represents a character linking the species of *Heterospatha* together, it is potentially problematic because it reduces information available for the group. Analyses produced 20 000 trees (the pre-set limit) with a length of 581, a CI of 0.51 and a RI of 0.71. The strict consensus tree with bootstrap percentages is shown in Fig. 1.

The Areceae (99 bootstrap percentage, BP), outgroup tribe Geonomateae (*Asterogyne* H. Wendl., *Geonoma* Willd., *Calyptogyne* H. Wendl., *Pholidostachys* H. Wendl.; 87 BP), the New World tribes Leopoldinieae (*Leopoldinia*), Manicarieae (*Manicaria*) and Euterpeae (*Neonicholsonia*), and the Pacific tribe Pelagodoxeae (*Pelagodoxa*, *Sommieria*; 100 BP) form a moderately supported clade (78 BP), but the relationships among these tribes are unclear. The topology supports the exclusion of Pelagodoxeae from Areceae.

Within the Areceae, a substantial clade consisting mainly of western Pacific and Australian genera from subtribes Archontophoenicinae, Basseliniinae, Carpoxylinae, Clinospermatinae, Linospadicinae, Ptychospermatinae, and Rhopalostylidinae, as well as several genera as yet unplaced to subtribe, is resolved, corresponding to the south-western Pacific and Australia clade of Lewis and Doyle (2002). This clade is hereafter termed the western Pacific clade. We recover only weak support for the group here (61 BP), and within it there is a large polytomy. One Indian Ocean genus, *Loxococcus* H. Wendl. & Drude from Sri Lanka, is included at the basal polytomy of the western Pacific clade. Subtribe Ptychospermatinae excluding *Ptychosperma micranthum* is weakly supported (clade wp3; 60 BP). *Ptychosperma micranthum* falls at the basal polytomy of the western Pacific clade. The monophyly of two other western Pacific subtribes Clinospermatinae (clade wp4; 98 BP) and Carpoxylinae (clade wp6; <50 BP) is also resolved. Subclades of a number of other subtribes are resolved, such as a group consisting of all members of subtribe Linospadicinae except *Calyptrocalyx* Blume (clade wp5; 75 BP).

Within the large western Pacific clade, all members of *Heterospatha* form a monophyletic group (72 BP) with the monotypic genus *Alsmithia* from Fiji, within which there is a basal polytomy such that the monophyly of neither genus is confirmed. Thus, the Philippine endemics *H. cagayanensis* Becc. and *H. sibuyanensis* Becc. are supported by 83 BP, in a group (<50 BP) otherwise consisting of the widespread *H. elata* Scheff. and an undescribed Philippine species (*Fernando*

Fig. 1. Strict consensus tree resulting from parsimony analysis of the *PRK* data set for tribe Areceae and outgroups. Bootstrap percentages >50% are indicated above the branches. Herbarium vouchers are indicated for clarity where necessary. Significant clades that appear in the combined analysis are marked with triangles. Higher level ranks follow the classification of Dransfield et al. (2005). Genera of Areceae that are not yet placed to subtribe by Dransfield et al. (2005) are indicated.

1624). A clade (100 BP) of the New Guinean species *H. delicatula* H. E. Moore and *H. humilis* Becc. is resolved within a clade (<50 BP) containing the Philippine species, *H. philippinensis* Becc. and *H. scitula* Fernando plus the Fijian *H. phillipsii*.

Rhopaloblaste is a strongly supported monophyletic group (97 BP) outside the western Pacific clade. *Rhopaloblaste ceramica* (Miq.) Burret, *R. ledermanniana* Becc. (*Heatubun* 191), and *R. singaporensis* (Becc.) Hook. f. are supported by 92 BP. The second accession of *R. ledermanniana* (*Maturbongs* 650) plus *R. augusta* (Kurz) H. E. Moore are not clearly resolved.

Most genera that resolve outside the western Pacific clade along with *Rhopaloblaste* are distributed in the Indian Ocean through to Malesia, and belong to subtribes Arecinae, Dypsidinae, Oncospermatinae, and Verschaffeltiinae or are as yet unplaced to subtribe. Of these subtribes, only the Arecinae is resolved as monophyletic here (69 BP).

RPB2 analysis—The final alignment had a total length of 1301 bp; 148 characters were excluded as ambiguously aligned due to microsatellite motifs or long homopolymer regions. Of the remaining positions, 257 (22.3%) were potentially parsimony informative (Table 2). Analysis produced 20 000 trees (the pre-set limit) with a length of 674, a CI of 0.58 and a RI of 0.71. The strict consensus tree with bootstrap percentages is shown in Fig. 2. No *RPB2* sequence was produced for *Heterospathe* sp. *Baker* 1130, and hence this analysis includes only 15 accessions for the genus.

Outgroup taxa are similarly arranged as in the *PRK* trees, but with slightly better resolution, although only a few groups are well-supported. Again, the topology supports the exclusion of Pelagodoxeae from Areceae. The Areceae is strongly supported (95 BP), although relationships within the clade are highly unresolved. The western Pacific clade is not recovered in this topology and only two subtribes, Clinospermatinae (clade wp4; 66 BP) and Carpoxylinae (clade wp6; 88 BP), are resolved as monophyletic. However, numerous subclades from various subtribes are resolved.

All species of *Heterospathe* form a clade, but it is weakly supported (<50 BP). There are within this clade two strongly supported subclades (1 and 2; 99, 98 BP, respectively). Clade 1 consists mainly of Philippine and Fijian species plus two species from New Guinea (*H. delicatula* and *H. humilis*). The other clade (2) contains New Guinean species only. The Fijian *Alsmithia* is, unlike in the *PRK* analysis, clearly nested within *Heterospathe*, in this case within the strongly supported clade 1. Also within clade 1, a weakly supported group (64 BP) contains all Philippine species plus the widespread *H. elata*. The New Guinean species, *H. humilis* and *H. delicatula*, form a strongly supported pair (94 BP).

Rhopaloblaste is again robustly resolved as monophyletic group (100 BP), within which *R. singaporensis* and *R. ledermanniana* (*Heatubun* 191) form a strongly supported pair (99 BP). The second accession of *R. ledermanniana* (*Maturbongs* 650) is resolved in an ambiguous position within the clade. A relationship not seen in the *PRK* tree is the strongly supported sister group relationship of *Rhopaloblaste* with the monotypic Mascarene genus, *Dictyosperma* H. Wendl. & Drude (96 BP).

Combined analysis—According to the partition homogeneity test, the two data sets were significantly incongruent ($P = 0.01$). However, comparing the single-gene topologies on

a node-by-node basis, which may give a better assessment of congruence than the sometimes unreliable partition homogeneity test (Dolphin et al., 2000; Reeves et al., 2001; Yoder et al., 2001), did not result in any highly supported, incongruent relationships (>85 BP). We have therefore chosen to combine the two datasets due to the generally high level of congruence between the single-gene trees. The increased number of well-supported clades resulting from analysis of the combined dataset relative to the separate data partitions (Table 2) lends further weight to our assertion that the two datasets are broadly congruent.

Due to the lack of an *RPB2* sequence for *Heterospathe* sp. *Baker* 1130, this accession was left out of the combined analysis. Analysis of the combined matrix produced 20 000 trees with a length of 1291, a CI of 0.53 and a RI of 0.69 (Table 2). One of the most parsimonious trees is shown in Fig. 3.

The combination of the two datasets resulted in a substantial increase in both resolution and numbers of nodes supported by the bootstrap (Table 2), especially within the Areceae. As found in the *RPB2* analysis, outgroup relations are well resolved. The topology moderately supports a large clade (90 BP) broadly comparable to relationships recovered by Lewis and Doyle (2002) composed of the following groups: Areceae (100 BP), Euterpeae, Geonomateae (100 BP), Leopoldinieae and Pelagodoxeae (100 BP). The New World tribe Manicarieae is strongly supported as sister to this clade (92 BP). The combined analysis supports the exclusion of Pelagodoxeae from Areceae, as did the individual data partitions.

The clade of Areceae is divided into two major subclades (Fig. 3), although neither is well-supported. The first clade (<50 BP) comprises genera occurring in the islands and continental regions of the Indian Ocean through Malesia to New Guinea, a few of which also extend into the western Pacific (*Cyrtostachys*, *Hydriastele*, *Rhopaloblaste*, and *Clino-stigma* H. Wendl.) and is hereafter referred to as the Indian Ocean clade. The second clade (the western Pacific clade; 63 BP) consists of genera from the Philippines, the Moluccas, and New Guinea through to Australia, New Caledonia, and the Areceae's easternmost limit in Samoa. The monotypic genus *Loxococcus* from Sri Lanka is supported as sister to the western Pacific clade (66 BP). Relationships within these two major clades are poorly resolved, but include several major groups of interest.

Within the Indian Ocean clade, only the Arecinae is resolved as monophyletic (87 BP). Three of the four genera of subtribe Dypsidinae resolve in a clade with *Iguanura* (clade io4; <50 BP). Subclades of subtribes Verschaffeltiinae and Oncospermatinae form a group (io1; <50 BP), with other constituent genera forming relationships elsewhere. Although these results call into question the monophyly of Dypsidinae, Oncospermatinae and Verschaffeltiinae, none of the relevant relationships is well supported.

Within the western Pacific clade, four of the five genera of Archontophoenicinae (*Actinokentia*, *Archontophoenix*, *Chambeyronia*, *Kentiopsis*) constitute a strongly supported group (100 BP) within clade wp2. Most members of subtribe Linospadicinae (excluding *Calypstrocalyx*) form a strongly supported clade (clade wp5; 100 BP). Subtribes Carpoxylinae (clade wp6; 96 BP) and Clinospermatinae (clade wp4; 100 BP) are strongly supported as monophyletic. Ptychospermatinae form a clade (clade wp3; 72 BP) that excludes *Ptychosperma micranthum*, which resolves as sister to clade wp4. Subclades of Basseliniinae receive bootstrap support within clades wp1

Fig. 2. Strict consensus tree resulting from parsimony analysis of the *RPB2* data set for tribe Areceae and outgroups. Bootstrap percentages >50% are indicated above the branches. Within *Heterospathe*, clade 1 comprises mainly Philippine and Fijian species and clade 2 consists of New Guinean species. See Results: *RPB2* analysis for further information. See Fig. 1 legend for an explanation of annotations.

Fig. 3. One of the most parsimonious trees from the combined analysis of PRK and RPB2 data for tribe Areceae and outgroups (length = 1291; CI = 0.53; RI = 0.69). Branch lengths under DELTRAN optimization are indicated above the branches. Bootstrap percentages >50% are indicated below the branches. Branches that collapse in the strict consensus tree are indicated by arrows. See Fig. 1 legend for an explanation of annotations.

(60 BP) and wp7 (98 BP), but the subtribe is not resolved as monophyletic.

All included members of *Heterospatha* resolve within the western Pacific clade as a strongly supported clade (94 BP) with the monotypic genus *Alsmithia* firmly nested inside. Within the *Heterospatha-Alsmithia* group, two clades with moderate to strong bootstrap support are found, consisting of (1) species occurring from the Philippines through to Fiji (clade 1, 100 BP) and (2) only species from New Guinea (clade 2, 95 BP). In clade 1, the widespread *H. elata* is resolved within a strongly supported clade (99 BP) also containing a group consisting of the two Philippine endemics, *H. cagayanensis* and *H. sibuyanensis* (99 BP). The two New Guinean species *H. delicatula* and *H. humilis* are resolved within clade 1 supported by 100 BP.

The five species of *Rhopaloblaste* are resolved within the Indian Ocean clade supported by 100 BP and with a strongly supported relationship (89 BP) to the monotypic Mascarene genus *Dictyosperma*. Complete resolution in this part of the consensus tree shows the first split to be between *Rhopaloblaste augusta* from Nicobar Island and a moderately supported group (72 BP) consisting of *R. ledermanniana* and *R. ceramica* (New Guinea and the Moluccas), plus *R. singaporensis* (peninsular Malaysia and Singapore). Within this group, *R. singaporensis* and *R. ledermanniana* (Heatubun 191) are strongly supported (100 BP) as sisters, with a well-supported relationship to *R. ceramica*. *Rhopaloblaste ledermanniana* (Maturbongs 650) is resolved separately from the other accession of *R. ledermanniana*.

DISCUSSION

Phylogenetic value of *PRK* and *RPB2*—The level of variation in the low-copy nuclear genes *PRK* and *RPB2* is significantly higher than that of the plastid genes used so far in palm phylogenetic studies. For example, the number of informative characters obtained for three plastid markers in a family-wide survey of palms ranged from 6–10% only (Asmussen and Chase, 2001), whereas *PRK* and *RPB2* yield 30.2% and 22.3%, respectively, in our study, despite the narrower taxonomic range of our sampling. However, consistency and retention indices are still relatively low, indicating an elevated number of homoplasious characters. Introns of plastid DNA seem to contain the least homoplasy reported for regions used in analyses of the palm family (e.g., for 94 taxa, RI = 0.89 for the *rps16* intron; Asmussen and Chase, 2001), but as mentioned above, they also contain a lower percentage of parsimony informative characters. In contrast, other nuclear ribosomal DNA regions like ITS and 5S spacer contain a high number of informative characters, but also high levels of homoplasy. In recent studies by Baker et al. (2000b, c) on the palm subfamily Calamoideae, values of RI varied between 0.39 (60 taxa, 415 informative characters) and 0.62 (108 taxa, 469 informative characters) for ITS, and between 0.48 (41 taxa, 267 informative characters) and 0.66 (77 taxa, 352 informative characters) for 5S. High levels of polymorphism within individuals were also discovered in these two multi-copy regions, rendering them relatively unappealing for phylogenetic reconstruction.

It is recognized that even if the partition homogeneity is rejected, combining data sets can still give more accurate phylogenetic estimates (Huelsenbeck et al., 1996; Wiens,

1998). The difference in the phylogenetic histories can potentially affect parts of the combined tree, but the overall accuracy may be increased by the larger number of characters applied to those parts of the tree unaffected by the mismatch (Wiens, 1998). We regard the fact that the combination of the two datasets resulted in a substantial increase in both resolution and numbers of nodes supported by the bootstrap as strong support for simultaneous analysis. It suggests that the significant incongruence identified by the partition homogeneity test is spurious and potentially a result of high levels of noise in the dataset (Dolphin et al., 2000).

This study has identified many new relationships within the Areceae and provided a more resolved and strongly supported estimate of the phylogeny relative to previous analyses that have included genera of Areceae (e.g., Asmussen and Chase, 2001; Lewis and Doyle, 2002; Asmussen et al., in press). It has also demonstrated that *PRK* and *RPB2* are capable of resolving relationships at high taxonomic levels (i.e., between tribes and subtribes) as well as at lower levels between closely related species. Unfortunately, a lack of resolution and bootstrap support leaves large areas of ambiguity in the tree that will have to be addressed before the classification, relationships, and evolutionary history of the Areceae can be fully appreciated.

Monophyly of *Heterospatha* and *Rhopaloblaste*—Our results demonstrate that the species of *Rhopaloblaste* form a strongly supported clade and are clearly not related to *Heterospatha*. The generic delimitations of Moore (1970) and Uhl and Dransfield (1987) are thus largely confirmed and well-established, despite the previous taxonomic confusion.

The species of *Heterospatha* are resolved as a strongly supported monophyletic group that also includes the monotypic Fijian genus *Alsmithia*. Several recent studies have indicated that some changes in generic delimitation will be necessary within tribe Areceae (Lewis and Doyle, 2002; Baker and Loo, 2004). In our study, the close relationship between the two genera is well-supported by the DNA analyses, not only by the combined dataset, but also by both datasets individually. The possibility of this relationship being the result of a technical error was ruled out by inclusion of two independent collections of *Alsmithia*.

Philippine species of *Heterospatha*—A full monograph of *Heterospatha* has yet to be undertaken and so far only the Philippine species have been revised. In other parts of its range (e.g., Papuasia), species limits are not clear, and relationships within the genus as a whole are not yet assessed. The results of our molecular analyses are congruent with the recent revision of *Heterospatha* from the Philippines (Fernando, 1990a, b). Fernando noted that the widespread *H. elata* (occurring in the Philippines, the Moluccas, Guam, and Palau Island) is morphologically most similar to the island endemics *H. sibuyanensis* and *H. cagayanensis*, due to their shared robust habits and large inflorescences; this relationship is reflected in our combined analysis (99 BP). Differences between *H. elata* and *H. sibuyanensis* are limited, and only the shape of fruits and seeds seem to distinguish these two species. In his revision, Fernando (1990b) noted that *H. sibuyanensis* is known only because of the absence of more herbarium material from the type locality. However, we question the identity of the DNA source, a cultivated palm originating far from the type

locality in Quezon Province. Unfortunately, the lack of fertile material precludes confirmation at this time, although sterile material closely resembles *H. elata*. Fernando furthermore described the Luzon endemic *H. scitula* as having the greatest resemblance to *H. philippinensis* (and *H. dransfieldii* Fernando, not included in this analysis), but differing in its neatly abscising leaves and shorter leaflet pairs and a seemingly infrapolar inflorescence. This relationship is supported by 91 BP in our combined analysis.

Relationships between *Heterospathe*, *Rhopaloblaste*, and other members of the Areceae—The monotypic Mascarene genus *Dictyosperma* is resolved with strong support as the nearest relative to *Rhopaloblaste* (89 BP). Uhl and Dransfield (1987) regarded the relationships of *Dictyosperma* as unclear, but noted that it was possibly related to *Clinostigma*, which is not supported by our analyses. Geographically, *Rhopaloblaste* and *Dictyosperma* are far separated, but morphological similarities between *Dictyosperma* and *Rhopaloblaste* do exist, e.g., the primary inflorescence branches of *Dictyosperma* become divaricate with age and are thus reminiscent of those in *Rhopaloblaste*. No such close relationship between *Heterospathe* and any of the other genera, apart from *Alsmithia*, is identified.

Higher-level relationships within the Areceae—The high support for the Areceae corroborates previous results (Hahn, 2002; Lewis and Doyle, 2002) and supports the exclusion of Pelagodoxeae from the tribe. Our study highlights some of the limitations of the previous classification (Uhl and Dransfield, 1987; Dransfield and Uhl, 1998), such as the non-monophyly of the large subtribe Iguanurinae, which is not accepted in the revised classification of Dransfield et al. (2005). It provides support for the delimitation of several subtribes recognized in the revised classification, principally Arecinae, Carpoxylinae, Clinospermatinae and Ptychospermatinae. Though the monophyly of the remaining subtribes of Areceae (Archontophoenicinae, Basseliniinae, Dyspidinae, Oncospermatinae, Rhopalostylidinae, and Verschaffeltiinae) is not supported, neither is strong support for their non-monophyly obtained. The sole exception to this generality is the curious relationship between *Hedyscepe* and the clade *Alloschmidia* and *Basselinia*, which is highly supported by the combined analysis (88 BP) and thus contradicts the monophyly of the Rhopalostylidinae.

Morphological characters—All morphological characters included in the study were homoplasious (Appendix S1; see Supplementary Data accompanying online version of this article). Such high levels of homoplasy have been reported several times for large and species-rich plant groups in the tropics (e.g., Moylan et al., 2004; Tam et al., 2004). Within the palm family, a recent study of subfamily Calamoideae found many of the characters previously used to delimit subtribes and genera of Calamoideae to be highly homoplasious (Baker et al., 2000a). Other palm genera that have been previously reported to be characterized by a combination of characters that are homoplasious include *Sommieria* and *Pelagodoxa* (Stauffer et al., 2004) and *Ptychococcus* Becc. (Zona, 2003).

Of the suite of characters associated with the crownshaft, only the presence of tubular leaf sheaths is unequivocally synapomorphic for the Areceae. Three of the remaining characters (infrapolar inflorescences, caducous prophylls, and peduncles shorter than the rachis) optimize equivocally at the

Areceae node, though are potentially synapomorphic for the group under some alternative optimizations (Appendix S1; see Supplementary Data accompanying online version of this article). However, these crownshaft characters are highly homoplasious with several occurrences outside the Areceae (e.g., Chamaedoreae, Cyclospatheae, Iriarteeae, Roystoneae) and numerous correlated reversals, resulting in a substantial number of genera without crownshafts (leaf sheath splitting to the base) that also bear interfoliar inflorescences with persistent or marcescent prophylls and peduncles that greatly exceed the rachis (*Alsmithia*, one species of *Brongniartikentia*, *Cyphosperma*, *Heterospathe*, *Iguanura*, *Nephrosperma*, *Phoenicophorium*, *Verschaffeltia*, some species of *Dypsis*, *Masoala*, *Marojejya* Humbert, *Calyptracalyx*, *Howea*, *Laccospadix*, and *Linospadix*; character four is inapplicable to last four genera). Our findings must be interpreted within the context of our choice of outgroups; a different selection (e.g., replacing *Neonicholsonia* with a crownshafted member of Euterpeae) may have yielded alternative optimizations at the Areceae node.

The remaining character states linked to the crownshaft (peduncular bract caducous; peduncular bract similar in length to prophyll or shorter) are not synapomorphic for the Areceae. Both states are widespread across our sample and plesiomorphic in the Areceae, suggesting that they may not be entirely dependent on the occurrence of a crownshaft as suspected initially. Although neither character provides a synapomorphy for the Areceae, they are both clearly closely linked to transformations in other members of the crownshaft character suite. Thus, we have demonstrated a high degree of correlation among the states that comprise the complex morphologies associated with the crownshaft. Such a strong correlation reflects the functional interdependence of these states. For example, the maturation of an inflorescence is hindered by a strictly tubular leaf sheath and can only proceed to anthesis on abscission of the subtending leaf; the inflorescence is thus exposed and presented in an infrapolar position. Although exceptions occur, e.g., in *Dypsis lutescens*, this correlation holds true in the majority of cases.

This study indicates that *Heterospathe*, including *Alsmithia*, can be diagnosed by a combination of character states that are homoplasious within the Areceae: leaf sheath splitting to the base, interfoliar inflorescences (at least at anthesis), peduncle longer than the rachis, peduncular bract longer than the prophyll, prophyll persistent, and peduncular bracts persistent. It is remarkable that the monophyly of a genus with such ambiguous morphology is so highly supported by molecular data. This combination does not preclude confusion with a small number of other genera of Areceae, namely *Brongniartikentia* [*B. vaginata* (Brongn.) Becc. only], *Cyphosperma*, *Iguanura*, and *Phoenicophorium*. However, these genera are readily distinguished on the basis of characters outside our data set, for example, didymous anthers in *Brongniartikentia* (which also has a crownshaft in *B. lanuginosa* and some individuals of *B. vaginata*; Pintaud and Hodel, 1998), an incomplete prophyll in *Cyphosperma*, praemorse leaflets in *Iguanura*, and spiny leaves in *Phoenicophorium*. Of these genera, only the geographical range of *Cyphosperma* overlaps with that of *Heterospathe* and *Alsmithia*. Notably, in the most parsimonious tree selected for exploring character optimization, *Cyphosperma* is sister to the *Heterospathe*-*Alsmithia* clade, and thus the character states listed before are optimized at the common node of

Heterospathe, *Alsmithia*, and *Cyphosperma* rather than at the basal node of the *Heterospathe*-*Alsmithia* clade. However, this relationship is not present in the strict consensus trees of any of our analyses.

The optimization of character nine (endosperm type) yields numerous independent transformations between the two states, homogeneous and ruminant, underscoring the labile nature of this condition. Ruminant endosperm was a defining character of *Heterospathe* in its delimitation according to Uhl and Dransfield (1987). Since 1987, however, a species of *Heterospathe* has been recorded with homogeneous endosperm (*H. uniformis* Dowe; Dowe and Cabalion, 1996). Furthermore, *Alsmithia* has homogeneous endosperm. Palm genera that include both ruminant and homogeneous species are well known (e.g., *Phoenix* L., *Reinhardtia*, *Hydriastele*, and *Calyptrocalyx*), but more recently infraspecific variation in this character has also been documented in *Ptychococcus* (Zona, 2003).

In addition to its homogeneous endosperm, *Alsmithia* deviates from *Heterospathe* in its highly sculptured endocarp (character eight) that is more reminiscent of those found in other western Pacific genera, such as *Burretio kentia*, *Veillonina*, and *Cyphosperma*, rather than the smooth endocarps typical of *Heterospathe*. However, Essig et al. (1999) noted some similarity between the pericarp composition in *Alsmithia* and *Heterospathe* (in particular *H. woodfordiana* Becc.) in that they both possess an expanded series of vascular bundles. A similar example is also seen in the coryphoid palm *Licuala* Wurm. in which some New Guinean species have developed a thick and highly ornamented endocarp, probably as an adaptation to specific dispersers (Barfod, 2000; Barfod et al., 2001). Optimization of this character suggests that ornate endocarps have arisen independently at least six times within the Areceae. Whether the developmental origin of endocarp sculpturing is the same in all Areceae genera is unknown and merits further investigation. From the limited material available to us, we suspect that the so-called endocarp of *Alsmithia* may in fact consist of hardened, elaborated mesocarp fibers. Regardless of this autapomorphy, the molecular evidence in combination with the numerous morphological similarities strongly supports the inclusion of *Alsmithia* in *Heterospathe* (Norup, 2005).

Despite possessing the suite of crownshaft character states that is common to so many Areceae genera, *Rhopaloblade* is in fact well defined morphologically. In contrast to *Heterospathe*, two almost uncontradicted synapomorphies have been identified for the genus: inflorescence branches convoluted in bud and a pinnate eophyll. Outside the Areceae, convoluted inflorescence branches are known to occur in species of *Geonoma* and a small number of other New World palms. Pinnate eophylls are unique to *Rhopaloblade* and *Acanthophoenix* in the Areceae, but are also found outside the clade, e.g., in *Podococcus*. However, this combination is not shared by any other Areceae genus. Several additional characters not included in our study can be added (Banka and Baker, 2004): an indumentum on the leaf rachis and petiole consisting of multiple laminar, membranous brownish-black scales; strongly divaricate primary inflorescence branches; and a club-shaped embryo.

As well as showing no phylogenetic relationship to *Heterospathe* and *Rhopaloblade*, the problematic taxon *Ptychosperma micranthum* does not conform to the character combinations previously listed or those of the genus *Ptychosperma*, indicating that the placement of this species

within these three genera by previous authors was in error. The similarity to *Heterospathe* is limited to the persistence of the prophyll and a ruminant endosperm. *Ptychosperma micranthum* possesses none of the characters of *Rhopaloblade* just listed and differs from *Ptychosperma* in, among other characters, its persistent rather than caducous prophyll and acute rather than praemorse leaflets. Crownshafted palms that possess persistent prophylls, such as *P. micranthum*, are rare and found only in *Roscheria* and some species of *Dypsis* and *Drymophloeus* Zipp. Our analysis demonstrates that *P. micranthum* is not related to any of these genera. Morphologically, it is readily distinguished from *Roscheria* in being unarmed rather than spiny, from *Drymophloeus* in bearing acute rather than praemorse leaflets, and from *Dypsis* in fruits bearing apical rather than basal stigmatic remains. The phylogenetic and morphological evidence presented here provides an unequivocal justification for the new genus *Dransfieldia* that has recently been erected to accommodate *P. micranthum* (Baker et al., 2006).

Biogeographical implications of the analysis for *Heterospathe* and *Rhopaloblade*—The high resolution at the species level within *Heterospathe* and *Rhopaloblade* makes it possible to explore the biogeographic implications of their modern distribution. We tentatively suggest that *Rhopaloblade* is of Indian Ocean origin, given its placement in the Indian Ocean clade, its sister group relationship to *Dictyosperma* from the Mascarenes, and the position of *R. augusta* from Nicobar Island, with subsequent dispersal eastwards into Malesia and Papua. The position of *R. singaporensis* could indicate a later dispersal back westward, possibly from New Guinea. However, in the absence of the remaining two species of *Rhopaloblade* (*R. gideonii* R. Banka from New Ireland and *R. elegans* H. E. Moore from the Solomon Islands) this scenario remains speculative. Optimal conditions for dispersal across Wallace's Line from New Guinea back to Peninsular Malaysia occurred in the Tertiary at the boundary between Miocene and Pliocene when a dispersal route was opened by the coherent landmasses that formed the Banda and Sunda arcs (Hall, 1998). Other possibilities to account for the present disjunct distribution include extinction, long-distance dispersal, or inadequate knowledge of the genus in the intervening areas, e.g., in Sulawesi.

Due to the small sample size in this study, conclusions on evolutionary and biogeographic patterns of *Heterospathe* are preliminary. As shown in Fig. 3, *Heterospathe* falls within the western Pacific clade. The two clades within *Heterospathe* have distinct geographies. Clade 1 consists of the Philippine species and the two New Guinean species, *H. humilis* and *H. delicatula*, as well as the Fijian *H. phillipsii* and *Alsmithia longipes*. These patterns cannot be explained by vicariance events and almost certainly represent dispersal events. In contrast, clade 2 is endemic to New Guinea. The following geological events described by Burrett et al. (1991) and Hall (1998) may have influenced this radiation and present-day distribution of *Heterospathe*: (1) Fiji, the Solomons, and Vanuatu were close enough to allow for amalgamation of biotas until the late Miocene (ca. 10 Ma, as parts of the Melanesian arc); (2) New Guinea was in contact with Fiji, the Solomons, and Vanuatu through the western end of the Melanesian arc (ca. 40 Ma) and later via the South Caroline arc (until ca. 10 Ma); and (3) there was a discontinuous land-bridge between New Guinea and the Philippines via the Philippine-

Halmahera arc from around 25 Ma. Thus, there have been numerous opportunities for dispersal both eastward and westward during the Miocene with subsequent radiation, especially in the Philippines, which continued to change until about 10 Mya (Burrett et al., 1991; Hall, 1998). Clear-cut biogeographic patterns cannot, therefore, be expected.

Concluding remarks—By combining the two low-copy nuclear genes *PRK* and *RPB2*, we have confirmed that the genera *Heterospathe* and *Rhopaloblaste* are distinct and unrelated groups that are both members of the Areceae. *Rhopaloblaste* is strongly supported as monophyletic, whereas the monotypic genus *Alsmithia* is clearly nested within *Heterospathe*, indicating that the taxonomic boundaries of *Heterospathe* should be altered slightly (Norup, 2005). Relative to the results of the molecular analyses, the morphological characters used in the delimitation of *Heterospathe* and *Rhopaloblaste* are homoplasious, especially in *Heterospathe*, for which no unique characters have been identified (cf. Hughes et al., 2004). Only by applying suites of characters is it possible to firmly establish generic limits, accounting for historical problems with generic delimitation.

Despite recovering many novel relationships, our phylogenies are frustratingly ambiguous in places. A low level of resolution in a large group of plants in the Indo-Pacific ocean, coupled with a lack of morphological synapomorphies, a non-fit between current infrafamilial classifications, and complex biogeographic history (as for *Heterospathe* and *Rhopaloblaste*) has also recently been documented within Araliaceae (Plunkett et al., 2004; Wen et al., 2004). The authors concluded that the large polytomies found in a clade consisting of taxa from Malesia, the Indian and Pacific Oceans, and Australia might be a result of a rapid simultaneous radiation early in the evolutionary history of the clade, although taxon and character sampling may also be influential. Whether or not this represents a plausible hypothesis for the poor resolution within the Indo-Pacific pseudomonomerous clade remains to be evaluated.

LITERATURE CITED

- ASMUSSEN, C. B., W. J. BAKER, AND J. DRANSFIELD. 2000. Phylogeny of the palm family (Arecaceae) based on *rps16* intron and *trnL-trnF* plastid DNA sequences. In K. L. Wilson and D. A. Morrison [eds.], *Monocots: systematics and evolution*, 525–537. CSIRO, Melbourne, Australia.
- ASMUSSEN, C. B., AND M. W. CHASE. 2001. Coding and noncoding plastid DNA in palm systematics. *American Journal of Botany* 88: 1103–1117.
- ASMUSSEN, C. B., J. DRANSFIELD, V. DEICKMANN, A. S. BARFOD, J.-C. PINTAUD, AND W. J. BAKER. In press. A new subfamily classification of the palm family (Arecaceae): evidence from plastid DNA phylogeny. *Botanical Journal of the Linnean Society*.
- BAKER, W. J. 2002. The palms of New Guinea project. *Flora Malesiana Bulletin* 13: 35–37.
- BAKER, W. J., C. B. ASMUSSEN, S. BARROW, J. DRANSFIELD, AND T. A. HEDDERSON. 1999. A phylogenetic study of the palm family (Palmae) based on chloroplast DNA sequences from the *trnL-trnF* region. *Plant Systematics and Evolution* 219: 111–126.
- BAKER, W. J., J. DRANSFIELD, AND T. A. HEDDERSON. 2000a. Phylogeny, character evolution, and a new classification of the calamoid palms. *Systematic Botany* 25: 297–322.
- BAKER, W. J., T. A. HEDDERSON, AND J. DRANSFIELD. 2000b. Molecular phylogenetics of subfamily Calamoideae (Palmae) based on nrDNA ITS and cpDNA *rps16* intron sequence data. *Molecular Phylogenetics and Evolution* 14: 195–217.
- BAKER, W. J., T. A. HEDDERSON, AND J. DRANSFIELD. 2000c. Molecular phylogenetics of *Calamus* (Palmae) and related rattan genera based on 5S nrDNA spacer sequence data. *Molecular Phylogenetics and Evolution* 14: 218–231.
- BAKER, W. J., AND A. H. B. LOO. 2004. A synopsis of the genus *Hydriastele*. *Kew Bulletin* 59: 61–68.
- BAKER, W. J., S. ZONA, C. D. HEATUBUN, C. E. LEWIS, R. A. MATURBONGS, AND M. V. NORUP. 2006. *Dransfieldia* (Arecaceae)—a new palm genus from western New Guinea. *Systematic Botany* 31: 60–68.
- BANKA, R., AND W. BAKER. 2004. A monograph of the genus *Rhopaloblaste* (Arecaceae). *Kew Bulletin* 59: 47–60.
- BARFOD, A. S. 2000. A new species of *Licuala* from New Guinea. *Palms* 44: 198–201.
- BARFOD, A. S., R. BANKA, AND J. L. DOWE. 2001. Field guide to palms in Papua New Guinea. *AAU Reports* 40: 1–77.
- BECCARI, O. 1877. Palmae Papuane. *Malesia* 1: 52–53.
- BECCARI, O. 1885. Reliquiae Schefferianae. Illustrazione di alcune palme viventi nel Giardino Botanico di Buitenzorg. *Annales du Jardin Botanique de Buitenzorg* 2:90, 126.
- BECCARI, O. 1909. Notes on Philippine palms. II. *Philippine Journal of Science, Botany* 4: 610–637.
- BURRETT, C., N. DUHIG, R. BERRY, AND R. VARNE. 1991. Asian and southwestern Pacific continental terranes derived from Gondwana, and their biogeographic significance. *Australian Systematic Botany* 4: 13–24.
- CHASE, M. W., AND H. G. HILLS. 1991. Silica gel: an ideal material for field preservation of leaf samples for DNA studies. *Taxon* 40: 215–220.
- DENTON, A. L., B. L. MCCONAUGHY, AND B. J. HALL. 1998. Usefulness of RNA polymerase II coding sequences for estimation of green plant phylogeny. *Molecular Biology and Evolution* 15: 1082–1085.
- DOLPHIN, K., R. BELSHAW, C. D. L. ORME, AND D. L. J. QUICKE. 2000. Noise and incongruence: interpreting results of the incongruence length difference test. *Molecular Phylogenetics and Evolution* 17: 401–406.
- DOWE, J. L., AND P. CABALION. 1996. A taxonomic account of Arecaceae in Vanuatu, with descriptions of three new species. *Australian Systematic Botany* 9: 1–60.
- DOWE, J. L., AND N. W. UHL. 1989. *Carpoxylon macrospermum*. *Principes* 33: 68–73.
- DOYLE, J. J., AND J. L. DOYLE. 1987. A rapid DNA isolation procedure from small quantities of fresh leaf tissue. *Phytochemistry Bulletin* 19: 11–15.
- DRANSFIELD, J. 1991. *Lemurophoenix* (Palmae: Arecoideae), a new genus from Madagascar. *Kew Bulletin* 46: 61–68.
- DRANSFIELD, J. 1999. Species and species concepts in Old World palms. In A. Henderson and F. Borchsenius [eds.], *Evolution, variation and classification of palms*. Memoirs of the New York Botanical Garden, vol. 83, 5–20. New York Botanical Garden Press, Bronx, New York, USA.
- DRANSFIELD, J., AND H. J. BEENTJE. 1995. The palms of Madagascar. Royal Botanic Gardens Kew and the International Palm Society, Norwich Print Services, Kew, UK.
- DRANSFIELD, J., AND N. W. UHL. 1998. Palmae. In K. Kubitzki [ed.], *The families and genera of vascular plants*, vol. 4, Flowering plants: monocotyledons, 306–389. Springer-Verlag, Berlin, Germany.
- DRANSFIELD, J., N. W. UHL, C. B. ASMUSSEN, W. J. BAKER, M. M. HARLEY, AND C. E. LEWIS. 2005. A new phylogenetic classification of the palm family, Arecaceae. *Kew Bulletin* 60: 559–569.
- ESSIG, F. B., T. J. MANKA, AND L. BUSSARD. 1999. A systematic histological study of palm fruits. III. Subtribe Iguanurinae (Arecaceae). *Brittonia* 51: 307–325.
- FARRIS, J. S., M. KÄLLERSJÖ, A. G. KLUGE, AND C. BULT. 1995. Testing significance of incongruence. *Cladistics* 10: 315–319.
- FELSENSTEIN, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783–791.
- FERNANDO, E. S. 1990a. A preliminary analysis of the palm flora of the Philippine Islands. *Principes* 34: 28–45.
- FERNANDO, E. S. 1990b. The genus *Heterospathe* (Palmae, Arecoideae) in the Philippines. *Kew Bulletin* 45: 219–234.

- FITCH, W. M. 1971. Towards defining the course of evolution: minimum change for a specific tree typology. *Systematic Zoology* 20: 406–416.
- GOVAERTS, R., AND J. DRANSFIELD. 2005. World checklist of palms. Royal Botanic Gardens, Kew, UK.
- GUNN, B. F. 2004. The phylogeny of the Cocoeae (Arecaceae) with emphasis on *Cocos nucifera*. *Annals of Missouri Botanical Gardens* 91: 505–522.
- HAHN, W. J. 2002. A phylogenetic analysis of the arecoid line of palms based on plastid DNA sequence data. *Molecular Phylogenetics and Evolution* 23: 189–204.
- HALL, R. 1998. The plate tectonics of Cenozoic SE Asia and the distribution of land and sea. In R. Hall and J. D. Holloway [eds.], *Biogeography and geological evolution of SE Asia*, 99–131. Backhuys Publishers, Leiden, Netherlands.
- HOOKE, J. D. 1883. Ordo 187. Palmae. In G. Bentham and J. D. Hooker [eds.], *Genera plantarum* 3, 870–948. Lovell Reeve, London, UK.
- HUELSENBECK, J. P., J. J. BULL, AND C. W. CUNNINGHAM. 1996. Combining data in phylogenetic analysis. *Trends in Ecology and Evolution* 4: 152–158.
- HUGHES, C. E., G. P. LEWIS, A. D. YOMONA, AND C. REYNEL. 2004. *Maraniona*. A new dalbergioid legume genus (Leguminosae, Papilionoideae) from Peru. *Systematic Botany* 29: 366–374.
- JACKSON, B. D. 1895. Index kewensis, vol. 2, part 4. Clarendon Press, Oxford, UK.
- KAWAGISHI, M., M. YAMAGISHI, AND A. ISHIHAMA. 1993. Cloning and sequence determination of the *Schizosaccharomyces pombe* *rpb2* gene encoding the subunit 2 of RNA polymerase II. *Nucleic Acids Research* 21: 469–473.
- KELCHNER, S. A. 2000. The evolution of non-coding chloroplast DNA and its application in plant systematics. *Annals of Missouri Botanical Garden* 87: 482–498.
- LEWIS, C. E., AND J. J. DOYLE. 2001. Phylogenetic utility of the nuclear gene malate synthase in the palm family (Arecaceae). *Molecular Phylogenetics and Evolution* 19: 409–420.
- LEWIS, C. E., AND J. J. DOYLE. 2002. Phylogenetics of tribe Areceae (Arecaceae) using low-copy nuclear genes. *Plant Systematics and Evolution* 236: 1–17.
- LEWIS, C. E., AND N. MARTÍNEZ. 2000. Identity of the *Hyophorbe* palms at the Botanical Garden of Cienfuegos, Cuba. *Palms* 44: 93–97.
- LOO, A. H. B., J. DRANSFIELD, M. W. CHASE, AND W. J. BAKER. 2006. Low copy nuclear DNA, phylogeny and the evolution of dichogamy in the betel nut palms and their relatives (Arecinae; Arecaceae). *Molecular Phylogenetics and Evolution*: in press.
- MADDISON, D. R., AND W. P. MADDISON. 2003. MacClade, version 4.06. Analysis of phylogeny and character evolution. Sinauer, Sunderland, Massachusetts, USA.
- MARTELLI, U. 1934. Generi, specie, e varietà nuove di palme gerontogee della tribù Arecaceae lasciate inedite dal Dr. Od. Beccari ed ordinate acura di U. Martelli. *Atti Società Toscana di Scienze Naturali residente in Pisa. Memorie* 44: 114–176.
- MARTELLI, U. 1935. La sinonimia delle palme gerontogee della tribù delle Areceae. *Nuovo Giornale Botanico Italiano, Nuova Serie* 42: 81.
- MOORE, H. E. JR. 1970. The genus *Rhopaloblade* (Palmae). *Principes* 14: 75–92.
- MOORE, H. E. JR., R. H. PHILLIPS, AND S. VODONAIVALU. 1982. Additions to the palms of Fiji. *Principes* 26: 122–125.
- MORLEY, R. J. 2000. Origin and evolution of tropical rain forest. Wiley, Chichester, UK.
- MORLEY, R. J. 2001. Tertiary history of the Malesian flora: a palynological perspective. In L. G. Saw, L. S. L. Chua, and K. C. Khoo [eds.], *Taxonomy: the cornerstone of biodiversity*, 197–210. Proceeding of the Fourth International Flora Malesiana Symposium 1998, The Research Institute Malaysia, Kepong, Kuala Lumpur.
- MORT, M. E., AND D. J. CRAWFORD. 2004. The continuing search: low-copy nuclear sequences for lower-level plant molecular phylogenetic studies. *Taxon* 53: 257–261.
- MOYLAN, E. C., J. R. BENNET, M. A. CARINE, R. G. OLMSTEAD, AND R. W. SCOTLAND. 2004. Phylogenetic relationships among *Strobilanthes* s. l. (Acanthaceae): evidence from ITS nrDNA, *trnL-F*, cpDNA, and morphology. *American Journal of Botany* 91: 724–735.
- NORUP, M. V. 2005. *Alsmithia* subsumed in *Heterospatha* (Arecaceae, Arecoideae). *Novon* 15: 455–457.
- NORUP, M. V., W. BAKER, C. LEWIS, A. LOO, AND J. DRANSFIELD. 2003. Phylogenetic and biogeographical studies of the Indo-Pacific clade of tribe Areceae (Palmae), based on two low-copy nuclear genes. Monocots III symposium, 2003, Ontario, California, USA. Rancho Santa Ana Botanical Gardens, California, USA (Abstract).
- OXELMAN, B., AND B. BREMER. 2000. Discovery of paralogous nuclear gene sequences coding for the second-largest subunit of RNA polymerase II (*RPB2*) and their phylogenetic utility in Gentianales of the asterids. *Molecular Biology and Evolution* 17: 1130–1145.
- OXELMAN, B., N. YOSHIKAWA, B. L. MCCONAUGHY, J. LUO, A. L. DENTON, AND B. D. HALL. 2004. *Rpb2* gene phylogeny in flowering plants, with particular emphasis on asterids. *Molecular Phylogenetics and Evolution* 32: 462–479.
- PFEIL, B. E., C. L. BRUBAKER, L. A. CRAVEN, AND M. D. CRISP. 2004. Paralogy and orthology in the Malvaceae *rpb2* gene family: investigation of gene duplication in *Hibiscus*. *Molecular Biology and Evolution* 21: 1428–1433.
- PINTAUD, J.-C., AND D. R. HODEL. 1998. The palms of New Caledonia. Allen Press, Lawrence, Kansas, USA.
- PLUNKETT, G. M., J. WEN, AND P. P. LOWRY II. 2004. Intrafamilial classifications and characters in Araliaceae: insights from the phylogenetic analysis of nuclear (ITS) and plastid (*trnL-trnF*) sequence data. *Plant Systematics and Evolution* 245: 1–39.
- POPP, M., AND B. OXELMAN. 2004. Evolution of a RNA polymerase gene family in *Silene* (Caryophyllaceae)—incomplete concerted evolution and topological congruence among paralogues. *Systematic Biology* 53: 914–932.
- REEVES, G., M. W. CHASE, P. GOLDBLATT, P. RUDALL, M. W. FAY, A. V. COX, B. LEJEUNE, AND T. SOUZA-CHIES. 2001. Molecular systematics of Iridaceae: evidence from four plastid DNA regions. *American Journal of Botany* 88: 2074–2087.
- RIDDER-NUMAN, J. W. A. 1996. Historical biogeography of the Southeast Asian genus *Spatholobus* (Legum.-Papilionoideae) and its allies. *Blumea Supplement* 10: 1–144.
- RONCAL, J., J. FRANCISCO-ORTEGA, C. B. ASMUSSEN, AND C. E. LEWIS. 2005. Molecular phylogenetics of tribe Geonomeae (Arecaceae) using nuclear DNA sequences of phosphoribulokinase and RNA polymerase II. *Systematic Botany* 30: 275–283.
- SALAMIN, N., M. W. CHASE, T. R. HODKINSON, AND V. SAVOLAINEN. 2003. Assessing internal support with large phylogenetic DNA matrices. *Molecular Phylogenetics and Evolution* 27: 528–539.
- SCHIEFFER, R. H. C. C. 1876. Sur quelques palmiers du groupe des Arécinées. Deuxième partie. *Annales du Jardin Botanique de Buitenzorg* 1: 103–164.
- SHEAHAN, M. C., AND M. W. CHASE. 2000. Phylogenetic relationships within Zygophyllaceae based on DNA sequences of three plastid regions, with special emphasis on Zygophylloideae. *Systematic Botany* 25: 371–384.
- SMALL, R. L., R. C. CRONN, AND J. F. WENDEL. 2004. Use of nuclear genes for phylogeny reconstruction in plants. *Australian Systematic Botany* 17: 145–170.
- STAUFFER, F. W., W. J. BAKER, J. DRANSFIELD, AND P. K. ENDRESS. 2004. Comparative floral structure and systematics of *Pelagodoxa* and *Sommieria* (Arecaceae). *Botanical Journal of the Linnean Society* 146: 27–39.
- SWOFFORD, D. L. 2002. PAUP*: phylogenetic analysis using parsimony (*and other methods), version 4. Sinauer, Sunderland, Massachusetts, USA.
- TAM, S.-M., P. C. BOYCE, T. M. UPSON, D. BARABÉ, A. BRUNEAU, F. FOREST, AND J. S. PARKER. 2004. Intergeneric and intrafamilial phylogeny of subfamily Monsteroideae (Araceae) revealed by chloroplast *trnL-F* sequences. *American Journal of Botany* 91: 490–498.
- THOMAS, M. M., N. C. GARWOOD, W. J. BAKER, S. A. HENDERSON, S. J. RUSSELL, D. R. HODEL, AND R. M. BATEMAN. 2006. Molecular phylogeny of the palm genus *Chamaedorea*, based on the low-copy

- nuclear genes *PRK* and *RPB2*. *Molecular Phylogenetics and Evolution*.
- TURNER, H. 1995. Cladistic and biogeographic analyses of *Arytera* Blume and *Mischarytera* gen. nov. (Sapindaceae). *Blumea Supplement* 9: 1–230.
- UHL, N. W., AND J. DRANSFIELD. 1987. Genera palmarum: a classification of palms based on the work of H. E. Moore Jr. International Palm Society and L. H. Bailey Hortorium. Allen Press, Lawrence, Kansas, USA.
- WEN, J., C. LEE, G. PLUNKETT, AND P. P. LOWRY, II. 2004. Evolution of Araliaceae in Asia: inference from the ITS, *ndhF*, *trnL-F*, and molecular data. Sixth International Flora Malesiana Symposium, 2004, Los Baños, Philippines. University of the Philippines, Los Baños, Philippines (Abstract).
- WIENS, J. J. 1998. Combining data sets with different phylogenetic histories. *Systematic Biology* 47: 568–581.
- WILSON, M. A., B. GAUT, AND M. T. CLEGG. 1990. Chloroplast DNA evolve slowly in the palm family (Arecaceae). *Molecular Biology and Evolution* 7: 303–314.
- YODER, A. D., J. A. IRWIN, AND B. A. PAYSEUR. 2001. Failure of the ILD to determine data combinability for slow loris phylogeny. *Systematic Biology* 50: 408–424.
- ZONA, S. 2003. Endosperm condition and the paradox of *Ptychococcus paradoxus*. *Telopea* 10: 1–7.

APPENDIX. Taxa used in this study, voucher information, GenBank accession numbers for the two regions studied. A dash indicates the region was not sampled. Herbarium acronyms are given in parentheses after accessions: AAU = University of Aarhus, BH = Cornell University, New York, BISH = Bishop Museum, Hawaii, FTG = Fairchild Tropical Botanic Garden, Miami, GUAM = University of Guam, JCT = James Cook University of Northern Queensland, K = Royal Botanic Gardens, Kew, LAE = Papua New Guinea Forest Research Institute, LBC = University of the Philippines at Los Baños, MAK = Tokyo Metropolitan University, NOU = Institut de Recherche pour le Développement, New Caledonia, NY = New York Botanical Garden, P = Muséum National d'Histoire Naturelle, Paris, SUVA = University of the South Pacific, Fiji, TL = Université Paul Sabatier, Toulouse, VEN = Fundación Instituto Botánico de Venezuela Dr. Tobías Lasser. Asterisks indicate sequences that were cloned. In the case of several clones per species, underlining indicates the clone used in the final matrix. Sequences published previously by Lewis and Doyle (2002) have GenBank numbers starting with AF.

Subfamily: Tribe; Subtribe

Taxon—Voucher specimen (Herbarium); GenBank accession: *PRK*, *RPB2*.

Arecoideae; Areceae; Archontophoenicinae

Actinorhytis calapparia (Blume) H. Wendl. & Drude ex Scheff.—*Heatubun* 192 (K); AJ831223, AJ830025. *Actinokentia huerlimannii* (Brongn.) Dammer—*Pintaud* 465 (NOU); AJ831222, AJ830023. *Archontophoenix purpurea* Hodel & Dowe—*Pintaud* 492 (TL); AJ831227, AJ830028. *Chambeyronia macrocarpa* (Brongn.) Vieill. ex Becc.—*Pintaud* 512 (P); AJ831260, AJ830056. *Kentiopsis oliviformis* (Brongn. & Gris.) Brongn.—*Pintaud* 358 (K); AF453353, AY543100.

Arecoideae; Areceae; Arecinae

Areca catechu L.—*Lewis* 98-093 (BH) for *PRK*, *TCMK* 21 (K) for *RPB2*; AF453333, AY543109. *Nenga pumila* (Blume) H. Wendl.) var. *pachystachya* (Blume) Fernando—*Baker* 994 (FTG); AY348914, AY543154. *Pinanga coronata* (Blume ex Mart.) Blume—*Baker* 1145 (K); AY348944, AY543156.

Arecoideae; Areceae; Basselininae

Alloschmidia glabrata (Becc.) H.E. Moore—*Pintaud* 468 (K); AJ831225, AJ830026. *Basselinia velutina* Becc.—*Pintaud* 553 (P); AJ831233, AJ830031. *BurretioKentia vieillardii* (Brongn. & Gris.) Pic.Serm.—*Pintaud* 197 (NY); AJ831243, AJ830039. *Campecarpus fulcitus* (Brongn.) H. Wendl. ex Becc.—*Pintaud* 524 (P); AJ831258, AJ830054. *Cyphophoenix nucele* H.E. Moore—*Pintaud* 372 (K); AJ831266, AJ830061. *Cyphosperma balansae* (Brongn.) H. Wendl. ex Salomon—*Baker* 89-030 (BISH); AF453340, AY543098. *Physokentia petiolata* (Burret) D. Fuller—*Pintaud* 452 (TL); AJ831322, AJ830138. *Veillonina alba* H.E. Moore—*Pintaud* 277 (K); AJ831336*, AJ831337*, AJ831338*, AJ831339*, AJ831340*, AJ831341*, AJ830149.

Arecoideae; Areceae; Carpoxylinae

Carpoxydon macrosperrum H. Wendl. & Drude—*Zona* 722 (FTG); AF453337, AJ830055. *Neoveitchia storckii* (H. Wendl.) Becc.—*Roncal* 73 (FTG); AJ831319, AJ830130. *Satakentia luikiuensis* (Hatus.) H.E. Moore—*Lewis* 99-051 (BISH); AF453376, AJ830146.

Arecoideae; Areceae; Clinospermatinae

Brongniartikentia lanuginosa H.E. Moore—*Pintaud* 368 (P); AJ831236*, AJ831237*, AJ831238*, AJ831239*, AJ831240*, AJ830033. *Clinosperma bracteale* (Brongn.) Becc.—*Pintaud* 349 (K); AJ831261, AJ830057. *Cyphokentia macrostachya* Brongn.—*Pintaud* 558 (P); AJ831264, AJ830060*. *Lavoixia macrocarpa* H.E. Moore—*Pintaud* 364 (P); AJ831302, AJ830110. *Moratia cerifera* H. E. Moore—*Pintaud* 347 (K); AJ831318, AJ830129.

Arecoideae; Areceae; Dypsidinae

Dypsis lutescens (H. Wendl.) Beentje & J. Dransf.—*Lewis* 00-004 (BH); AF453346, AJ830078. *Lemurophoenix halleuxii* J. Dransf.—*Lewis* 98-073 (BH); AF453354, AM260636. *Marojejya darianii* J. Dransf. & N.W. Uhl—*Lewis* 99-037 (BISH); AF453359, AJ830121*. *Masoala madagascariensis* Jum.—1992-3552 (K); AF453360, AJ830128.

Arecoideae; Areceae; Linospadicinae

Calyptracalyx hollrungii (Becc.) Dowe & M.D. Ferrero—*Baker* 1176 (K); AJ831251 AJ830043. *Howea belmoreana* (C. Moore & F. Muell.) Becc.—*Baker* 1154 (K); AJ831294, AJ830098. *Laccospadix australasicus* H. Wendl. & Drude—*Baker* 1172 (K); AJ831300, AJ830108. *Linospadix albertisiana* (Becc.) Burret—*Dowe* 720 (JCT); AJ831305, AJ830119.

Arecoideae; Areceae; Oncospermatinae

Acanthophoenix rubra (Bory) H. Wendl.—*Lewis* 98-067 (BH); AF453329, AJ830020. *Deckenia nobilis* H. Wendl. ex Seem.—*Lewis* 98-031 (BH); AF453342, AJ830063. *Oncosperma tigillarum* (Jack) Ridl.—*Lewis* 98-051 (BH); AF453364, AJ830134. *Tectiphiala ferox* H.E. Moore—*Lewis* 98-070 (BH); AF453380, AJ830148.

Arecoideae; Areceae; Ptychospermatinae

Adonidia merrillii (Becc.) Becc.—*Zona* 874 (FTG); AJ831224, AJ830193. *Balaka longirostris* Becc.—*Pintaud* 463 (SUVA); AJ831229, AJ830029. *Brassiophoenix drymophloeoides* Burret—*Coons* 1398 (FTG); AJ831235, AJ830195. *Carpentaria acuminata* (H. Wendl. & Drude) Becc.—*Zona* 827 (FTG); AJ831259, AJ830196. *Drymophloeus litigiousus* (Becc.) H.E. Moore—*Barrow* 125 (K); AJ831267, AJ830197. *Normanbya normanbyi* (F.Muell.) L.H. Bailey—*Lewis* 98-091 (BH); AF453363, AJ830132. *Ponapea palauensis* Kaneh—*Lewis* 99-055 (BISH); AJ831328, AJ830203. *Ptychococcus paradoxus* (Scheff.) Becc.—*Baker* 572 (K); AJ831324, AJ830200. *Ptychosperma salomonense* Burret—*Houghton* 1300 (FTG); AF453371, AY543105. *Solfia samoensis* Rech.—*Tipama'a* 001 (FTG); AJ831334, AJ830204. *Veitchia spiralis* H. Wendl.—*Zona* 724 (FTG); AJ831342, AJ830205. *Wodyetia bifurcata* A.K. Irvine—*Zona* 906 (FTG); AJ831343, AJ830206.

Arecoideae; Areceae; Rhopalostylidinae

Hedyscepe canterburyana (C. Moore & F. Muell.) H. Wendl. & Drude—*Pintaud* 407 (TL); AJ831276, AJ830081. *Rhopalostylis baueri* (Hook. f. ex Lem.) H. Wendl. & Drude—*Pintaud* 384 (NY); AJ831333, AJ830145.

Arecoideae; Areceae; Verschaffeltinae

Nephrosperma vanhoutteanum (H. Wendl. ex Van Houtt.) Balf. f.—*Lewis* 98-006 (BH); AF453362, AJ830131. *Phoenicophorium*

- borsigianum* (K. Koch) Stuntz—Lewis 98-024 (K); AF453368, AJ830136. *Roscheria melanochaetes* (H. Wendl.) H. Wendl. ex Balf. f.—Lewis 98-036 (BH); AF453374, AJ830140*. *Verschaffeltia splendida* H. Wendl.—Lewis 98-039 (BH); AF453381, AJ830150.
- Arecoideae; Areceae; unplaced
- Alsmithia longipes* H.E. Moore—Baker 1180 (FTG); AJ831226, AJ830027. *A. longipes* H.E. Moore—Zona 1039 (FTG); AM260638, AM260637. *Bentinckia condapanna* Berry ex Roxb.—1993-2989 (K); AF453336, AJ830032. *Clinostigma savoryanum* (Rehder & E.H. Wilson) H.E. Moore & Fosberg—Pintaud 442 (MAK); AJ831263, AJ830059. *Cyrtostachys renda* Blume—1982-5882 (K); AF453341, AJ830062. *Dictyosperma album* (Bory) Scheff.—Lewis 98-031 (BH); AF453343, AJ830064. *Heterospatha cagayanensis* Becc.—Kyburz s. n. (no voucher); AJ831277, AJ830082. *H. delicatula* H.E. Moore—Baker 1190 (K); AJ831278, AJ830083. *H. elata* Scheff.—Chapin 55 (K); AJ831279, AJ830084. *H. elata* Scheff.—Houghton & Hubbuck 1024 (FTG); AJ831292, AJ830096. *H. elata* Scheff.—Lewis 99-034 (GUAM); AF453350, AJ830085. *H. humilis* Becc.—Banka 2011 (K); AJ831280, AJ830086. *H. macgregorii* (Becc.) H. E. Moore—Baker 651 (K); AJ831281, AJ830087. *H. philippinensis* (Becc.) Becc.—Fernando 1623 (LBC); AJ831282, AJ833634. *H. phillipsii* D. Fuller & Dowe—Pintaud 454 (SUA); AJ831283, AJ830088*. *H. scitula* Fernando—Fernando 1625 (LBC); AJ831284, AJ830089. *H. sibuyanensis* Becc.—Zona 1050 (FTG); AJ831285, AJ830090. *H. sp.*—Baker 1115 (K); AJ831286, AJ830091. *H. sp.*—Baker 1130 (K); AJ831290, —. *H. sp.*—Baker 1142 (K); AJ831288, AJ830093. *H. sp.*—Banka 2008 (K); AJ831289, AJ830094. *H. sp. nov.*—Fernando 1624 (LBC); AJ831291, AJ830095. *Hydriastele beguinii* (Burret) W. J. Baker & Loo—Zona 799 (FTG); AY348951, AY543163. *H. brassii* (Burret) W.J. Baker & Loo—Baker 823 (K); AY348916, AY543116. *H. costata* F.M. Bailey—Baker 836 (K); AY348925, AY543127. *H. microspadix* (Warb. ex K. Schum. & Lauterb.) Burret—Baker 573 (K); AY348932, AY543136. *Iguanura wallichiana* (Mart.) Becc.—Lewis 99-049 (BISH); AF453352, AY543099. *Lepidorrhachis mooreana* (F. Muell.) O.F. Cook—Baker 1168 (K); AJ831304, AJ830118. *Loxococcus rupicola* (Thwaites) H. Wendl. & Drude—1990-2497 (K); AY348942, AY543151. *Ptychosperma micranthum* Becc.—Baker 1066 (K); AJ831326, AJ830139. *Rhopaloblaste augusta* (Kurz) H.E. Moore—Lewis 99-004 (FTG); AF453373, AY543107. *R. ceramica* (Miq.) Burret—Banka 2050 (LAE); AJ831329, AJ830141. *R. ledermanniana* Becc.—Heatubun 191 (K); AJ831331, AJ830144. *R. ledermanniana* Becc.—Maturbongs 650 (K); AJ831332, AJ830143. *R. singaporensis* (Becc.) Hook. f.—Baker 1174 (K); AJ831330, AJ830142.
- Arecoideae; Euterpeae
- Neonicholsonia watsonii* Dammer—Lewis 99-052 (BISH); AJ831356; AJ830172.
- Arecoideae; Geonomateae
- Asterogyne martiana* (H. Wendl.) H. Wendl. ex Drude—Baker 89021 (BISH); AF453334, AJ830154. *Calypogyne costatifrons* (L.H. Bailey) Nevers—Knudsen & Asmussen 603 (AAU); AJ831347, AJ830208. *Geonoma deversa* (Poir.) Kunth—Roncal 19 (FTG); AJ831354, AJ830210. *Pholidostachys pulchra* H. Wendl. ex Burret—Roncal 26 (FTG); AJ831360, AJ830211.
- Arecoideae; Iriarteae
- Socratea exorrhiza* (Mart.) H. Wendl.—Baker 992 (FTG); AF453378, AY543108.
- Arecoideae; Leopoldinieae
- Leopoldinia pulchra* Mart.—Romero 3060 (VEN); AF453355, AY543102.
- Arecoideae; Manicarieae
- Manicaria saccifera* Gaertn.—Henderson s.n. (NY); AF453358, AJ830173.
- Arecoideae; Pelagodoxeae
- Pelagodoxa henryana* Becc.—1988-2933 (K); AJ831321, AJ830135. *Sommieria leucophylla* Becc.—1992-3571 (K); AJ831335, AJ830147.
- Arecoideae; Podocceae
- Podococcus barteri* G. Mann & H. Wendl.—Reitsma 2840 (BH); AF453370, AJ830180.
- Coryphoideae; Livistoneae
- Chamaerops humilis* L.—Lewis 99-012 (BH); AF453339, AY543097.