Workshop on assessments of National Carbon Budgets within the Nordic Region

Current status and sensitivity to changes

Workshop on assessments of National Carbon Budgets within the Nordic Region

Current status and sensitivity to changes

*Kristina Hansen, Aki Koyama, Anne Sofie Lansø, Eva Thorborg Mørk
and Eva Podgrajsek*

Workshop on assessments of National Carbon Budgets within the Nordic Region
Current status and sensitivity to changes
Kristina Hansen, Aki Koyama, Anne Sofie Lansø, Eva Thorborg Mørk and Eva Podgrajsek

ISBN 978-92-893-2597-4
<http://dx.doi.org/10.6027/TN2013-560>
TemaNord 2013:560

© Nordic Council of Ministers 2013

Layout: Hanne Lebech
Cover photo: ImageSelect
Photo: p. 9 Johannes Jansson; p. 15. Johannes Jansson; p. 17 Johannes Jansson;
p. 27 Anne Sofie Lansø; p. 39 ImageSelect; p. 45 ImageSelect; p. 53 ImageSelect;
p. 57 Magnus Fröderberg; p. 61 Magnus Fröderberg

This publication has been published with financial support by the Nordic Council of Ministers.
However, the contents of this publication do not necessarily reflect the views, policies or recommendations of the Nordic Council of Ministers.

www.norden.org/en/publications

Nordic co-operation

Nordic co-operation is one of the world's most extensive forms of regional collaboration, involving Denmark, Finland, Iceland, Norway, Sweden, and the Faroe Islands, Greenland, and Åland.

Nordic co-operation has firm traditions in politics, the economy, and culture. It plays an important role in European and international collaboration, and aims at creating a strong Nordic community in a strong Europe.

Nordic co-operation seeks to safeguard Nordic and regional interests and principles in the global community. Common Nordic values help the region solidify its position as one of the world's most innovative and competitive.

Nordic Council of Ministers

Ved Stranden 18
DK-1061 Copenhagen K
Phone (+45) 3396 0200

www.norden.org

Content

Preface.....	7
Introduction	9
Three Nordic research projects	11
Abstracts.....	12
Keynote: Greenhouse gas inventories with focus on Land-Use and Land-Use Change.....	15
1. Observation of carbon sinks and sources	17
1.1 Summary of presentations.....	17
1.2 Abstracts.....	20
2. Modeling the carbon budget.....	27
2.1 Summary of presentations.....	27
2.2 Abstracts.....	29
3. Remote sensing data for carbon balance modeling.....	39
3.1 Summary of presentations.....	39
3.2 Abstracts.....	41
4. Impacts of future climate and land use scenarios.....	45
4.1 Summary of presentations.....	45
4.2 Abstracts.....	47
5. Posters.....	53
5.1 Abstracts.....	53
6. Discussion and concluding remarks	57
6.1 Current state of research.....	57
6.2 Sensitivities and impact	58
6.3 Research improvement	58
6.4 Largest uncertainties	59
6.5 Communication to policymakers.....	59
Workshop program	61
Program.....	61
Participants.....	64

Preface

The Nordic project “Workshop on assessments of the national carbon budget within the Nordic region – Current status and sensitivity to changes” was financed by the Nordic Council of Ministers’ Working Group on Climate and Air Pollution. The establishment of a new global climate agreement under UN climate convention (UNFCCC) is a primary objective for the Nordic countries and to contribute to this objective is of highest priority for the Nordic Working group on Climate and Air Pollution.

Further knowledge on the carbon balance and how emissions of greenhouse gases from anthropogenic activities translate into concentrations of greenhouse gases in the atmosphere could contribute to more cost-efficient and science-based mitigation measures. It would also enhance the possibility to get consensus in the international negotiations for a global agreement from 2020 on the accounting methods of emissions and removals of greenhouse gases from land-use, land-use changes and forestry. However, many uncertainties on the estimation of carbon exchange between the atmosphere and terrestrial and marine ecosystems still remain. An important objective of the workshop was to discuss the challenges on carbon budget estimations and how to improve the communication of results and conclusions of carbon budget research to policy makers.

The Nordic countries could have an important role to play in establishing the basic premises for the negotiation of the new climate agreement to apply from 2020. Individually the Nordic countries have implemented a wide spread of various climate mitigation measures and are therefore well positioned to exchange knowledge and to develop further research co-operation in this field. Joint Nordic analyses and networks can contribute to the results of the negotiations and to follow-up activities at national and regional level.

This workshop is a good example of when Nordic co-operation can give added value to an issue, where further knowledge is needed on a national, regional and global level. It is our hope that this report can make a significant contribution to the Nordic and international co-operation on the efforts to understand the carbon and other greenhouse gas exchanges between land, ocean and atmosphere and enhance the assessment of anthropogenic emissions and removals of greenhouse gases from land-use and land-use changes.

A handwritten signature in black ink, appearing to read 'Reino Abrahamsson', with a stylized flourish at the end.

Reino Abrahamsson

Chairman, Nordic Council of Ministers

Working Group on Climate and Air Pollution

Introduction

In the Nordic Strategy for Sustainable Development an objective is “to work to stem climate change and help prepare our regions and the rest of the world for the changes that will be needed” (Nordic Council of Ministers 2009). To do that, it is essential to understand the carbon balance as this defines how emissions of greenhouse gases, especially carbon dioxide and methane, from anthropogenic activities translate into concentrations of greenhouse gases in the atmosphere and control important climate feedback mechanisms (Houghton et al. 2007). Estimating and mapping carbon sinks and sources are therefore major issues concerning also the agreements of the United Nations Framework Convention on Climate Change (UNFCCC) in the Kyoto protocol. It is needed to ensure maximum cost effectiveness of actions taken to mitigate anthropogenic emissions of greenhouse gases. However, surface exchange of greenhouse gases are highly affected by anthropogenic forcing (changes in land use and management) and climate forcing. Therefore, to understand and predict these impacts using models, it is important to quantify realistically how much carbon is taken up by the terrestrial and marine ecosystems. Thus, detailed knowledge of the exchange of the greenhouse gases between land, ocean and atmosphere is important, and the processes involved must be fully understood and accurately quantified. However, large uncertainties on the estimations of carbon exchange remain.

It is clear that man- and climate driven land use changes impact the greenhouse gas budget, but the interplay between these changes still needs to be understood (Schulze et al. 2009). Improved information on the magnitude and location of natural carbon sources and sinks contrib-

utes to the development of new and improved tools to support politician's decision making in actions taken to mitigate anthropogenic greenhouse gas emissions and to develop management strategies. Thereby, the future environmental legislation can be based on more realistic carbon budget estimations.

Three Nordic research projects aim at estimating the carbon balance at national scale; the Danish ECOCLIM, the Swedish LAGGE and the Finnish SnowCarbo. These three projects are analysing the three major natural carbon reservoirs (terrestrial ecosystems, marine ecosystems and atmosphere) in respect to global warming, with particular focus on the future climate and land management. Scientists from these three projects, other researchers and politicians within this focus area were brought together for a three-day workshop to obtain an overview of the current status and knowledge on assessments of national carbon budgets as well as on projections and sensitivities to future changes.

The workshop was organised in four thematic sessions, one poster session, and one discussion session. The four themes for the sessions were:

- Observations of carbon sinks and sources
- Modeling the carbon budget
- Remote sensing data for carbon modeling
- Impacts of future climate and land use scenario

This report summarizes the presentations and discussions from each session of the workshop.

References

- Houghton, R. A. (2007). Balancing the Global Carbon Budget. *Annu. Rev. Planet. Sci.* 35, 313:347, 10.1146/annurev.earth.35.031306.140057.
- Nordic Council of Ministers (2009). Sustainable Development – New Bearings for the Nordic Region. Nordic Council of Ministers, Copenhagen, Denmark.
- Schulze, E. D., Luyssaert, S., Ciais, P., Freibauer, A., Janssens, I. A., Soussana, J. F., Smith, P., Levin, I., Thiruchittampalam, B., Heimann, M., Dolman, A. J., Valentini, R., Bousquet, P., Peylin, P., Peters, W., Rodenbeck, C., Etiope, G., Vuichard, N., Wattenbach, M., Nabuurs, G. J., Poussi, Z., Nieschulze, J., Gash, J. H., CarboEurope, Team (2009). Importance of methane and nitrous oxide for Europe's terrestrial greenhouse-gas balance. *Nature Geoscience*, 2, 12, 2009, 842-850, doi:10.1038/ngeo686.

Three Nordic research projects

The global carbon cycle, its sinks and sources, and how different future scenarios will affect the carbon balance are still not properly understood. ECOCLIM (Ecosystems surface exchange of greenhouse gases in an environment of changing anthropogenic and climate forcing), LAGGE (Landscape scale investigation of greenhouse gas exchange) and Snow-Carbo are three research projects from Denmark, Sweden and Finland respectively, which aim to shed some light into these unknowns.

ECOCLIM

ECOCLIM is an ongoing research project (2011–2014) and is funded by The Danish Council for Strategic Research. The project is coordinated by Aarhus University and other partners involved are Copenhagen University, Roskilde University, Technical University of Denmark, and ClimateLab. ECOCLIM's overall objective is collect new data at the air-sea interface and develop models that quantifies surface exchange of greenhouse gases (GHG) between terrestrial, marine and atmospheric components. With this system of models, an estimation of the annual surface flux of CO₂ in Denmark will be made and the effects of management scenarios affecting both land use and nitrogen load as well as climate scenarios will be studied.

LAGGE

The LAGGE project (2010–2013) is supported financially by The Swedish Research Council for Environment, Agricultural Science and Spatial Planning, BEGG, Vetenskapsrådet and University of Gothenburg and is coordinated by University of Gothenburg. Other involved partners in LAGGE are Linköping University, Ghent University, Belgium, Stockholm University, Örebro University, Royal Institute of Technology, Lund University, and Uppsala University. One aim of the Swedish project LAGGE is to quantify GHG balances on a landscape scale. This will include exchange between land-atmosphere, land-water, and water-atmosphere of CO₂, CH₄, N₂O and DOC/DON. Another aim of LAGGE is to estimate an overall watershed net carbon and GHG balance at different scales with new measurements and technologies from both available data and ongoing measurements and to link this data with models.

SnowCarbo

SnowCarbo (2009–2012) is funded by The European Commission Life+ program and is coordinated by the Finnish Meteorological Institute (FMI). Other project partners involved are the Finnish Environment Institute (SYKE) and Le Laboratoire des Sciences du Climat et l'Environnement (CEA.LSCE). The SnowCarbo project's main objective is to implement and demonstrate a new approach for mapping net carbon balance in Europe and northern Eurasian regions. In this new approach different information sources are combined; snow evolution, phenology, land cover, and CO₂ fluxes and concentrations. Local in situ observations will be combined with global Earth observation satellite data and land cover class information in a new way.

Abstracts

Ecosystem services in ECOCLIM

Lise Lotte Sørensen,¹ E. Bøgh, J. Bendtsen, E. Dellwik, T. Friborg, C. Geels, K. Hansen, A. Ibrom, E. T. Jørgensen, M. K. Sejr, K. Hansen, A. S. Lansø and C. A. Skjøth.

Surface exchange of greenhouse gases (GHG) between land/sea and the atmosphere is an important climate feedback mechanism. The processes involved in exchange of GHG needs to be fully understood and accurately quantified in order to increase reliability of climate projections and ensure that actions initiated to reduce anthropogenic GHG emissions are sustainable and not destructive to existing ecosystem services. Therefore it is important to address i.e. land use change in relation to the regulating services of the ecosystems, such as carbon sequestration and climate regulation. At present, the surface exchange of GHG in Denmark and Danish waters is unknown and qualified predictions of the effect of changes in climate or anthropogenic activities are not possible. In ECOCLIM it is a goal to quantify the natural sinks and sources controlling the atmospheric concentration of CO₂ and CH₄ thus a thorough understanding of the ecosystem processes controlling the uptake or emissions of

¹ Aarhus University, Department of Environmental Science, Frederiksborgvej 399, 4000 Roskilde, Denmark.

GHG is fundamental. Here we present ECOCLIM in the context of ecosystem services and the experimental studies within ECOCLIM which will lead to an enhanced understanding of Danish ecosystems.

A fertile peatland forest does not constitute a major greenhouse gas sink

Leif Klemetsson,² Meyer A., Tarvainen L., Nousratpour A., Björk, R.G., Ernfors M., Kasimir Klemetsson, Å., Lindroth A., Räntfors M., Rütting T., Wallin G., and Weslien P.

Afforestation has been proposed as a strategy to mitigate the often high greenhouse gas (GHG) emissions from agricultural soils with high organic matter content. However, the carbon dioxide (CO₂) and nitrous oxide (N₂O) fluxes after afforestation can be considerable, depending predominantly on site drainage and nutrient availability. Studies on the full GHG budget of afforested organic soils are scarce and hampered by the uncertainties associated with methodology. In this study we determined the GHG budget of a spruce-dominated forest on a drained organic soil with an agricultural history. Two different approaches for determining the net ecosystem CO₂ exchange (NEE) were applied: for the year 2008, direct (eddy covariance) and an indirect (analyzing the different components of the GHG budget), so that uncertainties in each method could be evaluated. The annual tree production in 2008 was 8.2 ($\pm$ 1.7) t C ha⁻¹ yr⁻¹ due to the high levels of soil nutrients, the favorable climatic conditions and the fact that the forest was in its optimum growth phase. N₂O fluxes were determined by the closed chamber technique and amounted to 3.3 ($\pm$ 2.4) t CO₂ eq ha⁻¹ yr⁻¹. According to the direct measurements from the eddy covariance technique, the site acts as a minor GHG sink of -4.1 ($\pm$ 2.6) t CO₂ eq ha⁻¹ yr⁻¹. This contrasts with the NEE estimate derived from the indirect approach which suggests that the site is a net GHG emitter of 3.3 ($\pm$ 10.1) t CO₂ eq ha⁻¹ yr⁻¹. Irrespective of the approach applied, the soil CO₂ effluxes counter large amounts of the C sequestration by trees. Due to major uncertainties involved in the indirect approach, the direct approach is considered the more reliable tool. As the site was in its optimum growth stage, i.e. the rate of C sequestration was at its maximum

² University of Gothenburg, Department of Biological and Environmental Science, Tellus, PO Box 460, SE 405 30 Göteborg, Sweden.

and will decrease with forest age; it will probably become a GHG source once again as the trees mature. Since forests in their younger stages are usually GHG sources or have no effect on GHGs, the overall sink potential of this afforested nutrient-rich organic soil is probably limited to only a short period.

Feasibility of Space-Borne Earth Observation Data to Investigate Carbon Cycle

Jouni Pulliainen,³ Ali Nadir Arslan, Kimmo Rautiainen, Anna Kontu, Tiina Markkanen, Tuula Aalto, Mika Aurela, Kristin Böttcher, Juha Lemmetyinen

Feasibility of space-borne microwave radiometers and optical sensors to support the mapping of CO₂ and CH₄ fluxes and annual balances is discussed investigated. First, satellite data retrievals and observations with tower-based reference sensors of satellite instruments are compared with available in situ data. This is carried out in order to find out the relevance of these specific satellite data products, such as day of snow melt information. Secondly, approaches to apply satellite data for mapping purposes are demonstrated. The investigated satellite instruments include current operational and Earth exploring systems such as SMOS, MODIS, AATSR, AMSR-E and SSM/I.

³ Finnish Meteorological Institute, Arctic Research, Erik Palmenin Aukio 1, 00101 Helsinki, Finland.

Keynote: Greenhouse gas inventories with focus on Land-Use and Land-Use Change

Steen Gyldenkærne⁴ and Ole-Kenneth Nielsen

The international regulation of issues related to climate change, e.g. emission of greenhouse gases, is done within the United Nations Framework Convention on Climate Change (UNFCCC). There are different requirements in terms of reporting of greenhouse gas emissions and only 37 countries have reduction commitments under the Kyoto Protocol (KP). The reporting under the UNFCCC and accounting under the KP only includes anthropogenic emissions. For most activities this is straight forward such as fuel combustion, agriculture and waste treatment. However, when it comes to land-use and land-use changes, it is more difficult to assess which emissions/removals are anthropogenic.

The precision of the submitted inventories for land-use and land use-change varies among countries depending on national circumstances (knowledge and resources for the inventory preparation). Over the years the submitted inventories has improved substantially.

⁴ Aarhus University, Department of Environmental Science, Frederiksborgvej 399, 4000 Roskilde, Denmark.

The agreed definition by the IPCC and adopted by the KP is to use a managed land proxy, i.e. all emissions/removals associated with managed land-use are to be included in the inventories to UNFCCC and in the accounting under the KP regardless that some can be argued not to be anthropogenic.

To meet the reporting requirements under the KP extensive work has been carried out in Denmark (the SINKS project) to elaborate a land-use matrix for 1990 onwards. In addition several variables used in the calculation of emissions/removals have been updated based on the SINKS research project. This has particularly improved the emission estimates from soils. The results have also shown a very high variability between years which should be taken into account in the negotiations for future reduction commitments.

1. Observation of carbon sinks and sources

1.1 Summary of presentations

To obtain a full understanding of the carbon balance and how greenhouse gases influence the climate at present and in the future, monitoring is needed. Monitoring carbon sinks and sources includes comprehensive experimental work across a wide range of ecosystems in the Nordic region covering both marine and terrestrial ecosystems. In session 1 “Observations of Carbon sinks and sources,” up-to-date findings of carbon dioxide (CO_2) and methane (CH_4) exchange between different types of ecosystems and the atmosphere as well as their biological and physical controlling processes were reported.

Greenhouse gas exchange between surfaces and the atmosphere is controlled by processes occurring over different spatial and temporal scales. Therefore, various observational methods covering different time and spatial scales are used when monitoring carbon sinks and sources. In the keynote presentation by A. Rutgersson, the question of whether we should care about variations in time and space concerning observations of carbon sinks and sources was addressed. The most direct method to estimate air-water carbon fluxes is the ecosystem scale Eddy-Covariance (EC) method. By using EC measurements, variations can be measured with relatively high temporal resolution, being a great advantage when investigating the mechanisms controlling the exchange. However, topographic complexity in surrounding landscapes influences

such measurements. E. Dellwik presented result from a modeling system that combines the flow model (SCADIS) that takes heterogeneities into account and a footprint calculator. The model is being developed to estimate the flux footprint of EC measurements and was tested for the Danish complex Risø Tower site.

Aquatic ecosystems as lakes and oceans are found to have significant effect on the carbon balance. On global scale the oceanic uptake is approximately 25–27% of the anthropogenic carbon emissions (Le Quere et al. 2013, Denman et al. 2007), while lakes might reduce the land carbon sink by 25% through CH₄ emissions (Bastviken et al. 2011). Furthermore, coastal ecosystems play a significant role linking the terrestrial, oceanic and atmospheric carbon reservoirs. They are often very biological active systems and therefore account for an important air-sea exchange of CO₂ (Borges et al. 2006). However, when calculating exchange of carbon between coastal zones and the atmosphere there exist uncertainties related to limited knowledge about physical and biological parameters and processes. And these processes occur over different time and spatial scales depending on i.e. meteorological conditions and biological processes. Recent micrometeorological observations from the Swedish Lake Tämnen were presented by E. Podgajsek and show diurnal cycle of methane, with high fluxes during night and low during day most likely controlled by diurnal variations of the convective mixing in the lake. Likewise, E. T. Mørk presented results from micrometeorological observations from the semi-enclosed sea, Kattegat, which indicated that physical stratification of the water column worked to separate high pCO₂ levels from the deeper waters.

In the Arctic region issues concerning climate changes, and especially temperature changes, are very important. Northern permafrost regions contain large stocks of soil organic carbon (Tarnocai et al. 2009), but the Arctic is very sensitive to temperature increase especially during growing season (Meehl et al. 2007). The continued warming may cause a reduction in the Arctic carbon sink and may even switch to a source. The surface temperature increase is predicted to be stronger in high latitudes and has for the Arctic in the period 1957–2006 been twice the global average. M. Lund pointed out the importance of long-term monitoring, as the Greenland Ecosystem Monitoring (GEM) program that has been very useful in this region. Further monitoring programs and collaborations such as the Arctic Science Partnership (ASP) have been established with the aim of monitoring long-term ecosystem atmosphere exchange of greenhouse gases and climate change effects and feedbacks in the Arctic and M. Lund announced that more programs are already planned. H. Njuabe presented

results from two weeks observations of CO₂ fluxes for 10 tundra sites from Subarctic to high Arctic. They showed that all sites were net sinks of CO₂ during the peak season; however, there is also a large spatial variability across the sites. These variations could be caused by differences in air temperature and leaf area index (LAI). This means that climate changes will influence these areas differently, which needs to be taken into consideration when predicting carbon sinks and sources under changing climate conditions. Furthermore, methane is a greenhouse gas which has been increasingly studied over the past decades due to its high global warming potential. Northern wetlands are thought to be a major natural source. Observations of atmospheric methane fluxes of a wet mire and a shallow lake in the Abisko region presented by M. Jammet shows large differences in emissions variability, indicating the importance of different environmental controls.

Finally, ice covered lakes are currently not considered in annual carbon budgets. However, ice cover duration must be seen as a defining aspect of high latitude regions as it works to regulate climate by i.e. the albedo and influencing the lake-atmosphere gas exchange (Prowse et al. 2011). B. Denfeld presented observations of pCO₂ dynamics for 335 lakes in Sweden that show that Nordic lakes build up CO₂ during ice cover which accumulates and might be highest shortly before the ice melts. It is therefore suggested to take ice cover duration of lakes into account when estimating annual carbon budgets in the Nordic region.

From the research presented in session 1, it is clear that acknowledging what scales of time and space our observations are representing is important in order to interpret data and improve our understanding of controlling processes on the carbon sinks and sources.

References

- Bastviken, D., Tranvik, L. J., Downing, J. A., Crill, P. M., and Enrich-Prast, A. (2011). Freshwater Methane Emissions Offset the Continental Carbon Sink. *Science*, 331, 6013, 50, doi: 10.1126/science.1196808.
- Borges, A.V., Schiettecatte, L.-S., Abril, G., Delille, B., and Gazeau, F., (2006). Carbon dioxide in European coastal waters. *Estuarine, Coastal and Shelf Science*, 70, 3, 375–387, doi:10.1016/j.ecss.2006.05.046,
- Denman, K. L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., da Silva Dias, P. L., Wofsy, S. C., and Zhang, X. (2007). Couplings Between Changes in the Climate System and Biogeochemistry. In: *Climate Change (2007). The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M.Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Le Quéré, C., Andres, R. J., Boden, T., Conway, T., Houghton, R. A., House, J. I., Marland, G., Peters, G. P., van der Werf, G. R., Ahlström, A., Andrew, R. M., Bopp, L., Canadell, J. G., Ciais, P., Doney, S. C., Enright, C., Friedlingstein, P., Huntingford, C., Jain, A. K., Jourdain, C., Kato, E., Keeling, R. F., Klein Goldewijk, K., Levis, S., Levy, P., Lomas, M., Poulter, B., Raupach, M. R., Schwinger, J., Sitch, S., Stocker, B. D., Viovy, N., Zaehle, S., and Zeng, N. (2013). The global carbon budget 1959–2011. *Earth Syst. Sci. Data*, 5, 165–185, doi:10.5194/essd-5-165-2013.

Meehl, G.A., T.F. Stocker, W.D. Collins, P. Friedlingstein, A.T. Gaye, J.M. Gregory, A. Kitoh, R. Knutti, J.M. Murphy, A. Noda, S.C.B. Raper, I.G. Watterson, A.J. Weaver, and Z.-C. Zhao (2007). Global Climate Projections. In: *Climate Change (2007). The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Tarnocai, C., J. G. Canadell, E. A. G. Schuur, P. Kuhry, G. Mazhitova, and S. Zimov (2009). Soil organic carbon pools in the northern circumpolar permafrost region. *Global Biogeochemical Cycles*, 23, doi:10.1029/2008GB003327.

1.2 Abstracts

1.2.1 **Key note: Observations of carbon sinks and sources**

Anna Rutgersson,⁵ Erik Sahlee, Eva Podgrajsek, Maria Norman.

Significant components of the marine carbon systems relates to the uptake/release to the atmosphere. Air-sea (air-lake) exchange is controlled by the difference in concentration between the water surface and the atmosphere, but also by the efficiency of the transfer. For emissions to the atmosphere of for example methane additional processes needs to be taken into account (like ebullition). The most direct method to estimate air-water carbon fluxes is by eddy-covariance (EC) measurements. By using EC measurements variation can be measured with relatively high temporal resolution, being a great advantage when investigating the mechanisms controlling the exchange.

⁵ Uppsala University, Villavägen 16, 75267 Uppsala, Sweden.

1.2.2 Diurnal cycle of lake methane flux

Eva Podgrajsek⁶

In this study the Eddy covariance method has been used for measuring methane fluxes from a lake situated in central Sweden. At several occasions during the long time measuring campaign (autumn 2010-autumn 2012), a diurnal cycle of methane, with high fluxes during night and low during day, has been captured.

During these occasions the difference between air and water temperature ($\Delta T = T_a - T_w$) also displayed an diurnal cycle, with ΔT being positive during day and negative during night with the corresponding change in the sensible heat flux i.e. negative during daytime and positive during nighttime.

The high nighttime methane fluxes could be explained with this difference in air and water temperature, which will cool the water surface during night, creating convective mixing in the lake, while during daytime the water will be stably stratified. Temperature measurements made at different vertical levels in the lake water confirm this water stratification. The nighttime convective mixing may act to disturb the bottom water, triggering methane ebullition events and bringing methane rich water up to the surface, which can be emitted to the atmosphere.

With this study we want to emphasize the necessity of introducing also complex physical processes when estimating air-water exchange fluxes and also measure methane fluxes not only on few occasions during daytime but also during night and for longer measuring periods.

1.2.3 Biological and Physical control of $p\text{CO}_2$ in a semi-enclosed sea: The Kattegat

Eva Thorborg Mørk,⁷ Peter Anton Stæhr and Mikael Sejr

Coastal ecosystems play a significant role in linking the terrestrial, oceanic and atmospheric carbon reservoirs and since they are often very biological active systems the coastal waters account for an important air-sea exchange of CO_2 . But because of the complexity of the coastal ecosystems

⁶ Uppsala University, Department of Earth Sciences, Villavägen 16, 75267 Uppsala, Sweden.

⁷ Aarhus University, Department of Bioscience, Frederiksborgvej 399, 4000 Roskilde, Denmark.

the exchange of CO₂ in these waters are connected with large uncertainties. Hence, understanding the controls of the air- sea CO₂ exchange in coastal waters is critical when developing robust carbon budgets.

We investigate the spatial distribution of water pCO₂ and the processes responsible for the observed variations in the shallow coastal sea Kattegat in August 2011. The results of the vertical variation show that pCO₂ most often increases with measuring depth and is controlled by the salinity and thereby the stratification of the water column. The variation of surface water pCO₂ show, that the lowest pCO₂ is observed in the northern part of Kattegat and in the shallow water inlet Smålandsfarvandet. The surface water pCO₂ is neither correlated with salinity, temperature or chlorophyll a concentration, which is the most common parameters describing open water pCO₂. The variations of surface water pCO₂ are instead connected to dissolved organic matter and water column light attenuation.

The presented results indicate that the vertical variation of pCO₂ is controlled by physical processes like stratification and mixing of saline bottom water from the North Sea and fresh surface water from the Baltic Sea. The surface water pCO₂ is estimated to be more controlled by biological processes with primary production controlled by light availability and the dissolved organic matter being oxidized by heterotrophic populations.

1.2.4 Risø tower measurements and analysis in the ECOCLIM project

Ebba Dellwik⁸ and Andrey Sogachev

In 2012, the 120m tall Risø tower was instrumented with concentration and flux measurements of carbon dioxide. A first analysis of the generated datasets is presented.

The area around the Risø tower is heterogeneous and a flow model, that can take the heterogeneities into account, together with a footprint calculator were used to estimate the flux footprint of the measurements.

The flux footprint was calculated for several measurement levels on the Risø mast, using the common wind direction from the south-west. The calculated footprints were narrow over the Roskilde fjord and wider

⁸ Technical University of Denmark, Wind Energy, Frederiksborgvej 399, 4000 Roskilde, Denmark.

over the land surface, due to the different turbulent regimes over these two types of surface.

1.2.5 Terrestrial monitoring of greenhouse gas exchange in Greenland

Magnus Lund,⁹ Torben R. Christensen, Birger U. Hansen, Mikkel P. Tamstorf

Within the Greenland Ecosystem Monitoring (GEM) programme, an integrated monitoring and long-term research programme on ecosystems and climate change effects and feedbacks in the Arctic, we are monitoring the exchange of greenhouse gases between Arctic terrestrial ecosystems and the atmosphere. In 1995, Zackenberg Basic in high Arctic northeast Greenland was established as a long-term monitoring site of Arctic ecosystem effects and feedback processes induced by climate change. A similar monitoring programme, Nuuk Basic, was established in 2007 in a low Arctic west Greenland site. At these locations, land-atmosphere exchange measurements of CO₂ and CH₄ in dry and wet areas using ICOS level instrumentation have been implemented as an integral part of the monitoring programmes, and several studies on land-atmosphere greenhouse gas exchange have been published. During 2012, another site on Disko Island (Arctic station) in west Greenland was established, and in 2014 a site in northernmost Greenland (Station Nord) will be installed. Within the Arctic Science Partnership (ASP), an extensive Greenlandic-Danish-Canadian research collaboration, further attention will be drawn towards studying gradients of ecosystem processes and climate change effects in Greenland as well as expanding current monitoring efforts towards full annual coverage. Recently, a GEM financed PhD project has been initiated to facilitate up-scaling of ecosystem processes, such as snow cover, vegetation greening and terrestrial carbon cycling, from local to regional scale.

⁹ Aarhus University, Frederiksborgvej 399, 4000 Roskilde, Denmark.

1.2.6 Spatial variability in peak season net ecosystem exchange across the Arctic tundra

Herbert Njuabe,¹⁰ Magnus Lund, Mikkel Tamstorf

We have investigated the spatial variability in carbon dioxide (CO₂) fluxes on 10 tundra sites from the Subarctic to high Arctic during the peak of the growing season. Flux data has been collected using the micro-meteorological eddy covariance technique. Sites include dry Arctic heaths, mixed tussock and polygonal tundra; and mires.

Comparison is based on common light response curve (LRC) parameterization during most productive two-week period in summer. The functional response of CO₂ exchange to photosynthetic photon flux density (PPFD) has been interpreted from the LRCs and their parameters used to investigate similarities and differences in aspects of ecosystem CO₂ exchange.

This study shows that there is a high variability between Arctic tundra sites. LRC parameters ranged between 1.55 and 8.04 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of CO₂ (fcsat); 0.78 and 3.85 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of CO₂ (Rd) and 0.01 to 0.06 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of CO₂ per $\mu\text{mol m}^{-2} \text{s}^{-1}$ of PPFD (alpha). All sites were net sinks of CO₂ during the peak season. Site variations could be explained mainly by differences in air temperature and leaf area index (LAI). Thus, climate change effect will vary considerably within the Arctic tundra due to heterogeneity and differences in climate characteristics, vegetation composition and phenology.

¹⁰ Aarhus University, Frederiksborgvej 399, 4000 Roskilde, Denmark.

1.2.7 Ecosystem-scale methane dynamics in a subarctic mire landscape, Northern Sweden: on the importance of lake emissions

Mathilde Jammot,¹¹ Thomas Friborg, Patrick Crill

Methane is a greenhouse gas which has been increasingly studied over the past decades due to its high global warming potential. Northern wetlands are thought to be a major natural source of methane, and the role of open water areas on methane dynamics is still an open question.

Surface carbon exchanges, with a focus on methane emissions, are monitored continuously in the Stordalen mire (68°20'N, 19°03'E) since late May 2012. The area is underlain by discontinuous permafrost, resulting in a dynamic landscape comprising dry palsa mounds elevated by permafrost, water-saturated areas with deep summer active layers, fen areas without permafrost, and open water bodies. Using high frequency eddy covariance (EC) observations, we take an ecosystem-scale approach to analyze emissions from different surface types and integrate the effect of landscape heterogeneity on methane dynamics.

The EC tower is situated at the shore of the wettest part of the mire, bordered by a shallow lake. Thanks to a stable bimodal east-west wind direction, we are able to clearly distinguish methane emissions between the vegetated productive fen and the lake. Results from 2012 growing season indicate much higher fluxes from the wetland quadrant: average methane emissions of $7.4 \pm 2.7 \text{ mg m}^{-2} \text{ h}^{-1}$ from the fen were six times higher than those observed at the lake ($1.2 \pm 2.6 \text{ mg m}^{-2} \text{ h}^{-1}$). We observe large differences in emissions variability between the two ecosystems, reflecting a difference in environmental controls.

Ongoing chamber measurements conducted in the vicinity of the tower validate the average summer emissions. Low lake fluxes were unexpected given recent publications about subarctic lakes. Eventually, yearly measurements will help assessing annual methane and carbon budgets of these different ecosystems. Permafrost in the area is expected to further degrade. Therefore, the fen is likely to represent a future state of the mire, that is to say a stable methane source.

¹¹ University of Copenhagen, Center for Permafrost, Øster Voldgade 10, 1350 København K, Denmark.

1.2.8 Under ice CO₂ dynamics in boreal lakes (aquatic ecosystems): Implications of a changing cryosphere

Blaize Denfeld,¹² Gesa Weyhenmeyer, Pirrko Kortelainen

It is well established inland waters have the capacity to efficiently transform carbon, resulting in substantial emissions of carbon dioxide (CO₂) to the atmosphere. However, annual CO₂ emission budgets do not account for the ice-cover period, which is a defining aspect of high latitude regions. Thus, there is a need to better understand and quantify pCO₂ dynamics during ice cover; in particular since climate change will undoubtedly alter the formation and duration of ice on lakes. We investigated under ice pCO₂ for a diverse set of 335 lakes in Sweden and related them to lake water chemistry, lake morphometry and ice duration data. We also measured under ice pCO₂ in a single lake ecosystem to evaluate vertical and horizontal pCO₂ variations. Our results show that pCO₂ under ice variations are dependent on nutrient conditions in the lake water and lake morphometry. Further, under the ice cover CO₂ concentration profile between bottom waters and surface waters was less sharp compared to the open water period. We conclude that the accumulation of CO₂ below lake ice is primarily driven by nutrient conditions and less by ice cover dynamics.

¹² Uppsala University, Norbyvägen 18D, 75236 Uppsala, Sweden.

2. Modeling the carbon budget

2.1 Summary of presentations

Within the research area of modeling the carbon cycle a wide range of research fields are present (Denman et al 2007). Many parameters influence the carbon cycle, thus it can be very complicated to model. Therefore, modelers have to decide on model complexity and model parameterization. The degree of complexity and parameterization is furthermore linked to the spatial scale of the model i.e. site specific, regional etc. Model wise the different spatial scales are connected by up- or down scaling; however the detailed description in site specific models might be parameterized in regional models. In Session 2 an introduction to some of the research fields within carbon cycle modeling were given starting with modeling of marine systems followed by terrestrial modeling both of forest and agricultural ecosystems, and concluded by regional modeling of the carbon cycle.

Many presentations in Session 2 showed how observations are important for modelers. Real time observations have several usages for developing and evaluating the performance of models such as tuning of model parameters, model validation and model inversions. B. Tupek uses an extensive data base of biomass of needles and leaves to estimate litter in forests, while A. Koyama uses observation in an inversion to estimate regional methane fluxes. R. Jensen uses observations of leaf chlorophyll and leaf photosynthesis to estimate V_{cmax} and parameterize a model to simulate LAI and GLAI. Of course the same observations cannot be used for both parameter settings and

validation – independent data is needed for that, and as such there is always a wish for more observations.

Input data are essential for model simulations and often weather data are one of the needed inputs. However, the choice of selected weather data can alter the results greatly as shown by A. Lethonen, where the soil carbon exchange varied significantly with the change of meteorological input data. Thus, modelers should genuinely consider which inputs to apply for their models.

A challenge for most models is the heterogeneity of many land surfaces, which can make the estimation of carbon exchange between the terrestrial biosphere and the atmosphere difficult both for footprint analysis and models of different spatial scales. This was shown for CH₄ emissions from wetlands by A. Budishchev.

The many branches of carbon cycle modeling are all contributing with a part of the puzzle to determine the carbon cycle. Models are needed, as observations have a very sparse spatial representation, and from observations alone you would not be able to get the full picture of the carbon budget. It is essential to understand the current processes influencing the carbon cycle in order to be able to predict future changes. Models are continuously being improved, and better representations of the physical, chemical and biological processes governing the carbon cycle are implemented. As shown by keynote speaker C. Heinze natural variation in atmospheric CO₂ has existed as long as our records go back in time (400,000 years) following the glacial cycles. However, our system is irreversible and with the present anthropogenic interference it is not certain that the atmosphere of the Earth ever again will experience a level of CO₂ being equal to that of the pre-industrial level (Archer 2005, Denman 2007). Thus, it is of greatest importance to further develop models for carbon cycle studies as to determine where the current situation will take us in the future.

References

- Archer, D., (2005). The fate of fossil fuel CO₂ in geologic time. *J. Geophys. Res.*, 110(C9), C09S05, doi:10.1029/2004JC002625.
- Denman, K. L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., da Silva Dias, P. L., Wofsy, S. C., and Zhang, X. (2007). Couplings Between Changes in the Climate System and Biogeochemistry. In: *Climate Change (2007). The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

2.2 Abstracts

2.2.1 Key note: Towards the quantification of marine carbon budgets with models

Christoph Heinze¹³

The ocean is important for carbon cycling primarily due to three main factors: 1. The Ocean is a huge carbon reservoir with a relative short turnover time. 2. Carbon dioxide in sea water is effectively dissociated inorganically into other substances. 3. Marine plankton is keeping the surface ocean carbon dioxide concentration at a lower level than a lifeless ocean would provide. For natural carbon dioxide variations, such as during the glacial-interglacial cycles, the biological carbon cycling presumably played a key role as shown by sediment core data. For the uptake of human produced carbon dioxide, the inorganic uptake by the ocean (dissolution, dissociation, transport, and mixing) plays the most prominent role. Ocean acidification, changes in ocean circulation, and other factors can, however, also lead to a modification of organic carbon cycling. On intermediate to long time scales, the ocean provides the most important uptake reservoir for anthropogenic carbon dioxide. The principle modeling concepts for simulating the marine carbon cycle will be described. An overview about present projects on establishing the global and regional marine carbon budgets will be presented. The inclusion of the coastal zone and the land-ocean transfer of matter provide specific challenges for Earth system modellers. New and emerging data syntheses provide a basis for combining models with observations in a systematic way in order to arrive at best possible estimates for the fate of anthropogenic carbon in the Earth system.

¹³ University of Bergen, Geophysical Institute & Bjerknes Centre for Climate Research, Allégaten 70, 5007 Bergen, Norway.

2.2.2 *CO₂ uptake in the inner Danish waters*

Jørgen Bendtsen¹⁴

The ocean takes up a significant part of the global anthropogenic CO₂ emissions. For example, about half of the total CO₂ emissions from fossil fuel combustion and cement-manufacturing production has been dissolved in the ocean since the industrialisation, equivalent to about 160 Pg C. Currently the ocean absorbs about one third of the total global anthropogenic CO₂ emissions of ~8 Pg C yr⁻¹ and this global carbon ocean sink is expected to continue in the near future, and even increase because of the increasing atmospheric pCO₂ level. However, the ocean CO₂ sink is characterized by large spatial and temporal variations with areas of large CO₂ out-gassing or uptake. Coastal areas also show a large spatial and temporal variation and, for example, in our region the current estimates varies from an annual CO₂ uptake of -1.2 mol m⁻² yr⁻¹ in the North Sea area to a net outgassing of about 1.6 mol m⁻² yr⁻¹ in the central Baltic Sea. The inner Danish waters, i.e. Kattegat, the Belt Seas and Sounds, are a part of the transition zone between the North Sea and the low-saline Baltic Sea and therefore located in this large spatial gradient of ocean carbon uptake. As a part of the on-going ECOCLIM project we analyse the CO₂ dynamics in the region through regional ocean modeling and preliminary results of the CO₂ uptake in the area will be presented. Finally the implications of further global warming on the future CO₂-uptake in the region will be discussed.

2.2.3 *Predicting damage vulnerability based on ground-based observations of forests and weather*

Mikko Peltoniemi,¹⁵ Petteri Muukkonen, Sanna Härkönen, Aleksi Lehtonen, Annikki Mäkelä, Seppo Nevalainen, Seppo Neuvonen

Climate has direct influence on forest carbon and water balances, but it also influences forest damages. Populations of some pests can grow enormously when climate and other factors turn favourable, and they can cause wide-

¹⁴ ClimateLab, Symbion Science Park, Fruebjergvej 3, Box 98, 2100 Copenhagen East, Denmark.

¹⁵ Finnish Forest Research Institute, METLA, Jokiniemenkuja 1, 01301 Vantaa, Finland.

spread forest damages, which have further consequences on forest carbon balances and growth. It is expected that the influence of forest damages to large-scale carbon budgets will increase with climate change.

We are building a model-data system, which draws from ground-based estimates of forest structure and weather in order to make inferences about forest carbon balances and damage vulnerability. Forest-inventory data provides the basic structural information about forests, which we scale wall-to-wall Finland using satellite images, and then use as input to a forest carbon-water balance model. Forest inventory data, together with permanent sample plots, also provides information about forest damages.

In this presentation, we will present on-going work that we are conducting in order to estimate vulnerability of forests to damages associated with low or high soil moisture. We will present our preliminary attempts to predict damage vulnerability of forests based on permanent sample plot and forest inventory data, topographic map data and weather. We further evaluate the potential of model predicted drought in explaining the occurrence of forest damages.

Climforisk, 2011–2014 LIFE09 ENV/FI/000571 Climate change induced drought effects on forest growth and vulnerability, www.metla.fi/life/climforisk

2.2.4 The uncertainty of modeled soil carbon stock change for Finland

Aleksi Lehtonen,¹⁶ Juha Heikkinen

Countries should report soil carbon stock changes of forests for Kyoto Protocol. Under Kyoto Protocol one can omit reporting of a carbon pool by verifying that the pool is not a source of carbon, which is especially tempting for the soil pool. However, verifying that soils of a nation are not a source of carbon in given year seems to be nearly impossible. The Yasso07 model was parameterized against various decomposition data using MCMC method. Soil carbon change in Finland between 1972 and 2011 were simulated with Yasso07 model using litter input data derived from the National Forest Inventory (NFI) and fellings time series.

¹⁶ Finnish Forest Research Institute, Jokiniemenkuja 1, 01301 Vantaa, Finland.

The uncertainties of biomass models, litter turnover rates, NFI sampling and Yasso07 model were propagated with Monte Carlo simulations. Due to biomass estimation methods, uncertainties of various litter input sources (e.g. living trees, natural mortality and fellings) correlate strongly between each other. We show how original covariance matrices can be analytically combined and the amounts of simulated components reduce greatly. While doing simulations we found that proper handling correlations may be even more essential than accurate estimates of standard errors.

As a preliminary results, from the analysis we found that both Southern- and Northern Finland were soil carbon sinks, coefficient of variations (CV) varying 10%–25% when model was driven with long term constant weather data. When we applied annual weather data, soils were both sinks and sources of carbon and CVs varied from 10%–90%. This implies that the success of soil carbon sink verification depends on the weather data applied with models. Due to this fact IPCC should provide clear guidance for the weather data applied with soil carbon models and also for soil carbon sink verification. In the UNFCCC reporting carbon sinks of forest biomass have been typically averaged for five years – similar period for soil model weather data would be logical.

2.2.5 *A synthesis on carbon budget in a Danish, temperate deciduous forest*

Jian Wu,¹⁷ Klaus Steenberg Larsen, Leon van der Linden, Claus Beier, Kim Pilegaard, Andreas Ibrom

A synthesis on carbon budget in a temperate deciduous forest, Sorø (Zealand, Denmark) was performed by combining all available data from eddy covariance, chamber, suction cups and biometric measurements. The net ecosystem exchange of CO₂ (NEE), soil respiration, tree growth, litter production and the leaching of dissolved organic carbon were independently estimated and used to calculate other unmeasured ecosystem carbon budget (ECB) components, based on mass balance equations. This provided a complete assessment of the carbon storage and allocation within the ecosystem. The results showed that this tempera-

¹⁷ Technical University of Denmark, Frederiksborgvej 399, 4000 Roskilde, Denmark.

ture deciduous forest was moderately productive ($259 \pm 41 \text{ g C m}^{-2} \text{ yr}^{-1}$) with both high rates of gross primary production (GPP, $1881 \pm 95 \text{ g C m}^{-2} \text{ yr}^{-1}$) and ecosystem respiration (Re, $1624 \pm 197 \text{ g C m}^{-2} \text{ yr}^{-1}$). Approximately 62% of the gross assimilated carbon was respired by the living plants, while 21% was contributed to the soil as litter production, the latter balancing the total heterotrophic respiration. The remaining 17% was either stored in the plants (mainly as aboveground biomass) or removed from the system as wood production. The SOC was considered to be unchanged over the period of observation, given the high degree of uncertainty associated with the small loss detected ($33 \pm 85 \text{ g C m}^{-2} \text{ yr}^{-1}$). The results of the study yield valuable information that helps checking the plausibility of the multiple datasets in an ecosystem context. The plausibility analyses reported here, using simple mass balance equations to constrain the uncertainty of estimated ecosystem components of ECB, may be widely applicable among other experimental sites.

2.2.6 Variation of pine, spruce, and birch litterfall data exceeds modeling inputs for soil carbon stock change

Boris Ľupek¹⁸, Aleksi Lehtonen, Tatu Hokkanen, Liisa Ukonmaanaho, Pekka Nöjd, Eero Nikinmaa

Prerequisite for precise estimates of soil carbon stock change is a correct model function; moreover, exact model inputs. Systematic errors in model inputs propagate into large offsets with soil carbon stocks and their changes on regional level estimates.

Here, we aim to test the differences between the existing Pine, Spruce and Birch litterfall data and litterfall estimates based on models. We applied biomass models and litter-turnover rates for litter trap sites and then compared the quantity of measured and modeled foliage litter.

The litterfall data showed in comparison to foliage estimates larger variation, which is presumably driven by differences in climate and environment, a subject of further investigations.

¹⁸ University of Helsinki, Department of Forest Sciences, Latokartanonkaari 7, 00014 Helsinki, Finland.

2.2.7 Validation of PEATLAND-VU methane emission model using eddy covariance observations and footprint modeling

Artem Budishchev,¹⁹ Y. Mi, J. van Huissteden, G. Schaepmann-Strub, F. J. Parmentier, G. Fratini, L. Belelli, A. Gallagher, A. J. Dolman, T. C. Maximov

Several methane emission models were developed recently. However, validation of these models is currently performed using data collected by closed chamber method. While being a simple and relatively inexpensive way to measure fluxes, this method has a number of limitations such as small measurement area and low temporal resolution. Furthermore, chamber measurements are unsuitable to register ebullition events, which can have a significant influence on observed fluxes. Eddy covariance (EC) measurements on the other hand provide data on a half-hourly basis and cover larger areas, while being non-intrusive way to measure fluxes and account for ebullition.

In this study, we present a validation of PEATLAND-VU methane emission model using eddy covariance data, collected in summer periods of 2008 and 2009 at the Indigirka lowland site in Eastern Siberia. High resolution vegetation map together with Kormann & Meixner, 2001 footprint model were used to produce footprint weighted methane fluxes. Consecutively, these values were compared to EC data.

Results of the comparison show good agreement of the modeled fluxes and EC flux. The model was able to reproduce daily variations of the fluxes albeit some mismatch.

¹⁹ VU University Amsterdam, De Boelelaan 1085, 1081HV Amsterdam, The Netherlands.

2.2.8 Temporal variation of leaf nitrogen: a key parameter for photosynthesis modeling

Rasmus Jensen²⁰

Leaf photosynthetic capacity is limited by leaf nitrogen concentration (Na) and leaf chlorophyll content (Cab) and these parameters often show a characteristic seasonal dynamic. When simulating CO₂ exchange over vegetated surfaces, model outputs are sensitive to leaf photosynthetic capacity.

In the present study we investigate the temporal change in nitrogen and thus photosynthetic capacity and the implications when parameterizing a CO₂ exchange model. Field measurements of leaf photosynthesis (LI-6400, LICOR Inc.), leaf reflectance (SPAD-502, Minolta), and LAI (LAI-2000, LICOR Inc.) were conducted on agricultural fields in Western Denmark during one growing season. The leaf photosynthesis measurements provided the basis for estimating photosynthetic capacity. SPAD measurements and LAI was measured with a higher spatial and temporal resolution. SPAD readings were calibrated against Cab and Na analyzed on leaf material in the laboratory and later correlated to photosynthetic capacity. These data were used to parameterize a coupled photosynthesis and stomatal model that was run for the growing season 2012 to estimate NEE. In parallel fluxes of greenhouse gases were continuously measured by eddy covariance systems at the field site, providing the basis for estimating the exchange of energy, water vapour, and CO₂ on canopy scale. We compare modeled NEE with the measured CO₂ fluxes.

The presented approach is shown to provide an efficient field sampling method for model parameterization with high temporal and spatial resolution and a physiological basis for scaling fluxes from leaf level to canopy scale.

²⁰ University of Copenhagen, Department of Geosciences and Natural Resource Management, Øster Voldgade 10, 1350 København K, Denmark.

2.2.9 Modeling of regional CO₂ balance in Life+ project SnowCarbo

Tiina Markkanen,²¹ Tea Thum, Jouni Susiluoto, Niina Puttonen, Tuula Aalto.

In SnowCarbo project a regional modeling framework was developed to estimate current land ecosystem CO₂ balance in region covering Fennoscandia and Baltic countries. The modeling framework consists of a Regional Climate Model (RCM), REMO, and a land surface scheme (LSS) JSBACH. The models are one-way coupled so that REMO provides the meteorology for the surface processes of LSS.

REMO is derived from the operational weather forecast model of the Deutscher Wetter Dienst (DWD), therefore it has been extensively evaluated for its capability to predict the synoptic scale meteorological phenomena. In this project the RCM have been driven with daily initialization from boundary conditions so that the climate remains close to the actual weather conditions. The boundary data is from ERA-INTERIM. As REMO model does not presently consider carbon cycle, JSBACH is used to simulate CO₂ exchange in the present work.

JSBACH is the LSS of general circulation model (GCM) ECHAM6 of MPI-Hamburg. The role of a LSS in GCM is to provide the lower boundary condition to the atmosphere for 1) the vertical diffusion scheme; 2) radiation scheme and 3) hydrological cycle such that the surface energy balance and water balance are closed. Stomatal functioning, that is the most crucial control of water vapor exchange between the vegetation and the atmosphere, constrains the photosynthetic CO₂ exchange as well. Thus LSS JSBACH, that treats the surface water cycle by considering the physiological response of vegetation to the climatic variables, is readily able to predict the photosynthetic CO₂ exchange rate. Moreover, in order to produce a reliable net ecosystem exchange of CO₂ (NEE), the allocation of carbon into various pools in soil and vegetation and the decomposition of these storages are described in JSBACH.

We present the land CO₂ balances estimated by the modeling framework for years 2001–2009 and discuss its reliability, practical applicability and future developments.

²¹ Finnish Meteorological Institute, P.O.Box 503, 00101 Helsinki, Finland.

2.2.10 Methane surface flux estimations by inverse modeling: CarbonTracker Data Assimilation Shell

Aki Koyama,²² Tuula Aalto, Peter Wouters, Leif Backman, Maarten Krol, Karolina Stanislawska.

In recent years, inverse models are becoming common tools to estimate greenhouse gas surface fluxes by source categories, such as anthropogenic, natural fire, biosphere and oceanic. One of such models, CarbonTracker Data Assimilation Shell (CTDAS), has been further developed to derive optimized estimates of global methane surface fluxes, using the forward and bottom-up estimates as priors. EDGAR v4.2, LPJ-WhyMe, GFED and NOAA GMD Carbon Cycle GHG MBL References are used as the prior emissions for anthropogenic, biosphere, fire and ocean respectively. Further, prior for termites added. The priors are transported by chemistry-transport model (TM5) to match time and location where atmospheric observations are available, and two of those (anthropogenic and biosphere) are optimized through the data assimilation system (Ensemble Kalman filter). The latest version of TM5, driven by ERA-Interim ECMWF meteorological fields, with a two-way nested European domain and $1^\circ \times 1^\circ$ grid over high Northern latitudes (up to 70°N), is used to focus the estimates over Europe and European Boreal regions. Atmospheric methane sink with OH is included with contributions from tropospheric and stratospheric OH, Cl and O(1D). The atmospheric observations are collected from WDCGG. The forward model result for year 2007 is compared to the atmospheric observations in Europe in order to prepare and validate the modeling system for inverse approach.

²² Finnish Meteorological Institute, P.O.Box 503, 00101 Helsinki, Finland.

3. Remote sensing data for carbon balance modeling

3.1 Summary of presentations

Remote sensing data have various applications to carbon balance modeling such as: model inputs, model validation, coordinating tools for mobile measurements, regional analysis, upscaling of local measurements, and several others (Anderson and Norman 2007, Rosenqvist et al. 2003, Turner et al. 2004). They all play important roles in regional and large spatial scale estimations, which cannot be discovered by only local measurements.

The invited speaker, K. Böttcher discussed the impacts of land cover maps as inputs to a one-way coupled REMO-JSBACH model, developed in SnowCarbo Life+ project. The land cover maps influence carbon budget estimations because most of the natural carbon cycle on land is controlled by the vegetation. The amount and types of vegetation determine the balances between sinks and sources, and affect the temporal variation of the carbon exchange. Therefore, a good land cover map is essential to carbon balance modeling. The comparison illustrated the large impacts on estimated carbon balances by using different land cover maps. Also, the validation of model results using MODIS normalized difference vegetation index (NDVI), water index and snow cover area data illustrated the difficulties in estimating snow cover seasons over the Nordic region. However, applying remote sensing data as an adjustment of model parameters could potentially improve the model.

Using high resolution land cover data generally improve model estimations, especially for heterogeneous land. T. Becker used remote sensing data of a coordinate reference system (CRS) in the estimation of local land surface map. He used a system, where a camera is placed in an airborne mobile platform, taking pictures of study area (Salmisuo). Combining those pictures, a very fine resolution map has been created. Such map cannot be obtained by satellite observations from space, and is useful in detailed local carbon cycle estimation.

Another application is discussed by A. Gallagher, where the remote sensing data were used for pond detection of the Siberian arctic tundra. The satellite images of reconnaissance satellite KH-11 KENNAN (also known as Key Hole) image from 1977 and earth observation satellite GeoEye-1 image from 2010 were compared to determine the changes in the number of ponds around the area. Although the two images differ much in resolutions, the comparison illustrated a clear increase in the number of ponds, which indicates the thawing of permafrost. As permafrost contains the vast amount of carbon, its changes has impacts on greenhouse gas emissions. The changes in the land also affect local vegetation.

The carbon cycle is not only dependent on natural processes, but also on human activities. A. Nordbo discussed the use of MODIS global land cover data for upscaling the local urban measurements to continental urban levels. From the remote sensing data, the fraction of natural area were determined and meta-analysis of data from 14 flux measurement sites were used as scaling factor for estimation of Net Urban Ecosystem Exchange (NUE). The results showed good agreements with inventory data.

It is important to be aware that remote sensing data does not reveal everything. Old data tend to have low resolutions, making it difficult to use for past analysis and small regional scale estimation. Furthermore, clouds have been problematic in retrieving the data; the satellite data are limited for the Nordic region in winter because the region is mostly clouded during the season.

To conclude, the applicability of remote sensing data to carbon balance modeling is diverse. Data such as land and snow cover map are especially useful for the Nordic region, where the natural land is extensive, and the carbon balance is especially sensitive to snow cover and melting of permafrost. Other remote sensing data are also useful in modeling carbon exchange of heterogeneous lands, such as swamps and cities. In summary new and improved understanding of regional to global scale carbon balance is obtained through the use of remote sensing data.

References

- Anderson M.C., Kustas W.P., and Norman J.M. (2007). Upscaling flux observations from local to continental scales using thermal remote sensing. *Agronomy J.*, 99, 1, 240–254, doi:10.2134/agronj2005.0096S.
- Rosenqvist, Å., Milne, A., Lucas, R., Imhoff, M., and Dobson, C. (2003). A review of remote sensing technology in support of the Kyoto Protocol. *Environmental Science and Policy*, 6, 5, 441–455, doi:10.1016/S1462-9011(03)00070-4.
- Turner, D. P., Ollinger, S. V., and Kimball, J. S. (2004). Integrating remote sensing and ecosystem process models for landscape- to regional-scale analysis of the carbon cycle. *BioScience*, 54, 573–584, doi:10.1641/0006-3568(2004)054[0573:IRSAEP]2.0.CO;2.

3.2 Abstracts

3.2.1 *Key note: Remote sensing data for carbon balance modeling*

K. Böttcher,²³ M. Törmä, P. Härmä, T. Markkanen, O.-P. Mattila, S. Metsämäki, A. Arslan, T. Aalto, J. Pullianen

In the framework of the project SnowCarbo (<http://snowcarbo.fmi.fi>), we investigated the use of remote sensing data in support of carbon balance modeling. The work focussed on (i) the implementation of up-to date land cover information and (ii) the evaluation of modelled phenology and snow cover. For the modeling of annual carbon balances in Northern Europe, the land surface model JSBACH (Jena Scheme for Biosphere-Atmosphere) of Max Planck Institute (MPI) in Hamburg was applied in an instantaneous form. JSBACH was forced with regional meteorological data in hourly time resolution that was generated with the regional climate model REMO (MPI, Hamburg). Era-INTERIM data from the European Centre for Medium-Range Weather Forecasts (ECMWF) was used as boundary data for the regional model.

Land cover information affects both the regional climate (REMO) and the carbon cycle (JSBACH) and alternative land cover data sets (CORINE Land Cover, GlobCover version 2.2) were evaluated and compared to the standard model version (Global Land Cover Characteristics Database by USGS). Additionally, allocated surface parameters (e.g. forest ratio) for

²³ Finnish Environment Institute, Mechelininkatu 34 A, 00251 Helsinki, Finland.

land cover categories were evaluated and revised using local estimates from Finland.

The performance of the JSBACH model regarding the start of the growing season (hereafter referred to as Growing Season Start Date, GSSD) and the melting of snow cover in spring were assessed with satellite observations from Moderate Resolution Imaging Spectrometer (MODIS). The satellite retrieval method for GSSD in evergreen coniferous forest was calibrated to in situ observations of CO₂ fluxes at eddy covariance (EC) sites in Finland. A fixed fraction of peak growing season gross primary production (GPP) was used as a threshold value for the growing season onset from EC measurements. Modelled GSSD from regional GPP predications by JSBACH showed good spatial correspondence, but a delay (about 2 weeks) compared to satellite observations.

3.2.2 *Climate driven permafrost degradation and the resulting increase in pond cover impacts on the Siberian tundra greenhouse gas emissions*

Angela Gallagher,²⁴ A. Budishev, B. Li, R. Petrov, J. van Huissteden, M.M.P.D.Heijmans, T.C. Maximov, A.J. Dolman.

Rising global temperatures threatens the stability of continuous permafrost environment, resulting in the release of previously frozen carbon, creating a positive climate feedback from terrestrial systems. The resulting permafrost degradation will impact on both a large scale, expansion of thaw lakes, and small-scale features, such as superficial pond formation and mass wasting.

The increased occurrence of thaw ponds and their impact on vegetation is been studied at the Kytalyk research station, located in Indigirka lowlands, Northeast Siberia. This area is located on the drained bed of an Early Holocene thaw lake. The area is characterised by the presence of low palsas (flat ice mounds), covered with mosses and *Betula nana*. The edges of these palsas are subject to frequent thawing, creating shallow ponds with decaying palsa vegetation.

Comparison, using high resolution satellite images from 1977 (American Keyhole project image) and 2010 (Geoeye), showed increased oc-

²⁴ Vrije Universiteit Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands.

currence of thaw ponds over a 33 year period. Flux measurements from a selection of these ponds showed elevated emission of CO₂ and CH₄. Dead *Betula nana* produced fluxes of 106.84mg CO₂ m⁻² hr⁻¹ and 3.58mg CH₄ m⁻² hr⁻¹ in the summer of 2011.

However, a decrease of GHG fluxes occurs when *Carex* and *Eriophorum* (sedges) vegetation invades these ponds. The CH₄ emission from sedges is still high, 9.1mg CH₄ m⁻² hr⁻¹ in 2010 and 3.9mg CH₄ m⁻² hr⁻¹ in 2011, but this is compensated by rapid CO₂ uptake. It is therefore likely that GHG emission from this type of shallow permafrost degradation is strongly influenced by ecosystem recovery rates.

3.2.3 The use of high resolution aerial photography to estimate source area contributions to heterogeneous ecosystem fluxes

Thomas Becker,²⁵ J. Sievers, I. Forbrich, J. Schneider, M. Gazovic, L

Accurate determination of carbon balances in heterogeneous ecosystems requires an estimate of the spatial extent of the different surface types. In peatlands, small-scale variations of surface structures, vegetation cover and hydrology influence the carbon exchange on the microform scale. To extrapolate plot data, detailed knowledge of the small-scale variations is required. But even for the evaluation of ecosystem-scale measurements estimates of the landcover are needed. Here, we evaluate the influence of landcover maps of differing high resolution on the estimation of vegetation period net ecosystem exchange of a heterogeneous peatland.

The studies have been conducted in the near-pristine boreal peatland complex Salmisuo located in North Karelia, eastern Finland. An eddy covariance system was running all year round in 2006 and 2007, while closed chamber measurements were performed from May–October 2006 and May–July 2007. Both chamber measurements and eddy covariance measurements were modelled based on irradiance and temperature functions. Aerial photographs have been obtained utilising a blimp and a digital camera, resulting in a surface classification of the 200m radius fetch of the eddy covariance system. To evaluate dependencies of the ground reso-

²⁵ Aarhus University, Frederiksborgvej 399, 4000 Roskilde, Denmark.

lution with respect to coverage changes of land use classes, we resampled to resolutions of 13cm (best), 60cm and 100cm. The resulting maps are used to extrapolate the chamber measurements and in combination with an analytical footprint model (Kormann & Meixner) to estimate the 30min source area fractions of the different microform types.

We will demonstrate the advantages of using very high resolution aerial photographs in combination with an analytic footprint model to achieve more accurate model results at higher resolution. Additionally, we assess the relative merits of using extrapolated modelled chamber measurements and eddy covariance measurements in modeling gas fluxes in the study area.

3.2.4 Fraction of natural area predicts CO₂ budgets of cities

*Annika Nordbo*²⁶

Cities are the largest source of greenhouse-gases, especially that of CO₂. We perform a meta-analysis of annual CO₂ budget estimates from 14 cities. The estimates are based on direct eddy-covariance flux measurements that take into account sources within a few hectares around a measurement tower. We show that the fraction of natural land area can explain 84% of the intercity variation in the CO₂ budget. Thus, this dependency is used as a proxy for predicting CO₂ budgets around the world based on satellite-driven land cover data. The upscaling procedure is verified by extracting 56 case study cities from the mapping and by comparing their inventory-based estimates to our estimates based on land cover. The correspondence is high ($r^2=0.72$) and thus the fraction of natural area can be corroborated to be a valid proxy for CO₂ budgets of cities. Furthermore, the annual budget is zero when the fraction of natural area is about 80%.

²⁶ University of Helsinki, Department of Physics, P.O. Box 64, 00014 Helsinki, Finland.

4. Impacts of future climate and land use scenarios

4.1 Summary of presentations

In order to assess the future evolution of the carbon cycle, scenarios are needed. In the framework of IPCC various climate scenarios have been conducted. They span a wide range in which the actual future climate change most likely will lie (Meehl et al. 2007, Moss et al. 2010).

Many scientist working within research areas where natural responses to climate change are investigated are dependent on climate scenarios in order to assess the future changes. However, due to the large spread in the climate scenarios it can be troublesome to select which and how many climate scenarios to use in a study, since scientists can be limited by both accessibility to the climate data, time and money. The question is not just which scenario to use, but also from which Global Circulation Model (GCM) these data should come. Among GCMs large variations can be seen for the same applied scenario, and as explained by invited speaker Martin Drews one cause for these variations is the different land use models used in the GCMs. Thus, it is important to consider both the scenarios and GCMs used when applying such data in a study.

Another important consideration is how to present the results – should it be a mean across scenarios, a mean across GCMs, should it be communicated as just one figure or a range, and with or without uncertainties? These reflections were all pointed out by Tuomo Kalliokoski during his presentation on climate induced change in Finnish forest for which the use of climate scenarios are very important.

There is no doubt that a changing climate will induce changes in ecosystems, and with this changing climate a need to change our habits and our perspective on land use management exists. For this land use scenarios can be very useful and for which the same considerations as climate scenarios can be conducted. Examples of change in habits are management changes of ecosystems such as forests and agricultural sites, but also the idea of biomass utilization for energy, sustainability and carbon storage could be considered as presented by Kim Pilegaard.

Assessment of climate induced changes on ecosystems is most often investigated by models, but at the heathland of Brandbjerg an experimental site has been built. Here the effect of three parameters is being tested – enhanced CO₂, higher temperature and drought, alterations which all are likely in the future climate. The influence from these variables was found not necessarily to be additive revealing the nonlinearities and complexity in our natural system and climate.

The sinks from terrestrial ecosystem have continued to grow with increasing anthropogenic emissions. It is, however, uncertain how efficient the sinks will continue to be in the future. The uncertainties lies within errors in the estimation of land-use changes, ecosystems possible recovery from past disturbance or induced change in metabolism caused by alterations in CO₂, nitrogen (N), temperature and water. As mentioned by Camilla Geels fertilization by N has widely been used throughout the 20th century, but as its harmful effect on natural ecosystem was discovered legislation has decrease the use. Nevertheless, the decreased input of N might reduce ecosystems ability to uptake CO₂.

Climate and land-use scenarios are important tools for scientist in this field of research. But one should keep in mind that these scenarios do not tell exactly what will happen in the future, rather they are made to give possible pathways of the future development. No one in the scientific community can tell exactly what will be happening in the future. Scientist can only give qualified guesses based on the assumptions made in their modeling framework.

References

- Meehl, G. A., Stocker, T. F., Collins, W. D., Friedlingstein, P., Gaye, A. T., Gregory, J. M., Kitoh, A., Knutti, R., Murphy, J. M., Noda, A., Raper, S. C. B., Watterson, I. G., Weaver, A. J., and Zhao, Z.-C. (2007). Global Climate Projections. In: *Climate Change (2007): The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P., and Wilbanks, T. J. (2010). The next generation of scenarios for climate change research and assessment. *Nature* 463(7282), 747–756, doi:10.1038/nature08823.

4.2 Abstracts

4.2.1 **Key note: Climate models and land surface**

Martin Drews²⁷ and Jens Hesselbjerg Christensen

The land surface constitutes an important component in climate models. For example the terrestrial surface plays a key role in dividing the available energy between sensible and latent heat fluxes, it determines whether water drains or remains available for evaporation, it influences the surface fluxes of carbon, and it defines the surface albedo. Due to the relatively coarse resolution of coupled atmosphere-ocean circulation models (AOGCM), however, land processes are only described in such models in a fairly simplistic manner. Higher resolution regional climate models are generally better equipped to resolve the land surface, but many problems remain and there is general agreement that the land surface component in climate models should be improved. For example Christensen and Boberg (2012) has suggested that the temperature dependent climate projection deficiencies in CMIP5 models and in ENSEMBLES RCM's are linked with the ability of models to capture complex feedbacks accurately, in particular land surface-atmosphere interactions. Likewise, Wei et al (2010) argue about the importance of land models for climate simulations. Recent developments in land surface schemes in climate models include the Noah land surface scheme in WRF, the LPJ-GUESS ecosystem model and the LSM advanced land surface scheme used in the COSMO-CLM regional climate model.

²⁷ Climate Change and Sustainable Development, Technical University of Denmark, Risø Campus Frederiksborgvej 399, 4000 Roskilde, Denmark.

References

- Boberg, F., J.H. Christensen. Overestimation of Mediterranean summer temperature projections due to model deficiencies, 2012, *Nature Climate Change* 2, 433–436(2012)
- Christensen, J.H., F. Boberg. Temperature dependent climate projection deficiencies in CMIP5 models, *Geophysical Research Letters*, Volume 39, Issue 24 (2012)
- Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.). *Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, 2007, Cambridge University Press, Cambridge, United Kingdom
- Wei, J., P.A. Dirmeyer, Z. Guo, L. Zhang, V. Mishra. How much do different land models matter for climate simulation? Part I: Climatology and Variability, 2010, *Journal of Climate* 23 (11), 3135–3145(2010)

4.2.2 Biomass utilization for energy, sustainability, carbon storage, and sinks

*Kim Pilegaard*²⁸

There are few possibilities to mitigate climate change by actually removing CO₂ from the atmosphere. Use of biomass for energy is generally not CO₂ neutral since some energy is needed for the production and conversion of biomass. One possibility to improve the situation is BIO-CCS, where biomass utilization is combined with carbon capture and storage. Another possibility is to store carbon in the soil in the form long-lived C-compounds such as biochar and ash. When growing biomass for energy-purpose it should be carefully assessed how much benefit there is, when all the necessary inputs are taken into account. Also it is important to assess the stocks created. From a climate point of view long-term storage of C in the soil as seen in old forests might be more beneficial than high uptake in above ground biomass in short rotation crops.

The terrestrial sink for carbon has been increasing over some decades, but might be vulnerable due to climate change and extreme climate events. It is demonstrated how the heat-wave in Europe in 2003 led to a net release of carbon from terrestrial ecosystems equal to 4 years of normal net carbon sequestration.

²⁸ Department of Chemical and Biochemical Engineering, Technical University of Denmark, 2800 Kongens Lyngby, Denmark.

4.2.3 Effects of elevated CO₂, warming and summer drought on the carbon balance in a Danish heathland after seven treatment years – results from the CLIMAITE project

Klaus Steenberg Larsen,²⁹ Per Ambus, Claus Beier, Andreas Ibrom, Johannes Ransijn, Inger Kappel Schmidt, Jian Wu

In a Danish heathland co-dominated by heather (*Calluna vulgaris*) and grasses (*Deschampsia flexuosa*) we simulated realistic future climate scenarios in a full-factorial design of elevated atmospheric CO₂ (510ppm), increased temperatures (0.5–1.5 °C) and intensified summer drought events (4–6 weeks per year). Treatments were initiated in 2005. Using manual chamber techniques, we measured soil respiration (SR), ecosystem respiration (ER) and net ecosystem exchange of CO₂ (NEE) and determined gross ecosystem photosynthesis (GEP) as NEE – ER. We also monitored carbon losses in the form of dissolved organic carbon (DOC) in leached soil water.

Across all combinations of treatments with elevated CO₂, SR rates increased by 20–30%, whereas GEP rates increased by <5% leading to a net loss of carbon under elevated CO₂ compared to the carbon balance under ambient CO₂. Elevated CO₂ did not significantly affect above-ground ecosystem respiration rates or below-ground DOC leaching. The warming treatment had only small effects on both GEP and respiration rates, i.e. slightly decreased uptake rates by photosynthesis and slightly increased loss rates by respiration, resulting in increased overall net carbon losses from the system. In contrast, the drought treatment led to reduced rates of both photosynthesis and respiration, while the net balance was almost unaffected.

The drought treatment therefore exhibited the smallest change in climate change feedback (i.e. no change in NEE), whereas both warming, but especially elevated CO₂ treatments overall showed a surprising positive feedback to climate change even after seven years of treatments. Furthermore, the effects of elevated CO₂ interacted with the other climate drivers. The results highlight that not all ecosystems may be expected to increase their net carbon uptake in a future CO₂ enriched at-

²⁹ Technical University of Denmark, ECO Centre, Department of Chemical and Biochemical Engineering, Frederiksborgvej 399, 4000 Roskilde, Denmark.

mosphere especially because other climate drivers may significantly affect the final ecosystem response.

4.2.4 Climate change induced changes in gross primary production in Finland

Tuomo Kalliokoski,³⁰ Mikko Peltoniemi, Annikki Mäkelä

During this century, mean annual temperature may rise 2–6 °C and precipitation increase 10–20% in Finland. These changes combined with higher atmospheric CO₂-concentration have yielded several tens of percentages increase in growth of trees in some modeling studies. However, the local and regional level climate change scenarios derived from Global Circulation Models (GCM) includes lot of uncertainty. Timing of precipitation is particularly difficult to predict and possible drought spells during growing seasons may influence the expected growth increase of trees. Further development of both regional climate and growth models is required to capture these effects.

Here we present preliminary results of on-going study in which we will estimate the change of the Finnish forests gross primary production (GPP) due to climate change. Implemented climate scenarios are constructed on basis of SRESB1, SRESAB1 and SRESA2 emission scenarios and predictions of altogether of 14 different GCMs. Based on the GCM predictions, the daily mean change of four variables, temperature, precipitation, vapour pressure and global radiation, have been estimated for regular 10 x 10 km grid of Finland for three periods, 2011–2040, 2041–2070 and 2071–2100.

The gross primary production of forests will be estimated with PRELES model, which has been developed and parameterized for the prediction of GPP, evapotranspiration (ET) and soil water balance in boreal conditions. First preliminary results show up to 40% increase in GPP and over 10% increase in ET at whole country level.

³⁰ University of Helsinki, Department of forest sciences, Latokartanokaari 7, 00014 Helsinki, Finland.

4.2.5 Past, present and future changes in the atmospheric input of Nitrogen to the biosphere

Camilla Geels,³¹ Ellermann, T., Hansen, K. M., Christensen, J., Hertel, O., Lansø, A. S., Ambelas Skjøth, C., and Brandt, J.

The atmospheric deposition of nitrogen to natural surfaces is an important component in the nutrient cycling at the local to global scale. However, the spatial distribution of the deposition is highly heterogeneous due to the complex nature of sources and especially deposition processes.

International and national agreements and regulations (e.g. the EU National Emission Ceiling (NEC) directive) have through the last two decades led to emission reductions of sulphur, nitrogen oxides and ammonia in Northern Europe. This has led to a decrease in the atmospheric input of Nitrogen (N) through deposition to both land and sea areas. In Denmark measurements show for example a ca. 30% decrease from 1990 to 2008 in the atmospheric N input to the land areas. Emission projections used in the newest IPCC work points in the direction of even lower N emissions towards year 2100. Significant changes can also be foreseen in the future due to projected changes in the climate.

In order to assess past, present and future changes in the atmospheric deposition of N related to local, regional and international emissions a model framework has been setup based on the atmospheric chemistry transport model DEHM (the Danish Eulerian Hemispheric Model). The DEHM model has been driven on past, present and future meteorology projected by the Atmosphere-Ocean Climate model ECHAM5/MPI-OM and anthropogenic emissions prescribed by the RCP4.5 emission scenario.

Assessments of the past and current as well as projections of future N deposition levels and patterns will be discussed. Simulations of the combined effect of both future climate and emission changes (i.e. our best guess of the future), show that the effect of emission changes dominates in most areas. Changes up to +/- 40% are seen across the Northern Hemisphere.

Projections like this are an important input to terrestrial biosphere models including the nutrient cycling and describing the current and future vegetation-atmosphere interactions.

³¹ Aarhus University, National Environmental Research Institute, Frederiksborgvej 399, 4000 Roskilde, Denmark.

5. Posters

5.1 Abstracts

5.1.1 *First step towards a mesoscale model system modeling the carbon cycle and carbon greenhouse gases for Denmark*

Anne Sofie Lansø,³² C. Geels, M. Tum, J. Bendtsen, J.H. Christensen, M. S. Plejdrup, J. Brandt, L. L. Sørensen

As part of the Danish project ECOCLIM (Ecosystems Surface Exchange of Greenhouse Gases in an Environment of Changing Anthropogenic and Climate forcing) a mesoscale model system is under development.

Numerical models are an important tool for studies of the processes related to atmospheric CO₂ and have been used for decades. More recently mesoscale models have been exploited to simulate the exchange of CO₂ between terrestrial ecosystems and the atmosphere at a high spatio-temporal scale. The fine resolution of mesoscale models help capture the heterogeneity of terrestrial ecosystems and its corresponding exchange of CO₂ with the atmosphere. Not many mesoscale studies have included the exchange of CO₂ between marine ecosystems and the

³² Aarhus University, National Environmental Research Institute, Frederiksborgvej 399, 4000 Roskilde, Denmark.

atmosphere in coastal regions. However, global measurements have indicated that open shelves at high latitudes act as sinks for atmospheric CO₂, while inner estuaries and near-shore coastal areas act as sources of atmospheric CO₂.

For the area of Denmark it is of greatest importance to include the marine exchange as Denmark is surrounded by estuaries and coastal shelves. In this study a modeling framework of high spatio-temporal resolution is developed for the area of Denmark and its surroundings with the aim of determining the exchange processes between the atmosphere and terrestrial and marine ecosystems, and the resulting concentration of CO₂ in the atmosphere. The model system is based on the atmospheric model DEHM (The Danish Eulerian Hemispheric Model) using a nest over Denmark with a spatial resolution of 5.6 x 5.6 km. Anthropogenic emissions are taken from the national emission inventory of Denmark aggregated onto an hourly 1x1 km grid according to source regions and CRF sectors. The marine component to be implemented will be based on the COHERENCE ocean model, while the terrestrial exchange will be facilitated by the Biosphere Energy Transfer Hydrology model (BETHY/DLR) operated at the German Aerospace Center. Validation of the simulated atmospheric CO₂ will be enabled by real time measurements from a new Danish tall mast positioned at Risø.

5.1.2 Surface Energy Balance of Scandinavian Peatlands

Pavel Alekseychik,³³ Annalea Lohila, Anders Lindroth, Jurgen Sagerfors, Narasinha Shurpali, Mats Nilsson, Matthias Peichl, Janne Rinne, Timo Vesala

Wetlands, and specifically peatlands, represent an important class of biomes on Earth. Besides covering a major fraction of the land surface, they contain vast amounts of organic carbon, which they have accumulated since the last glacial era. Since wetlands remain undersampled and less understood compared to boreal forests and tundra, some of their features have received insufficient attention. Surface energy balance (SEB) remains one of the most unexplored aspects in the wetland stud-

³³ University of Helsinki, Department of Physics, P.O. Box 64, 00014 Helsinki, Finland.

ies. The problem deserves more effort, since SEB links such important aspects as temperature, hydrology and carbon budget.

There is a concern that the modern anthropogenic activities may drive the climate towards warming, so that the peatlands will get free of permafrost and climatic constraints, emitting their carbon resources into the atmosphere. However, the effects of anthropogenic intervention will not be limited to the altered peatland carbon budgets: various management practices, such as drainage, peat harvesting and afforestation, will also affect peatland-atmosphere energy exchange.

We intend to study SEB in several boreal peatlands, including Siikaneva (two sites of different trophicity), Kalevansuo, Alkkia and Linnansuo in Finland; Degerö Stormyr and Fajemyr in Sweden. The listed sites represent both the natural and managed peatlands, and have together provided more than 30 site-years of data on peatland-atmosphere interactions.

Our study will attempt to explore the potential shifts in the peatland SEB caused by management. To address this question, we will: compare the SEB closure at the said locations; determine the main controllers of the energy balance in peatlands; assess inter-and intra-site variation in SEB and its components; finally, draw a comparison between the boreal peatland and forest SEB. This is a pioneering effort to present a compiled evidence of the boreal peatland energy balances.

5.1.3 Simulating CO₂ fluxes for a Danish beech forest using a coupled photosynthesis and stomatal conductance model – The first step in the development of a coupled NH₃ – CO₂ model

Kristina Hansen,³⁴ R. Jensen, T. Friborg and E. Boegh

Future changes in climate and land use affect the carbon (C) balance of the Danish forests. Particularly, nutrient availability and nitrogen (N) deposition represent uncertainties for the prediction of future greenhouse gas exchange for Danish forests as the mechanisms describing nutritional effects are not well developed in climate and ecosystems models. However, measurements of N fluxes between the forests and

³⁴ Roskilde University, Department of Environmental, Social and Spatial Change, Universitetsvej 1, 4000 Roskilde, Denmark.

atmosphere are limited, and large uncertainties are related to the modeling of the dry deposition velocity (v_d) of ammonia (NH_3). Currently, the parameterisation of the canopy resistance is based on the early empirical Jarvis (1976) approach to describe the stomatal resistance. However, we wish to exchange this description using a more dynamical description such that the impact of nutrient availability and N deposition on CO_2 fluxes can be represented in terms of leaf photosynthetic capacity.

This study presents the first step in the development of an integrated modeling system to simulate CO_2 and NH_3 fluxes for a Danish beech forest which includes the dynamical modeling of the CO_2 flux. A coupled photosynthesis and stomatal conductance model based on the Collatz approach (1991, 1992) has been applied to the beech forest Lille Bøgeskov for the year 2010. The model was adapted to the forest ecosystem based on input parameters from the literature, measured climatic parameters and measured LAI for the forest. The preliminary results show that the coupled photosynthesis and stomatal conductance model matches very well the diurnal variations during the photosynthetic active season, however, the model underestimates the max CO_2 flux both during daytime and night time.

5.1.4 Climate changes in the Nordic region 8-km resolution according to the RCP4.5 and RCP8.5 scenarios

Cathrine Fox Maule³⁵ and P. Thejll

As climate changes the conditions for plant growth in the Nordic region will change. A new downscaling experiment using an 8-km horizontal resolution in the Nordic region has recently been conducted for the two future greenhouse gas emission scenarios RCP4.5 and RCP8.5 at the Danish Climate Centre. Here we will present results of the expected climate changes in the 21st century for temperature, precipitation, cloud cover, sunshine duration and sea surface temperatures according to the HIRHAM regional climate model for both scenarios.

³⁵ Danish Climate Centre, Danish Meteorological Institute, Lyngbyvej 100, 2100 Copenhagen Ø, Denmark.

6. Discussion and concluding remarks

Multiple sensitivities and uncertainties exist in the field of research covering carbon budgeting in the Nordic Regions. During the workshop “Workshop on assessments of the national Carbon budget within the Nordic region – Current status and sensitivity to changes,” an important objective was to discuss these challenges and to improve the ways to communicate results and conclusions to policy makers. A discussion session covered five research areas; terrestrial carbon modeling, measuring terrestrial carbon fluxes, measuring and modeling marine and lake carbon, remote sensing and climate and land use scenarios. From this session and discussion throughout the workshop the following were agreed upon.

6.1 Current state of research

We currently know that lakes are an important source of both CH₄ and CO₂ on a regional scale. We estimate that CO₂ is important because inorganic carbon can be formed from organic carbon in the lakes. However, it is still unknown if the lakes are an important CO₂ source on a global scale. It was agreed upon that we still know very little on exchange of CH₄ and CO₂ in both the lakes and in the marine environment.

The boreal region is fairly well covered with regards to measurements of terrestrial carbon fluxes. However, studies of carbon fluxes in agricultural areas and urban areas are highly underrepresented and

more measurements in these areas are of high interest to the research community.

In modeling the terrestrial carbon exchange, we currently experience a lack of usable input and validation data on a regional level. There exists a deficiency in understanding the key processes controlling the carbon exchange on the plant and stand level. These gaps of knowledge combined with the current up-scaling difficulties leave a requirement of further studies on carbon exchange and the coupling to carbon models.

6.2 Sensitivities and impact

We estimate that the fraction of oceanic carbon uptake will be reduced over time. It is uncertain what will happen in the coastal areas but the current knowledge indicates that the outgassing of carbon will increase in these areas. The primary production in the water phase will likely decrease in the future which will reduce the ability of the marine environment to take up carbon. At the moment very little is known about the sensitivities of the CH₄ exchange in lakes with regards to future climate change but it is known that the impact depends on the latitude of the lakes.

The sensitivities of the terrestrial carbon exchange with respect to the effect of land use changes are investigated through measurements in restored ecosystems, which at the current time are extensively being done in the Scandinavian areas.

At the ecosystem level the terrestrial modeling of the temperature response is handled well compared to the response of CO₂ or H₂O. It is known that the snow cover in the high-latitude regions will diminish in the future; hence the growing season will be longer in terrestrial ecosystems which will have implications for the carbon cycle. What is still unknown is the acclimation of soil carbon stock in the future climate.

6.3 Research improvement

The close connection between the freshwater systems on land and the coastal waters might be of importance to the regional carbon budget; hence we need further investigation on how lakes and ocean research can be connected. Measurements of carbon exchange in oceans and lakes are complicated and expensive to execute and we still need more measurements and improvement of the sensors. It is of high importance that

focus is laid on developing accurate continuous measuring sensors for the water phase, which would add to the coverage of available data.

On the terrestrial measurements, improvements to the instrumentation and a standardisation of measurements and post processing are needed. We need further research on the effect of the variety of eddy covariance footprint areas and we need more consistency in the filtering and gap-filling process in data sets. More interaction and exchange of data and duties between research groups of different profiles would improve our research substantially.

More collaboration between different fields of science will in carbon modelling play an important role in the future research through increased available resources both financial and work force. In the future new questions will arise, creating new needs that might change the former traditions of research.

6.4 Largest uncertainties

Large uncertainties of the future climate scenarios still exist due to the unknown feedback processes in the marine environment. The remineralisation processes are at this point not described to satisfaction and the physical forcing and stoichiometry in the marine and lake ecosystems is very uncertain in the future scenarios.

The link between the aquatic and the terrestrial ecosystems is still not fully understood which may cause large uncertainties. In connection with need of improving research on eddy covariance footprint we see large uncertainties on the eddy covariance footprint models; hence the models need further verification through comparisons of footprint areas and point measurements. The terrestrial measurements of carbon exchange are possibly containing large uncertainties caused by the lack of knowledge of the role of non-turbulent transport in eddy covariance measurements.

6.5 Communication to policymakers

National and international politics are often established through good communication between scientists and policy makers. An example is how scientific work has revealed the rapid increase of greenhouse gases since the Industrial Revolution, and the effect of the greenhouse gases on environment and societies. Close work between the two groups lead to

the Kyoto Protocol, setting international targets on emission reduction. Communication between the policy makers and the scientists is often important in creating new guidance in our societies.

However, the communication sometimes becomes difficult because of a gap between the two groups due to their different perspective. For policy makers it can be hard to assess uncertainties levels, and whether reported results lie reasonable within these levels. For example, when future scenarios are discussed, policy makes preferred one number from scientific studies to base their policy on. Thus, to answer questions such as “how much anthropogenic emissions should be reduced to prevent further global warming,” the quantity “how much” should be reported in the simplest possible manner. Policy makers need clear answers to build their policy upon.

On the other hand, in the scientific community it is important to report the uncertainties because none of the scientific models used for scenario studies are perfect in predicting unknowns. The uncertainties tell us features of the models, such as sensitivities and representativeness. Furthermore, no models are identical and neither are their results. There are many assumptions behind the scientific framework of a given modeling study, and the obtained scientific results are not what will happen, but the estimation of what will happen based on the framework sat for the study. This is an important point for policy makers to remember.

How should the scientists communicate with the policy makers, to avoid misinterpretation of model results and uncertainties connected with them? One way could be to put more emphasis on clarifying the scientific frameworks. It will help the policy makers to understand the meaning of the uncertainties, and the influences the frameworks have over the obtained results, and hopefully at the same time make the policy makers better to recognise the importance of the scientific framework and its related uncertainties.

The question on how best to report actual scientific results have still not been answered – should it be with or without uncertainties and confidence levels, one figure or a range etc.? One should keep in mind the “type of study,” and what question the study tries to answer when assessing how best to pass on scientific results. To conclude, scientific work gives suggestions and of guidance to the policy makers. Let us hope that the two groups continue and improve their communication in the future.

Workshop program

Workshop on assessments of the national Carbon budget within the Nordic region – Current status and sensitivity to changes, 20–22 March 2013, Roskilde University, Denmark.

Program

Wednesday 20 March – Day 1

13.00 –13.30	Arrival and registration
13.30 –13.45	Welcome and practical information
13.45 –14.15	<i>Keynote: Green house gas inventories with focus on Land-Use and Land-Use Change Steen Gyldenkerne</i>
14.15–15.45	Opening Session: Project Presentations <i>Chair: Eva Podgrajsek</i> Ecosystems Surface Exchange of Greenhouse Gases in an Environment of Changing Anthropogenic and Climate forcing (ECOCLIM) <i>Lise Lotte Sørensen</i> Landscape Scale Investigation of Greenhouse Gas Exchange (LAGGE) <i>Leif Klemetsson</i> Overview of project achievements and future prospects (SnowCarbo) <i>Jouni Pulliainen</i>
15.45–16.00	<i>Coffee break</i>
16.00–17.30	Session 1: Observations of carbon sinks and sources <i>Chair: Kristina Hansen</i> <i>Keynote: Observations of carbon sinks and sources Anna Rutgersson</i> Diurnal cycle of lake methane flux <i>Eva Podgrajsek</i> Biological and Physical control of pCO ₂ in a semi-enclosed sea: The Kattegat <i>Eva Thorborg Mørk</i> Risø tower measurements and analysis in the ECOCLIM project <i>Ebba Dellwik</i>
18.30	<i>Dinner at Rådhuskælderen</i>

Thursday 21 March – Day 2

08.30–09.00 *Coffee and roll*

09.00–10.20 Session 1: Observations of carbon sinks and sources (*continued*)

Chair: Kristina Hansen

Terrestrial monitoring of greenhouse gas exchange in Greenland
Magnus Lund

Spatial variability in peak season net ecosystem exchange across the Arctic tundra
Herbert Njuabe

Ecosystem-scale methane dynamics in a subarctic mire landscape, Northern Sweden: on the importance of lake emissions
Mathilde Jammet

Under ice CO₂ dynamics in boreal lakes (aquatic ecosystems): Implications of a changing cryosphere
Blaize Denfeld

10.20–10.40 *Coffee break*

10.40–12.30 Session 2: Modeling The Carbon Budget

Chair: Anne Sofie Lansø

Keynote: Towards the quantification of marine carbon budgets with models
Christoph Heinze

CO₂ uptake in the inner Danish waters
Jørgen Bendtsen

Predicting damage vulnerability based on ground-based observations of forests and weather
Mikko Peltoniemi

The uncertainty of modelled soil carbon stock change for Finland
Aleksi Lehtonen

A synthesis on carbon budget in a Danish, temperate deciduous forest
Jian Wu

12.30 –13.30 *Lunch*

13.30–15.10 Variation of pine, spruce and birch litterfall data exceeds modeling inputs for soil carbon stock change
Boris Tupek

Validation of PEATLAND-VU methane emission model using eddy covariance observations and footprint modeling
Artem Budishchev

Temporal variation of leaf nitrogen: a key parameter for photosynthesis modeling
Rasmus Jensen

Modeling of regional CO₂ balance in Life+ SnowCarbo
Tiina Markkanen

Methane surface flux estimations by inverse modeling: Carbon Tracker Data Assimilation Shell
Aki Koyama

15.10–16.10 Session 3: Posters

First step towards a mesoscale model system modeling the carbon cycle and carbon greenhouse gases for Denmark
Anne Sofie Lansø

Surface energy balance of Scandinavian peatlands
Pavel Alekseychik

	Simulating CO ₂ fluxes for a Danish beech forest using a coupled photosynthesis and stomatal conductance model – The first step in the development of a coupled NH ₃ – CO ₂ model <i>Kristina Hansen</i>
	Climate changes in the Nordic region 8-km resolution according to the RCP4.5 and RCP8.5 scenarios <i>Cathrine F. Maule</i>
16.10–17.30	Session 4: Discussion Session <i>Chair: Eva Thorborg Mørk</i>
19.00	<i>Young scientist networking at Café Satchmo</i>

Friday 22 March – Day 3

08.30–09.00	<i>Coffee and roll</i>
09.00–10.40	Session 5: Impacts On Future Climate And Land Use Scenarios <i>Chair: Anne Sofie Lansø</i> <i>Keynote: Climate models and land surface</i> <i>Martin Drews</i> Climate change induced changes in gross primary production in Finland <i>Tuomo Kallikowski</i> Biomass utilization for energy, sustainability, carbon storage and sinks <i>Kim Pilegaard</i> Effects of elevated CO ₂ , warming and summer drought on the carbon balance in a Danish heath-land after seven treatment years – results from the CLIMATE project <i>Klaus S. Larsen</i> Past, present and future changes in the atmospheric input of Nitrogen to the biosphere <i>Camilla Geels</i>
10.40–11.00	<i>Coffee break</i>
11.00–12.30	Session 6: Remote Sensing Data For Carbon Balance Modeling <i>Chair: Aki Koyama</i> <i>Keynote: Remote sensing data for carbon balance modeling</i> <i>Kristin Böttcher</i> The use of high resolution aerial photography to estimate source area contributions to heterogeneous ecosystem fluxes <i>Thomas Becker</i> Climate driven permafrost degradation and the resulting increase in pond cover impacts on the Siberian tundra greenhouse gas emissions <i>Angela Gallagher</i> Fraction of natural area predicts CO ₂ budgets of cities <i>Annika Nordbo</i>
12.30–13.00	Concluding remarks and closing
13.00–14.00	<i>Lunch</i>
14.00	<i>End</i>

Participants

Alekseychik, Pavel	University of Helsinki
Becker, Thomas	Aarhus University
Bendtsen, Jørgen	ClimateLab
Bøgh, Eva	Roskilde University
Budishchev, Artem	VU University of Amsterdam
Böttcher, Kristin	Finnish Environment Institute
Dellwik, Ebba	Technical University of Denmark
Denfeld, Blaize	Uppsala University
Drews, Martin	Technical University of Denmark
Fox Maule, Cathrine	Danish Meteorological Institute
Gallagher, Angela	VU University Amsterdam
Geels, Camilla	Aarhus University
Gran, Anna	Nordic Council of Ministers
Gyldenkerne, Steen	Aarhus University
Hansen, Kristina	Roskilde University
Heinze, Christoph	University Of Bergen
Ibrom, Andreas	Technical University of Denmark
Jammet, Mathilde	University of Copenhagen
Jensen, Rasmus	University of Copenhagen
Kalliokoski, Tuomo	University of Helsinki
Klemedtsson, Leif	University of Gothenburg
Koyama, Aki	Finnish Meteorological Institute
Lansø, Anne Sofie	Aarhus University
Larsen, Klaus Steenberg	Technical University of Denmark
Lethonen, Aleksi	Finnish Forest Research Institute
Lund, Magnus	Aarhus University
Markkanen, Tiina	Finnish Meteorological Institute
Mørk, Eva Thorborg	Aarhus University
Njuabe, Herbert	Aarhus University
Nordbo, Annika	University of Helsinki
Peltoniemi, Mikko	Finnish Forest Research Institute METLA
Pilegaard, Kim	Technical University of Denmark
Podgrajsek, Eva	Uppsala University
Pulliaainen, Jouni	Finnish Meteorological Institute
Rutgersson, Anna	Uppsala University
Sogachev, Andrey	Technical University of Denmark
Stendel, Martin	Danish Meteorological Institute
Sørensen, Lise Lotte	Aarhus University
Tupek, Boris	University of Helsinki
Wu, Jian	Technical University of Denmark

norden

Nordic Council of Ministers

Ved Stranden 18
DK-1061 Copenhagen K
www.norden.org

TemaNord 2013:560

Workshop on assessments of National Carbon Budgets within the Nordic Region

The three-day workshop organized by the three Nordic research projects; ECOCLIM, LAGGE and SnowCarbo brought together scientists and other actors from Nordic countries to communicate and discuss research on carbon budget estimations in the Nordic region. Through presentations of most recent research in the field and following scientific discussions, the workshop contributed to strengthen the scientific basis of the identification and quantification of major natural carbon sinks in the Nordic region on which integrated climate change abatement and management strategies and policy decisions is formed from.

This report summarizes presentations and discussions from the four thematic sessions; Observations of carbon sinks and sources, Modeling the carbon budget, Remote sensing data for carbon modeling, and Impacts of future climate and land use scenarios and gives an overview of the current status and knowledge on research on assessments of national carbon budgets as well as on projections and sensitivity to future changes in e.g. management and climate change in the Nordic Region.

TemaNord 2013:560
ISBN 978-92-893-2597-4

9 789289 325974