









# Accounting for environmental variation in co-occurrence modelling reveals the importance of positive interactions in root-associated fungal communities

Nerea Abrego<sup>1</sup>  | Tomas Roslin<sup>1,2</sup>  | Tea Huotari<sup>1</sup> | Ayco J. M. Tack<sup>3</sup>  |  
Björn D. Lindahl<sup>4</sup>  | Gleb Tikhonov<sup>5</sup>  | Panu Somervuo<sup>6</sup>  | Niels Martin Schmidt<sup>7</sup>  |  
Otso Ovaskainen<sup>6,8</sup> 

<sup>1</sup>Department of Agricultural Sciences, University of Helsinki, Helsinki, Finland

<sup>2</sup>Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

<sup>3</sup>Department of Ecology, Environment and Plant Sciences, Stockholm University, Stockholm, Sweden

<sup>4</sup>Department of Soil and Environment, Swedish University of Agricultural Sciences, Uppsala, Sweden

<sup>5</sup>Computational Systems Biology Group, Department of Computer Science, Aalto University, Espoo, Finland

<sup>6</sup>Organismal and Evolutionary Biology Research Programme, University of Helsinki, Helsinki, Finland

<sup>7</sup>Arctic Research Centre, Department of Bioscience, Aarhus University, Roskilde, Denmark

<sup>8</sup>Centre for Biodiversity Dynamics, Department of Biology, Norwegian University of Science and Technology, Trondheim, Norway

## Correspondence

Nerea Abrego, Department of Agricultural Sciences, University of Helsinki, Helsinki, Finland.  
Email: nerea.abrego@helsinki.fi

## Funding information

Vetenskapsrådet, Grant/Award Number: 2015-03993; LUOVA graduate school of University of Helsinki; Academy of Finland, Grant/Award Number: 1273253, 250444, 276909, 285803 and 308651; Norges Forskningsråd, Grant/Award Number: 223257; Ella ja Georg Ehrnroothin Säätiö

## Abstract

Understanding the role of interspecific interactions in shaping ecological communities is one of the central goals in community ecology. In fungal communities, measuring interspecific interactions directly is challenging because these communities are composed of large numbers of species, many of which are unculturable. An indirect way of assessing the role of interspecific interactions in determining community structure is to identify the species co-occurrences that are not constrained by environmental conditions. In this study, we investigated co-occurrences among root-associated fungi, asking whether fungi co-occur more or less strongly than expected based on the environmental conditions and the host plant species examined. We generated molecular data on root-associated fungi of five plant species evenly sampled along an elevational gradient at a high arctic site. We analysed the data using a joint species distribution modelling approach that allowed us to identify those co-occurrences that could be explained by the environmental conditions and the host plant species, as well as those co-occurrences that remained unexplained and thus more probably reflect interactive associations. Our results indicate that not only negative but also positive interactions play an important role in shaping microbial communities in arctic plant roots. In particular, we found that mycorrhizal fungi are especially prone to positively co-occur with other fungal species. Our results bring new understanding to the structure of arctic interaction networks by suggesting that interactions among root-associated fungi are predominantly positive.

## KEYWORDS

Arctic, co-occurrence, endophyte, interaction network, mycorrhiza, symbiotic network

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Molecular Ecology* published by John Wiley & Sons Ltd

## 1 | INTRODUCTION

Understanding the role of interspecific interactions in shaping ecological communities is one of the central goals in community ecology (Bailey, Kelsic, & Kishony, 2016; Werner & Peacor, 2003; Wootton, 1994). Species can interact in two ways: directly, when individuals of one species affect the fitness of a second species, or indirectly, when individuals of one species affect the fitness of a second species through their direct interaction with a third species. In interaction networks, such as those formed by plants and microbes in the rhizosphere, direct and indirect interactions play important roles in determining the community composition and fitness of both plants and microbes (Artursson, Finlay, & Jansson, 2006; Frey-Klett, Garbaye, & Tarkka, 2007; Gould et al., 2018; Johansson, Paul, & Finlay, 2004). Furthermore, interactions in the rhizosphere influence ecosystem processes, such as nutrient cycling (e.g., Lindahl, de Boer, & Finlay, 2010; Toro, Azcon, & Barea, 1997). Therefore, unravelling the structure of rhizosphere interactions and in particular disentangling direct and indirect interactions has important implications for understanding both plant diversity and ecosystem functioning.

In this paper, we focus on root-associated fungi, which form highly interactive communities with important effects on host plant fitness (Rodriguez, White, Arnold, & Redman, 2009; Smith & Read, 2008). Root-associated fungi directly and indirectly interact with each other, at the same time as they interact with the host plants. Mycorrhizal fungi facilitate nutrient and water uptake as well as protect roots from pathogens and toxic compounds. In return, they gain direct access to carbohydrates from plant roots (Smith & Read, 2008). A less studied group of root-associated fungi are endophytic fungi, which use living plant tissues as their habitat and may sometimes improve plant fitness (Newsham, 2011; Rodriguez et al., 2009). Yet, root-associated fungal-plant interactions are not always positive, and mycorrhizal and endophytic fungi may also act as parasites (Johnson, Graham, & Smith, 1997; Rodriguez et al., 2009).

Within host plants, root-associated fungi may interact directly or indirectly with each other in a number of ways. Direct interactions among root-associated fungi involve trophic relationships in which some species constitute the resource for others (Lindahl et al., 2010; Tedersoo et al., 2009). Another type of interaction is direct competition during colonization of roots and for root resources (Moeller & Peay, 2016). Root-associated fungi may also interact indirectly, by some species modifying exudate production by plant roots (Ingham & Molina, 1991). However, the outcomes of interactions between fungal species are not always consistent and may vary with the timing of root colonization (i.e., priority effects, see Kennedy & Bruns, 2005; Kennedy, Peay, & Bruns, 2009) or with the abiotic environment (Erland & Finlay, 1992; Mahmood, Finlay, Fransson, & Wallander, 2003).

There is increasing evidence that interactions among co-occurring microbes within host plant individuals affect plant fitness. Fungi and bacteria can modify the nutritional influence of each other on the associated plants (Frey-Klett et al., 2011; Johansson et al., 2004). The so-called "Mycorrhiza Helper Bacteria" stimulate

the symbiotic association between plants and mycorrhizal fungi, ultimately improving plant growth (Founoune et al., 2002; Frey-Klett et al., 2007). Among root-associated fungi, mycorrhizal fungi and endophytes are usually studied separately, and thus their interactions and the effects of their interaction outcomes on plant fitness are largely unknown. Some authors have found positive co-occurrence patterns between endophytic and mycorrhizal fungi and proposed that the positive co-occurrences may mostly be due to them occupying different niches within plant roots (Wagg, Pautler, Massicotte, & Peterson, 2008), or alternatively due to their facilitative interactions (Yamamoto et al., 2014). In contrast, Wearn, Sutton, Morley, and Gange (2012) found that mycorrhizal and endophytic fungi mostly avoided each other within herbaceous plants, and suggested that mycorrhizal fungi and endophytic fungi interact antagonistically by competing for plant roots.

Evidence of interactions among root-associated fungi is best achieved through direct observations in the laboratory. However, root-associated fungal communities are highly species-rich and laboratory experiments are feasible for only a small proportion of the whole root-associated fungal diversity (see Singh, Millard, Whiteley, & Murrell, 2004). An indirect way of assessing the role of interspecific interactions in determining community structure from field observations is to evaluate patterns of co-occurrence (Gotelli, 2000). Studies of co-occurrence in root-associated fungal communities have found co-occurrences to range from predominantly positive (Gorzalak, Hambleton, & Massicotte, 2012; Kennedy, Nguyen, Cohen, & Peay, 2014) to predominantly negative (Koide, Xu, Sharda, Lekberg, & Ostiguy, 2005; Pickles, Genney, Anderson, & Alexander, 2012; Pickles et al., 2010; Wubet et al., 2012). However, and as noted in the studies cited, co-occurrence patterns do not solely reflect interactive associations, but also environmental filtering. If the effect of the environment is not taken into account, positive co-occurrences may reflect that species tend to occur under similar environmental conditions, whereas negative co-occurrence may reflect that species sort according to different environmental niches. To formulate more reliable hypotheses on species interactions based on co-occurrence patterns, it is thus crucial to determine which co-occurrences cannot be explained by the environmental variation (Ovaskainen, Hottola, & Siitonen, 2010).

In this study, we investigated co-occurrences among root-associated fungi, asking whether mycorrhizal and endophytic fungi co-occur with each other more or less strongly than expected based on their responses to environmental conditions and the host plant species. For this, we generated DNA-based community data on root-associated fungi of five plant species evenly sampled along an elevational gradient at a high arctic site (Zackenberget Valley, Northeast Greenland). We hypothesized that co-occurrence patterns among arctic root-associated fungi would be predominantly positive, reflecting facilitative interactions. In particular, we expected positive co-occurrences to be particularly prevalent between mycorrhizal and endophytic fungi, reflecting facilitative interactions as those previously shown for mycorrhizal fungi and bacteria (Frey-Klett et al., 2011; Johansson et al., 2004).

## 2 | MATERIALS AND METHODS

The DNA sequence data used in this paper were originally published by Abarenkov, Somervuo, Nilsson, et al. (2018a), in Abarenkov, Somervuo, Nilsson, et al. (2018b). Abarenkov, Somervuo, Nilsson, Kirk et al. (2018b) used the data solely for methodological developments related to molecular species identification, and did not provide a detailed description of the sampling design or environmental covariates. We therefore describe these aspects of the sampling design in full detail.

### 2.1 | Study design

Sampling was carried out on the western hillside of the Aucella Mountain located in the Zackenberg Valley in northeast Greenland (74°30'N, 21°00'W). The Zackenberg Valley is characterized by an arctic climate, with mean monthly temperatures ranging from -20 to +7°C and an annual precipitation of 260 mm. The vegetation consists of low tundra, dominated by arctic willow (*Salix arctica*) and arctic bell-heather (*Cassiope tetragona*). The western hillside of the Aucella Mountain spans from 0 to 1,040 m above sea level (a.s.l.).

We selected five plant species which occurred along the whole elevational gradient and which are known to establish mycorrhizal and endophytic fungal associations: (a) alpine bistort (*Bistorta vivipara*) associates with ectomycorrhizal fungi (Gardes & Dahlberg, 1996); (b) mountain avens (the crossbreed *Dryas octopetala* × *integrifolia*) associates with ectomycorrhizal fungi (Gardes & Dahlberg, 1996); (c) purple saxifrage (*Saxifraga oppositifolia*) forms associations with both ectomycorrhizal fungi and arbuscular mycorrhizal fungi (Fujimura & Egger, 2012); (d) arctic willow associates with ectomycorrhizal fungi (Gardes & Dahlberg, 1996); and (e) moss campion (*Silene acaulis*) forms associations with both ericoid mycorrhizal and ectomycorrhizal fungi (Kohn & Stasovski, 1990; Read & Haselwandter, 1981). Endophytic fungi are expected to be found in all plant species (Strobel, 2018).

In July 2015, the plant individuals were sampled at 18 locations randomly distributed along the elevational gradient. The lowest sampling location was located at 33 m a. s. l. and the highest sampling location at 479 m a. s. l. which was the upper limit for the distributions of the focal plants. The sampling locations were separated by an average distance of 2.7 km, with the shortest distance between two sampling locations being 373 m and the longest 6.4 km. Within each sampling location, we uprooted five plant individuals from each of the focal species along a transect of at most 50 m. The plant individuals were selected by moving from a starting point towards the southeast along an elevational isocline. Sampled individuals of different species were sampled at any distance, but individuals of the same species were separated by at least 1 m from each other. The whole root system of each plant individual was uprooted, and the fine roots (<2 mm) were collected as samples for the forthcoming molecular analyses, which were initiated in September 2015. The fine root samples were hand-cleaned for soil particles, first in the field and

then more thoroughly in the laboratory. Within the next 2 days after the first root cleaning in the field, the roots were further cleaned in the laboratory once or twice, where the absence of soil particles was verified with the use of a magnifying lens. The root samples were stored by wrapping them in tissue paper and placing them in plastic bags containing moisture-indicating silica gel. The following environmental variables, expected to shape niches of plants and/or fungi (e.g., Erlandson, Savage, Cavender-Bares, & Peay, 2015), were directly measured at each sampling location: soil pH (in soil-water suspension using a Direct Soil Measurement pH Portable Meter, Hanna Instruments), soil water content (using a HydroSense Handheld Soil Moisture Sensor, Campbell Scientific), the depth of the active soil layer (measuring the distance until the frozen horizon with a metal bar) and vegetation cover (visually estimated cover percentage of all vascular plants in an 1 × 1-m area). These measurements were averaged over three points for each sampling location.

### 2.2 | DNA extraction and sequencing

The root samples were first weighed and then ground into fine powder using a ball mill (Retsch Mixer Mill MM400). Ten milligrams of each sample was used for DNA extraction (all available material was used if less than 10 mg). DNA was extracted using the NucleoSpin Plant II kit (Macherey-Nagel).

As fungal DNA marker, we used the internal transcribed spacer (ITS). The ITS region shows marked variation among fungal species but less variation within species and it is thus considered the universal genetic barcode for fungi (Schoch et al., 2012). We amplified the internal transcribed spacer region 2 (ITS2) using the forward primer fITS7 (GTGARTCATCGAATCTTTG) (Ihrmark et al., 2012) and the reverse primer ITS4 (TCCTCCGCTTATTGATATGC) (White, Bruns, Lee, & Taylor, 1990). We note, however, that the ITS2 is not an optimal marker for discriminating among arbuscular mycorrhizal taxa (Krüger, Krüger, Walker, Stockinger, & Schüßler, 2012), and thus this group of root-associated fungi is likely to be underrepresented in our data.

Before polymerase chain reaction (PCR), all the DNA extracts were diluted to 0.5 ng/μl with ddH<sub>2</sub>O based on NanoDrop Lite (Thermo Scientific) measurements. PCR amplifications were performed using primers fITS7 and ITS4 tagged with 104 unique identification tags following Clemmensen, Ihrmark, Durling, and Lindahl (2016). PCRs were run for 22–35 cycles in a total volume of 50 μl. The number of cycles was adjusted on a sample-by-sample basis to yield bands of approximately the same strength for all samples on the agarose gel. PCR products were cleaned using the AMPure kit (Beckman Coulter), and DNA concentrations were determined using the Qubit dsDNA HS Assay Kit (Life Technologies). All 450 samples were successfully amplified and pooled into six composite samples, which were further purified using the Cycle-Pure Kit (Omega), and verified for quality on a Bioanalyzer (Agilent Tech). Pooled amplicon mixes were sequenced on a PacBio RS II system (Pacific Biosciences) at SciLifeLab (Uppsala, Sweden) using six SMRT cells. The sequence

data have been published in the Dryad data repository (Abarenkov, Somervuo, Nilsson, et al., 2018a).

### 2.3 | Bioinformatics analyses

We clustered the sequences to operational taxonomic units (OTUs) with a 98.5% clustering threshold using the SCATA pipeline (Sequence Clustering and Analysis of Tagged Amplicons, <http://scata.mykopat.slu.se>). Sequences were screened for tags and primer sequences, requiring a 90% match to the primer sequence. Sequences with a mean quality score lower than 20 or containing bases with a score lower than 3 were discarded. The remaining sequences were aligned pairwise, using USEARCH, and clustered into OTUs by single linkage clustering with 1.5% maximum distance allowed for sequences to enter clusters, homopolymers collapsed to 3 bp, miss match penalty 1, gap open penalty 0 and gap extension 1. Remaining singletons were removed. Species identification of the OTUs was conducted using the probabilistic taxonomic placement method of Protax-Fungi (Abarenkov, Somervuo, Nilsson, Kirk, et al., 2018b). This method quantifies the probabilities of all possible taxonomic placements for each query sequence. Importantly, the identification probabilities given by Protax-Fungi account for the possibility that some of the reference sequences are mislabelled, or that the species behind the environmental sequences are missing from the taxonomy or that there are no reference sequences for them. Such sources of uncertainty are particularly common among data on the fungal kingdom (Somervuo et al., 2017). As parameterization data, Protax-Fungi used a fungal taxonomy derived from Index Fungorum (Index Fungorum, 2016), and a fungal reference database derived from UNITE (Kõljalg et al., 2013).

Identified fungal taxa were classified as mycorrhizal or endophytic based on the FUNguild database (Nguyen et al., 2016) as well as the expertise of the authors (N.A. and B.L.). Mycorrhizal species were mostly ectomycorrhizal, but also some ericoid and arbuscular mycorrhizal fungal species were included. Fungal species with uncertain interaction ecology, for which different data sources provided contrasting information or that could not be taxonomically assigned (i.e., no hit), were grouped as “unclassified.”

The molecular data included 2,874 OTUs, out of which 695 were classified as mycorrhizal and 659 as endophytic fungi. Because our interest here was to estimate species associations, OTUs occurring in less than 5% of the sampling units were excluded from the statistical analyses, leaving 231 OTUs (78 mycorrhizal and 63 endophytic, see Table S1).

### 2.4 | Statistical analyses

To estimate the pairwise co-occurrences among the root-associated fungi, we used the joint species distribution modelling approach (Warton et al., 2015) of Hierarchical Modelling of Species Communities (HMSC; Ovaskainen et al., 2017). HMSC is a

hierarchical generalized linear mixed model and is thus structured by fixed effects and random effects. This approach allowed us to jointly model the occurrences and abundances of the fungal species identified in each plant individual as a function of the environmental variables, while accounting for the structure of the study design. Importantly for reaching the aim of this paper, this modelling approach allowed us to estimate to what extent co-occurrence could be explained by the environmental conditions, and to identify species pairs for which statistically supported co-occurrence remained also after accounting for the environmental conditions and the host plant species. This is achieved by comparing the co-occurrences estimated from a model which does not include any explanatory variables (i.e., raw co-occurrences) with the co-occurrences estimated from a model including explanatory variables (i.e., residual co-occurrences). Residual co-occurrences are more likely to reflect interactive associations than raw co-occurrences (Ovaskainen, Abrego, Halme, & Dunson, 2016; Ovaskainen et al., 2010; Zurell, Pollock, & Thuiller, 2018). The structure of the two fitted models is as follows:

1. “Raw co-occurrences model”—In this model, we accounted only for the variable of sequencing depth (continuous variable, log-transformed), which represents observation effort rather than variation in environmental conditions.
2. “Residual co-occurrences model”—In this model, in addition to sequencing depth, we included the plant species (categorical variable), elevation (continuous variable), the squared term of elevation (to account for the nonlinear responses of root-associated fungi to altitude; see Miyamoto, Nakano, Hattori, & Nara, 2014), soil pH (continuous variable), soil water content (continuous variable), depth of the active soil layer (continuous variable) and vegetation cover (continuous variable) as explanatory variables. We further included the community-level random effect (implemented through latent variables following Ovaskainen et al., 2016) of the sampling site to control for additional site-level variation that was not accounted for by the above-mentioned variables.

While in both models the random effect of the plant individual models the species pair that co-occurs more or less often than expected by random, the two models differ in what “by random” exactly means, as this depends on which fixed effects are controlled for in the models. Thus, in the raw co-occurrences model, “by random” refers to the null expectation that root-associated species would be distributed among the plant individuals fully independently of each other. In the residual co-occurrences model, “by random” refers to the environmentally constrained null expectation that, after accounting for the species affinities to the environmental covariates and host plant species, the root-associated species would be distributed among the plant individuals independently of each other.

To account for the zero-inflated nature of the data, we fitted hurdle-type models. For this, we first modelled presence-absences using a probit-link function, and then abundances (sequence counts) conditional on presence with a log-normal model. We fitted the models using the R-package HMSC 3.0 (Tikhonov et al., 2020) assuming the

default prior distributions. The settings applied for posterior sampling through a Markov chain Monte Carlo (MCMC) method and the results of MCMC convergence are provided in the Appendix S1.

We considered that a co-occurrence between a pair of fungal species was well supported if the posterior probability of the association being positive or negative was at least 0.9. To assess how different types of fungi varied in the proportion of positive and negative co-occurrences that they established, we computed the proportions of species pairs that showed well-supported positive or negative co-occurrences within and among the three groups of root-associated fungi.


To evaluate the phylogenetic structure in the co-occurrence patterns, we focused on the taxonomic composition of the groups of positively co-occurring species revealed by the residual co-occurrences model (Figure 1b; Table S1). We evaluated the differences in phylogenetic composition in two ways. We first tested whether higher taxonomic levels (families in the case of mycorrhizal species and orders in the case of endophytic fungi because for the latter the family-level taxonomy is unresolved for many species) were under- or over-represented among OTUs that show statistically supported co-occurrences. To make the comparison possible, we conducted this analysis for those taxonomic levels that were represented by at least five species. For each family, we counted how many species were included in the four groups, and compared this number with the null expectation based on 10,000 permutations. In our second phylogenetic analysis, we asked whether the groups of positively co-occurring species differed from each other in their phylogenetic composition. For this, we counted the number of species in each family or order, and computed Bray–Curtis dissimilarity to evaluate how different the groups of positively co-occurring species were in

terms of phylogenetic composition. We compared this dissimilarity to the null expectation based on 10,000 permutations.

To evaluate the environmental drivers that cause the difference between the raw and the residual co-occurrences, we quantified how much of the community variation explained by the residual co-occurrence model was attributed to the plant species, elevation, soil pH, soil water content, depth of the active soil layer and vegetation cover, using the variance partitioning approach of Ovaskainen et al. (2017). We evaluated the explanatory power of the presence–absence model by the area under the curve (AUC) statistic and that of the abundance model by the  $R^2$  value, both averaged over the species (Norberg et al., 2019).


### 3 | RESULTS

The number of statistically supported positive raw co-occurrences was higher than the number of statistically supported negative raw co-occurrences (Figures 1 and 2). Taking the effects of the environmental covariates into account, the majority of the positive co-occurrences remained, whereas the majority of the negative co-occurrences disappeared (Figures 1 and 2). Thus, in line with our original hypothesis, we found the root-associated fungal communities of our high-arctic site to be predominantly structured by positive co-occurrences. The co-occurrences that disappeared between the “raw co-occurrences model” and the “residual co-occurrences model” were mostly attributed to plant species and elevation, the other environmental variables explaining a small part of the community variation (Table 1). The “residual co-occurrences model” had a high explanatory power, with an AUC value of .86 for the presence–absence model and  $R^2$  of .30 for the abundance model.



**FIGURE 1** Raw and residual co-occurrences among root-associated fungal OTU pairs estimated from the presence–absence model. The co-occurrences are shown only for cases for which the co-occurrence was either positive (red) or negative (blue) with at least 90% posterior probability (the remaining cases are indicated in white). The OTUs have been ordered emphasizing the network structure, so that OTUs within group 1 (including the mycorrhizal group M1, the endophyte group E1 and the unclassified group U1) are found especially often together, the OTUs in group 2 (including M2, E2 and U2) are found especially often together, and the OTUs of group 1 are found especially seldom with the OTUs of group 2. The identity of the OTUs displayed in the panels is given in Table S1, based on identification through Protax-Fungi [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

**FIGURE 2** Proportions of species pairs showing well-supported (90% posterior probability) associations among the different groups of root-associated fungi (mycorrhizal, endophytic and unclassified fungi). Results from both the presence-absence model and the abundance conditional on presence model are shown. Proportions of species pairs with a well-supported positive association are indicated in red and proportions of species pairs with a well-supported negative association are indicated in blue [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**TABLE 1** Variance partitioning among the fixed and random effects included in the “residual co-occurrences model”

	Covariate	Presence-absence (%)	Abundance (%)
Fixed effects	Plant species	24.5	28.2
	Elevation	16.4	12.1
	Vegetation cover	4.4	8.7
	pH	3.6	5.2
	Soil moisture	1.8	4.3
	Depth of active soil layer	1.1	3.8
	Random effects	Sampling location	12.9
Plant individual		20.7	9.6

Interestingly, most of the positive residual co-occurrences were found among mycorrhizal fungi (30%) or between mycorrhizal and endophytic fungi (25%, Figure 2). Thus, especially mycorrhizal fungi co-occurred positively with either other mycorrhizal or endophytic fungi. The first permutation test showed that no family or order was over- or under-represented among the groups of positively co-occurring species ( $p > .05$  for all families or orders, Table 2; Table S2). The second permutation test, which asked whether the groups of positively co-occurring species were phylogenetically similar, revealed that the two groups of positively co-occurring mycorrhizal species were represented by different families more distinctly than expected by random (Bray-Curtis dissimilarity = 0.90, null expectation 0.69,  $p < .001$ ), whereas there were no significant differences in the phylogenetic composition between the two groups of positively co-occurring endophytic fungi (Bray-Curtis dissimilarity = 0.76, null expectation 0.73,  $p = .45$ ).

**TABLE 2** Number of OTUs (in parentheses) of each of the families forming the positively co-occurring groups depicted in Figure 1

Group 1	Mycorrhizal (Group M1)	Sebacinaceae (4), Thelephoraceae (2), Russulaceae (2), Clavariaceae (1), Inocybaceae (1)
	Endophyte (Group E1)	Helotiales (4), Chaetothyriales (1), Archaeorhizomycetales (1)
Group 2	Mycorrhizal (Group M2)	Thelephoraceae (15), Sebacinaceae (12), Cortinariaceae (9), Inocybaceae (8), Pyronemataceae (2), Clavulinaceae (1), Gloniaceae (1), Tuberaceae (1)
	Endophyte (Group E2)	Chaetothyriales (12), Helotiales (6), Hyaloscyphaceae (3), Hypocreales (4), Capnodiales (1), Coniochaetaceae (1), Leotiales (1), Nectriaceae (1), Pleosporales (1), Sordariales (1), Thelebolaceae (1)

Note: The OTUs have been taxonomically assigned using the probabilistic method Protax-Fungi, provided in Table S1. In cases where the family-level taxonomic assignment was unknown, the order-level taxonomic assignment is given.

The vast majority of statistically supported co-occurrences were detected by the model fitted to the presence-absence data, whereas the model fitted to the abundance conditional on presence data revealed few associations (Figure 2). In other words, among those species pairs for which we found statistically supported positive co-occurrences, the presence of one of the species implied the presence of the other species. However, if considering only those

sampling units where both species are present, a high abundance of one of the species did not imply high abundance of the other species.

## 4 | DISCUSSION

In both macroscopic and microbial communities, negative interactions have, thus far, gained more attention than positive interactions. However, community ecologists now increasingly acknowledge the importance of positive interspecific interactions in shaping communities (e.g., Bruno, Stachowicz, & Bertness, 2003; Tirado, Pugnaire, & Eriksson, 2005), also among microorganisms (Elias & Banin, 2012; Freilich et al., 2011; Nadell, Xavier, & Foster, 2008; but see Foster & Bell, 2012). Our results from arctic plant root systems suggests that positive interactions play an important role in shaping microbial communities. By showing that co-occurrences among fungal species are predominantly positive, our results have important consequences for understanding the structure of arctic interaction networks, where associations with symbiotic fungi play a pivotal role in plant nutrient acquisition (Hobbie & Hobbie, 2006).

In line with our main hypothesis, root-associated fungi formed predominantly positive co-occurrences that were particularly prevalent between mycorrhizal fungi and other root-associated fungi. Some previous studies have that found predominantly positive co-occurrences among root-associated fungi have suggested that such patterns reflect facilitative interactions (Pan & May, 2009; Yamamoto et al., 2014). This possibility seems plausible also in the present study, especially given that mycorrhizal fungi are known to establish facilitative interactions with bacteria (Frey-Klett et al., 2011; Johansson et al., 2004). Some bacteria enhance fungal growth through improved nutrition and reduction of the impact of adversities (Frey-Klett et al., 2007). We propose that in the same way as mycorrhizal fungi exhibit facilitative interactions with bacteria, they may also form facilitative interactions with other mycorrhizal or endophytic fungi. Just as similar associations between the mycorrhizal helper bacteria and mycorrhizal fungi have been found not to be restricted to particular bacterial or fungal taxa (Frey-Klett et al., 2007), we found no phylogenetic structure among fungi–fungi co-occurrence patterns. As one example of facilitative interactions, some fungi may change the host plant's physiology in a way stimulating or benefitting colonization by other fungi (Kennedy et al., 2014). Another possibility is that rather than facilitative interactions, the positive co-occurrences detected in our study reflect either myco-parasitic or trophic interactions among pairs of root-associated fungi (Lindahl et al., 2010; Tedersoo et al., 2009). Finally, it might be that positive co-occurrences among fungi arise due to their shared helper bacteria, so that the presence of some fungi and their associated helper bacteria may make it easier for other fungi to colonize the plant.

Most studies analysing observational data have concluded that root-associated fungi mainly compete during colonization of plant roots (e.g., Koide et al., 2005; Pickles et al., 2012; Pickles et al., 2010; Wubet et al., 2012). However, studies that have reported such

negative co-occurrences have typically been based on samples that do not represent the roots of single plant individuals, and that have applied co-occurrence analyses without accounting for the effects of environmental variation or spatial structure in the data. Compared to these previous studies, our sampling design and joint species distribution modelling approach allowed us to unveil patterns of co-occurrences that were not explained by the environment. These co-occurrences were scored at the level of the roots of a plant individual (i.e., at the scale at which fungi–fungi interactions are expected to primarily take place). We found that negative co-occurrences were mostly explained by environmental variation, whereas most positive co-occurrences remained unexplained by environmental variation. However, we note that because our approach of estimating species interactions relies on co-occurrence patterns rather than experimental validation, it is highly susceptible to the choice of environmental covariates that are controlled for in the analysis (Dormann et al., 2018). For instance, our models do not include microscale environmental data (e.g., soil conditions changing at the scale of 1 m or less), which may have explained part of the variation in species occurrences that remained unexplained in our data. While a statistical model will always miss at least some of the factors behind the data, given the high explanatory power achieved by our models, we argue that the general patterns captured by our analyses would not drastically change even if we had accounted also for those “unknown” environmental constraints.

We note that one reason why we captured more positive than negative associations may be that co-occurrence data are typically more informative about positive associations than about negative associations. Namely, if two species are rare in the sense that they inhabit only a small fraction of sampling units, it is unlikely that just by chance they would co-occur in the very same sampling units, and thus the data may contain a strong signal on positive co-occurrence of such rare species. While this may partially explain why we recorded a higher fraction of positive than negative co-occurrences in the raw co-occurrence model, we do not consider that this is the case for the residual co-occurrence model. This is because most of the positive co-occurrences remained statistically supported after environmental variation was taken into account, even if the environmental variables explained a large part of the variation in species occurrences. Hence, we consider the dominance of residual positive co-occurrences to reflect ecological signal rather than a statistical artefact.

Previous studies have shown that root-associated fungal communities show marked variation along elevational gradients, due to the changes in vegetation structure and abiotic conditions (Bahram, Pölme, Kõljalg, Zarre, & Tedersoo, 2012; Coince et al., 2014; Jarvis, Woodward, & Taylor, 2015; Matsuoka, Mori, Kawaguchi, Hobara, & Osono, 2016). In line with these results, we found that plant species and elevation were major factors explaining fungal occurrences. The effect of elevation is likely to reflect the effect of temperature, with high-elevation sites being colder than low-elevation sites (Bahram et al., 2012; Jarvis et al., 2015). The effect of plant species reflects the specialization of root-associated fungal communities to the host

plant species (Pölme et al., 2018). One environmental aspect that was not taken into account in our study is differences in microhabitat between mycorrhizal and endophytic fungi (i.e., that the two groups colonize different tissues in the roots). Such differences have been previously suggested to explain their positive co-occurrences within root systems (Wagg et al., 2008). Yet, even if mycorrhizal and endophytic fungi colonize different parts of the root, the chemical changes caused by a given fungus are likely to cause systemic rather than local effects, affecting the physiology of the whole plant individual (Bonfante & Genre, 2010).

Given our intriguing finding of widespread positive co-occurrences between root-associated fungi from a high-arctic site, we propose that future studies should focus on assessing the generality of our results for arctic ecosystems and the mechanisms driving fungal co-occurrences and their significance for plant growth. What remains unresolved is whether the patterns found in our study extend to other fungal groups inhabiting other plant growth forms than those considered here. In our study, the focal plant species establish mostly ectomycorrhizal associations, and thus plants establishing arbuscular associations such as herbaceous plants were not studied. Also, effectively all fungal species classified as mycorrhizal were ectomycorrhizal, because to efficiently detect arbuscular mycorrhizal fungi, a different set of primers is needed (Lekberg et al., 2018). Studies focusing on the mechanisms driving co-occurrence patterns will be particularly relevant for arctic ecosystems, where plant growth is hampered by the short growing season and the scarcity of nutrients. At a more applied level, added proof for the ecological significance of fungus–fungus co-occurrences might have important implications in biotechnology and soil restoration. Some studies have demonstrated that co-inoculation of some specific pairs of mycorrhizal and endophytic fungal species is more beneficial for plant growth than single inoculations (Reininger & Sieber, 2012, 2013). Our study provides a starting point for investigating fungal species pairs with beneficial effects on plant fitness.

## ACKNOWLEDGEMENTS

We thank the three anonymous reviewers for the constructive comments on an earlier version of the manuscript. N.A. and O.O. were supported by the Research Council of Norway (CoE grant no. 223257), N.A. by the Academy of Finland (grant no. 308651), O.O. by the Academy of Finland (grant nos. 1273253 and 250444), A.T. by the Swedish Research Council Vetenskapsrådet (grant no. 2015-03993), T.R. by grants from the Ella and Georg Ehrnrooth Foundation, G.T. by the LUOVA graduate school of University of Helsinki, and T.R. and T.H. by the Academy of Finland (grant nos. 276909 and 285803). Aarhus University is thanked for providing access to the Zackenberg Research Station, and the Logistics team and the Biobasis team created an excellent working environment.

## AUTHOR CONTRIBUTIONS

A.T., T.R., O.O. and N.A. conceived the idea and designed the study. N.A. and O.O. collected the data; B.L. supervised and T.H. conducted the laboratory work; B.L. supervised and T.H. and P.S. carried

out the bioinformatic analyses; N.A., O.O. and G.T. performed the statistical analyses; N.A. wrote the first draft of the manuscript with specific contributions from T.R., T.H. and O.O., and all authors contributed substantially to the revisions.

## DATA AVAILABILITY STATEMENT

The species and environmental data matrices used in the data analysis as well as R-scripts for reproducing the results are published in the Dryad data repository, <https://doi.org/10.5061/dryad.9kd51c5dp> (Abrego, 2020). The original sequence data are published in the Dryad data repository, <https://doi.org/10.5061/dryad.9dr6j0c>.

## ORCID

Nerea Abrego  <https://orcid.org/0000-0001-6347-6127>

Tomas Roslin  <https://orcid.org/0000-0002-2957-4791>

Ayco J. M. Tack  <https://orcid.org/0000-0002-3550-1070>

Björn D. Lindahl  <https://orcid.org/0000-0002-3384-4547>

Gleb Tikhonov  <https://orcid.org/0000-0003-3040-0307>

Panu Somervuo  <https://orcid.org/0000-0003-3121-4047>

Niels Martin Schmidt  <https://orcid.org/0000-0002-4166-6218>

Otso Ovaskainen  <https://orcid.org/0000-0001-9750-4421>

## REFERENCES

- Abarenkov, K., Somervuo, P., Nilsson, R. H., Kirk, P. M., Huotari, T., Abrego, N., & Ovaskainen, O. (2018a). Protax-fungi: A web-based tool for probabilistic taxonomic placement of fungal internal transcribed spacer sequences. *New Phytologist*, 220, 517–525. <https://doi.org/10.1111/nph.15301>
- Abarenkov, K., Somervuo, P., Nilsson, R., Kirk, P. M., Huotari, T., Abrego, N., & Ovaskainen, O. (2018b). Data from: PROTAX-fungi: A web-based tool for probabilistic taxonomic placement of fungal ITS sequences. Dryad. <https://doi.org/10.5061/dryad.9dr6j0c>
- Abrego, N. (2020). Accounting for environmental variation in co-occurrence modeling reveals the importance of positive interactions in root-associated fungal communities. Dryad Dataset, <https://doi.org/10.5061/dryad.9kd51c5dp>
- Artursson, V., Finlay, R. D., & Jansson, J. K. (2006). Interactions between arbuscular mycorrhizal fungi and bacteria and their potential for stimulating plant growth. *Environmental Microbiology*, 8, 1–10. <https://doi.org/10.1111/j.1462-2920.2005.00942.x>
- Bahram, M., Pölme, S., Kõljalg, U., Zarre, S., & Tedersoo, L. (2012). Regional and local patterns of ectomycorrhizal fungal diversity and community structure along an altitudinal gradient in the Hyrcanian forests of northern Iran. *New Phytologist*, 193, 465–473. <https://doi.org/10.1111/j.1469-8137.2011.03927.x>
- Bairey, E., Kelsic, E. D., & Kishony, R. (2016). High-order species interactions shape ecosystem diversity. *Nature Communications*, 7, 12285. <https://doi.org/10.1038/ncomms12285>
- Bonfante, P., & Genre, A. (2010). Mechanisms underlying beneficial plant–fungus interactions in mycorrhizal symbiosis. *Nature Communications*, 1, 48. <https://doi.org/10.1038/ncomms1046>
- Bruno, J. F., Stachowicz, J. J., & Bertness, M. D. (2003). Inclusion of facilitation into ecological theory. *Trends in Ecology & Evolution*, 18, 119–125. [https://doi.org/10.1016/S0169-5347\(02\)00045-9](https://doi.org/10.1016/S0169-5347(02)00045-9)
- Clemmensen, K. E., Ihrmark, K., Durling, M. B., & Lindahl, B. D. (2016). Sample preparation for fungal community analysis by high-throughput sequencing of barcode amplicons. In F. Martin, & S. Uroz (Eds.), *Microbial Environmental Genomics (MEG)* (pp. 61–88). New York, NY: Springer.
- Coince, A., Cordier, T., Lengellé, J., Defosse, E., Vacher, C., Robin, C., ... Marçais, B. (2014). Leaf and root-associated fungal assemblages



- do not follow similar elevational diversity patterns. *PLoS One*, 9, e100668. <https://doi.org/10.1371/journal.pone.0100668>
- Dormann, C. F., Bobrowski, M., Dehling, D. M., Harris, D. J., Hartig, F., Lischke, H., ... Kraan, C. (2018). Biotic interactions in species distribution modelling: 10 questions to guide interpretation and avoid false conclusions. *Global Ecology and Biogeography*, 27, 1004–1016. <https://doi.org/10.1111/geb.12759>
- Elias, S., & Banin, E. (2012). Multi-species biofilms: Living with friendly neighbors. *FEMS Microbiology Reviews*, 36, 990–1004. <https://doi.org/10.1111/j.1574-6976.2012.00325.x>
- Erland, S., & Finlay, R. (1992). Effects of temperature and incubation time on the ability of three ectomycorrhizal fungi to colonize *Pinus sylvestris* roots. *Mycological Research*, 96, 270–272. [https://doi.org/10.1016/S0953-7562\(09\)80936-2](https://doi.org/10.1016/S0953-7562(09)80936-2)
- Erlandson, S. R., Savage, J. A., Cavender-Bares, J. M., & Peay, K. G. (2015). Soil moisture and chemistry influence diversity of ectomycorrhizal fungal communities associating with willow along an hydrologic gradient. *FEMS Microbiology Ecology*, 92, fiv148. <https://doi.org/10.1093/femsec/fiv148>
- Foster, K., & Bell, T. (2012). Competition, not cooperation, dominates interactions among culturable microbial species. *Current Biology*, 22, 1845–1850. <https://doi.org/10.1016/j.cub.2012.08.005>
- Founoune, H., Duponnois, R., Bâ, A. M., Sall, S., Branget, I., Lorquin, J., ... Chotte, J. L. (2002). Mycorrhiza Helper Bacteria stimulate ectomycorrhizal symbiosis of *Acacia holosericea* with *Pisolithus alba*. *New Phytologist*, 153, 81–89. <https://doi.org/10.1046/j.0028-646X.2001.00284.x>
- Freilich, S., Zarecki, R., Eilam, O., Segal, E. S., Henry, C. S., Kupiec, M., ... Ruppin, E. (2011). Competitive and cooperative metabolic interactions in bacterial communities. *Nature Communications*, 2, 589. <https://doi.org/10.1038/ncomms1597>
- Frey-Klett, P., Burlinson, P., Deveau, A., Barret, M., Tarkka, M., & Sarniguet, A. (2011). Bacterial-fungal interactions: Hyphens between agricultural, clinical, environmental, and food microbiologists. *Microbiology and Molecular Biology Reviews*, 75, 583–609. <https://doi.org/10.1128/MMBR.00020-11>
- Frey-Klett, P., Garbaye, J., & Tarkka, M. (2007). The mycorrhiza helper bacteria revisited. *New Phytologist*, 176, 22–36. <https://doi.org/10.1111/j.1469-8137.2007.02191.x>
- Fujimura, K. E., & Egger, K. N. (2012). Host plant and environment influence community assembly of High Arctic root-associated fungal communities. *Fungal Ecology*, 5, 409–418. <https://doi.org/10.1016/j.funeco.2011.12.010>
- Gardes, M., & Dahlberg, A. (1996). Mycorrhizal diversity in arctic and alpine tundra: An open question. *New Phytologist*, 133, 147–157. <https://doi.org/10.1111/j.1469-8137.1996.tb04350.x>
- Gorzalak, M. A., Hambleton, S., & Massicotte, H. B. (2012). Community structure of ericoid mycorrhizas and root-associated fungi of *Vaccinium membranaceum* across an elevation gradient in the Canadian Rocky Mountains. *Fungal Ecology*, 5, 36–45. <https://doi.org/10.1016/j.funeco.2011.08.008>
- Gotelli, N. J. (2000). Null model analysis of species co-occurrence patterns. *Ecology*, 81, 2606–2621. [https://doi.org/10.1890/0012-9658\(2000\)081\[2606:NMAOSC\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2000)081[2606:NMAOSC]2.0.CO;2)
- Gould, A. L., Zhang, V., Lamberti, L., Jones, E. W., Obadia, B., Korasidis, N., ... Ludington, W. B. (2018). Microbiome interactions shape host fitness. *Proceedings of the National Academy of Sciences of the United States of America*, 115, E11951–E11960. <https://doi.org/10.1073/pnas.1809349115>
- Hobbie, J. E., & Hobbie, E. A. (2006). <sup>15</sup>N in symbiotic fungi and plants estimates nitrogen and carbon flux rates in Arctic tundra. *Ecology*, 87, 816–822. [https://doi.org/10.1890/0012-9658\(2006\)87\[816:NISFA P\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[816:NISFA P]2.0.CO;2)
- Ihrmark, K., Bödeker, I. T. M., Cruz-Martinez, K., Friberg, H., Kubartova, A., Schenck, J., ... Lindahl, B. D. (2012). New primers to amplify the fungal ITS2 region – evaluation by 454-sequencing of artificial and natural communities. *FEMS Microbiology Ecology*, 82, 666–677. <https://doi.org/10.1111/j.1574-6941.2012.01437.x>
- Index Fungorum (2016). 2016. <http://www.indexfungorum.org>
- Ingham, E. R., & Molina, R. (1991). Interactions among mycorrhizal fungi, rhizosphere organisms and plants. In P. Barbosa, V. A. Krischik, & C. J. Jones (Eds.), *Microbial mediation of plant-herbivore interactions* (1st ed., pp. 169–199). Hoboken, NJ: John Wiley & Sons.
- Jarvis, S. G., Woodward, S., & Taylor, A. F. S. (2015). Strong altitudinal partitioning in the distributions of ectomycorrhizal fungi along a short (300 m) elevation gradient. *New Phytologist*, 206, 1145–1155. <https://doi.org/10.1111/nph.13315>
- Johansson, J. F., Paul, L. R., & Finlay, R. D. (2004). Microbial interactions in the mycorrhizosphere and their significance for sustainable agriculture. *FEMS Microbiology Ecology*, 48, 1–13. <https://doi.org/10.1016/j.femsec.2003.11.012>
- Johnson, N. C., Graham, J., & Smith, F. A. (1997). Functioning of mycorrhizal associations along the mutualism–parasitism continuum. *New Phytologist*, 135, 575–585. <https://doi.org/10.1046/j.1469-8137.1997.00729.x>
- Kennedy, P. G., & Bruns, T. D. (2005). Priority effects determine the outcome of ectomycorrhizal competition between two Rhizopogon species colonizing *Pinus muricata* seedlings. *New Phytologist*, 166, 631–638. <https://doi.org/10.1111/j.1469-8137.2005.01355.x>
- Kennedy, P., Nguyen, N., Cohen, H., & Peay, K. (2014). Missing checkerboards? An absence of competitive signal in *Alnus*-associated ectomycorrhizal fungal communities. *PeerJ*, 2, e686. <https://doi.org/10.7717/peerj.686>
- Kennedy, P. G., Peay, K. G., & Bruns, T. D. (2009). Root tip competition among ectomycorrhizal fungi: Are priority effects a rule or an exception? *Ecology*, 90, 2098–2107. <https://doi.org/10.1890/08-1291.1>
- Kohn, L. M., & Stasovski, E. (1990). The mycorrhizal status of plants at Alexandra Fiord, Ellesmere island, Canada, a high arctic site. *Mycologia*, 82, 23–35. <https://doi.org/10.1080/00275514.1990.12025836>
- Koide, R. T., Xu, B., Sharda, J., Lekberg, Y., & Ostiguy, N. (2005). Evidence of species interactions within an ectomycorrhizal fungal community. *The New Phytologist*, 165, 305–316. <https://doi.org/10.1111/j.1469-8137.2004.01216.x>
- Köljal, U., Nilsson, R. H., Abarenkov, K., Tedersoo, L., Taylor, A. F. S., Bahram, M., ... Larsson, K.-H. (2013). Towards a unified paradigm for sequence-based identification of fungi. *Molecular Ecology*, 22, 5271–5277. <https://doi.org/10.1111/mec.12481>
- Krüger, M., Krüger, C., Walker, C., Stockinger, H., & Schübler, A. (2012). Phylogenetic reference data for systematics and phylotaxonomy of arbuscular mycorrhizal fungi from phylum to species level. *New Phytologist*, 193, 970–984. <https://doi.org/10.1111/j.1469-8137.2011.03962.x>
- Lekberg, Y., Vasar, M., Bullington, L. S., Sepp, S.-K., Antunes, P. M., Bunn, R., ... Öpik, M. (2018). More bang for the buck? Can arbuscular mycorrhizal fungal communities be characterized adequately alongside other fungi using general fungal primers? *New Phytologist*, 220, 971–976. <https://doi.org/10.1111/nph.15035>
- Lindahl, B. D., de Boer, W., & Finlay, R. D. (2010). Disruption of root carbon transport into forest humus stimulates fungal opportunists at the expense of mycorrhizal fungi. *ISME Journal*, 4, 872–881. <https://doi.org/10.1038/ismej.2010.19>
- Mahmood, S., Finlay, R. D., Fransson, A., & Wallander, H. (2003). Effects of hardened wood ash on microbial activity, plant growth and nutrient uptake by ectomycorrhizal spruce seedlings. *FEMS Microbiology Ecology*, 43, 121–131. <https://doi.org/10.1111/j.1574-6941.2003.tb01051.x>
- Matsuoka, S., Mori, A. S., Kawaguchi, E., Hobara, S., & Osono, T. (2016). Disentangling the relative importance of host tree community, abiotic environment and spatial factors on ectomycorrhizal fungal

- assemblages along an elevation gradient. *FEMS Microbiology Ecology*, 92(5), fiw044. <https://doi.org/10.1093/femsec/fiw044>
- Miyamoto, Y., Nakano, T., Hattori, M., & Nara, K. (2014). The mid-domain effect in ectomycorrhizal fungi: Range overlap along an elevation gradient on Mount Fuji, Japan. *ISME Journal*, 8, 1739–1746. <https://doi.org/10.1038/ismej.2014.34>
- Moeller, H. V., & Peay, K. G. (2016). Competition-function tradeoffs in ectomycorrhizal fungi. *PeerJ*, 4, e2270. <https://doi.org/10.7717/peerj.2270>
- Nadell, C. D., Xavier, J. B., & Foster, K. R. (2008). The sociobiology of biofilms. *FEMS Microbiology Reviews*, 33, 206–224. <https://doi.org/10.1111/j.1574-6976.2008.00150.x>
- Newsham, K. K. (2011). A meta-analysis of plant responses to dark septate root endophytes. *New Phytologist*, 190, 783–793. <https://doi.org/10.1111/j.1469-8137.2010.03611.x>
- Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J., ... Kennedy, P. G. (2016). FUNGUILD: An open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecology*, 20, 241–248. <https://doi.org/10.1016/j.funeco.2015.06.006>
- Norberg, A., Abrego, N., Blanchet, F. G., Adler, F. R., Anderson, B. J., Anttila, J., ... Ovaskainen, O. (2019). A comprehensive evaluation of predictive performance of 33 species distribution models at species and community levels. *Ecological Monographs*, 89, e01370. <https://doi.org/10.1002/ecm.1370>
- Ovaskainen, O., Abrego, N., Halme, P., & Dunson, D. (2016). Using latent variable models to identify large networks of species-to-species associations at different spatial scales. *Methods in Ecology and Evolution*, 7, 549–555. <https://doi.org/10.1111/2041-210X.12501>
- Ovaskainen, O., Hottola, J., & Siitonen, J. (2010). Modeling species co-occurrence by multivariate logistic regression generates new hypotheses on fungal interactions. *Ecology*, 91, 2514–2521. <https://doi.org/10.1890/10-0173.1>
- Ovaskainen, O., Tikhonov, G., Norberg, A., Guillaume Blanchet, F., Duan, L., Dunson, D., ... Abrego, N. (2017). How to make more out of community data? A conceptual framework and its implementation as models and software. *Ecology Letters*, 20, 561–576. <https://doi.org/10.1111/ele.12757>
- Pan, J. J., & May, G. (2009). Fungal-fungal associations affect the assembly of endophyte communities in maize (*Zea mays*). *Microbial Ecology*, 58, 668–678. <https://doi.org/10.1007/s00248-009-9543-7>
- Pickles, B. J., Genney, D. R., Anderson, I. C., & Alexander, I. J. (2012). Spatial analysis of ectomycorrhizal fungi reveals that root tip communities are structured by competitive interactions. *Molecular Ecology*, 21, 5110–5123. <https://doi.org/10.1111/j.1365-294X.2012.05739.x>
- Pickles, B. J., Genney, D. R., Potts, J. M., Lennon, J. J., Anderson, I. C., & Alexander, I. J. (2010). Spatial and temporal ecology of Scots pine ectomycorrhizas. *New Phytologist*, 186, 755–768. <https://doi.org/10.1111/j.1469-8137.2010.03204.x>
- Pölmé, S., Bahram, M., Jacquemyn, H., Kennedy, P., Kohout, P., Moora, M., ... Tedersoo, L. (2018). Host preference and network properties in biotrophic plant–fungal associations. *New Phytologist*, 217, 1230–1239. <https://doi.org/10.1111/nph.14895>
- Read, D. J., & Haselwandter, K. (1981). Observations on the mycorrhizal status of some alpine plant communities. *New Phytologist*, 88, 341–352. <https://doi.org/10.1111/j.1469-8137.1981.tb01729.x>
- Reininger, V., & Sieber, T. N. (2012). Mycorrhiza reduces adverse effects of dark septate endophytes (DSE) on growth of conifers. *PLoS One*, 7, e42865. <https://doi.org/10.1371/journal.pone.0042865>
- Reininger, V., & Sieber, T. N. (2013). Mitigation of antagonistic effects on plant growth due to root co-colonization by dark septate endophytes and ectomycorrhiza. *Environmental Microbiology Reports*, 5, 892–898. <https://doi.org/10.1111/1758-2229.12091>
- Rodríguez, R. J., White, J. F., Jr., Arnold, A. E., & Redman, R. S. (2009). Fungal endophytes: Diversity and functional roles. *New Phytologist*, 182, 314–330. <https://doi.org/10.1111/j.1469-8137.2009.02773.x>
- Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C. A., ... Schindel, D. (2012). Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences of the United States of America*, 109, 6241–6246. <https://doi.org/10.1073/pnas.1117018109>
- Singh, B. K., Millard, P., Whiteley, A. S., & Murrell, J. C. (2004). Unravelling rhizosphere–microbial interactions: Opportunities and limitations. *Trends in Microbiology*, 12, 386–393. <https://doi.org/10.1016/j.tim.2004.06.008>
- Smith, S. E., & Read, D. J. (2008). *Mycorrhizal symbiosis* (3rd ed.). London, UK: Academic Press.
- Somervuo, P., Yu, D. W., Xu, C. C. Y., Ji, Y., Hultman, J., Wirta, H., & Ovaskainen, O. (2017). Quantifying uncertainty of taxonomic placement in DNA barcoding and metabarcoding. *Methods in Ecology and Evolution*, 8, 398–407. <https://doi.org/10.1111/2041-210X.12721>
- Strobel, G. (2018). The emergence of endophytic microbes and their biological promise. *Journal of Fungi*, 4, 57. <https://doi.org/10.3390/jof4020057>
- Tedersoo, L., Pärtel, K., Jairus, T., Gates, G., Pöldmaa, K., & Tamm, H. (2009). Ascomycetes associated with ectomycorrhizas: Molecular diversity and ecology with particular reference to the Helotiales. *Environmental Microbiology*, 11, 3166–3178. <https://doi.org/10.1111/j.1462-2920.2009.02020.x>
- Tikhonov, G., Opedal, Ø. H., Abrego, N., Lehikoinen, A., de Jonge, M. M. J., Oksanen, J., & Ovaskainen, O. (2020). Joint species distribution modelling with the R-package Hmsc. *Methods in Ecology and Evolution*, 11, 442–447. <https://doi.org/10.1111/2041-210X.13345>
- Tirado, R., Pugnaire, F. I., & Eriksson, O. (2005). Community structure and positive interactions in constraining environments. *Oikos*, 111, 437–444. <https://doi.org/10.1111/j.1600-0706.2005.14094.x>
- Toro, M., Azcon, R., & Barea, J. (1997). Improvement of arbuscular mycorrhiza development by inoculation of soil with phosphate-solubilizing rhizobacteria to improve rock phosphate bioavailability ( $^{32}\text{P}$ ) and nutrient cycling. *Applied and Environmental Microbiology*, 63, 4408–4412.
- Wagg, C., Pautler, M., Massicotte, H. B., & Peterson, R. L. (2008). The co-occurrence of ectomycorrhizal, arbuscular mycorrhizal, and dark septate fungi in seedlings of four members of the Pinaceae. *Mycorrhiza*, 18, 103–110. <https://doi.org/10.1007/s00572-007-0157-y>
- Warton, D. I., Blanchet, F. G., O'Hara, R. B., Ovaskainen, O., Taskinen, S., Walker, S. C., & Hui, F. K. C. (2015). So many variables: Joint modeling in community ecology. *Trends in Ecology & Evolution*, 30, 766–779. <https://doi.org/10.1016/j.tree.2015.09.007>
- Wearn, J. A., Sutton, B. C., Morley, N. J., & Gange, A. C. (2012). Species and organ specificity of fungal endophytes in herbaceous grassland plants. *Journal of Ecology*, 100, 1085–1092. <https://doi.org/10.1111/j.1365-2745.2012.01997.x>
- Werner, E. E., & Peacor, S. D. (2003). A review of trait-mediated indirect interactions in ecological communities. *Ecology*, 84, 1083–1100. [https://doi.org/10.1890/0012-9658\(2003\)084\[1083:AROTI\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2003)084[1083:AROTI]2.0.CO;2)
- White, T. J., Bruns, T. D., Lee, S., & Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In M. A. Innis, D. H. Gelfand, J. J. Sninsky, & T. J. White (Eds.), *PCR protocols: A guide to methods and applications* (pp. 315–322). New York, NY: Academic Press.
- Wootton, J. T. (1994). The nature and consequences of indirect effects in ecological communities. *Annual Review of Ecology and Systematics*, 25, 443–466. <https://doi.org/10.1146/annurev.es.25.110194.002303>
- Wubet, T., Christ, S., Schöning, I., Boch, S., Gawlich, M., Schnabel, B., ... Buscot, F. (2012). Differences in soil fungal communities between European beech (*Fagus sylvatica* L.) dominated forests are related to soil and understory vegetation. *PLoS One*, 7, 1–14. <https://doi.org/10.1371/journal.pone.0047500>

- Yamamoto, S., Sato, H., Tanabe, A. S., Hidaka, A., Kadowaki, K., & Toju, H. (2014). Spatial segregation and aggregation of ectomycorrhizal and root-endophytic fungi in the seedlings of two *Quercus* species. *PLoS One*, 9, e96363. <https://doi.org/10.1371/journal.pone.0096363>
- Zurell, D., Pollock, L. J., & Thuiller, W. (2018). Do joint species distribution models reliably detect interspecific interactions from co-occurrence data in homogenous environments? *Ecography*, 41, 1812–1819. <https://doi.org/10.1111/ecog.03315>

**How to cite this article:** Abrego N, Roslin T, Huotari T, et al. Accounting for environmental variation in co-occurrence modelling reveals the importance of positive interactions in root-associated fungal communities. *Mol Ecol*. 2020;29:2736–2746. <https://doi.org/10.1111/mec.15516>

#### SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.